References

Adler, I. (1974). A model of contact pressure in phyllotaxis. *J. theor. Biol.* 45, 1–79.

Ataullakhanov, F.I. and Guriya, G.T. (1994). Spatial aspects of the dynamics of blood clotting - I. Hypothesis. *Biophysics* 39, 91–97.

Bär, M., Eiswirth, M., Rotermund, H.H. and Ertl, G. (1992). Solitary-wave phenomena in an excitable surface-reaction. *Phys. Rev. Lett.* 69, 945–948.

Baier, H. and Bonhoeffer, F. (1992). Axon guidence by gradients of a target-derrived component. *Science* 255, 472–475.

Bellwinkel, H.W. (1992). Naturwissenschaftliche Themen im Werk von Thomas Mann. *Naturw. Rundsch.* 45, 174–183.

Bernasconi, G.P. (1994). Reaction-diffusion model for phyllotaxis. *Physica D* 70, 90–99.

Bode, H.R. (2003). Head regeneration in hydra. *Dev. Dyn.* 226, 225–236.

Bode, P.M., Awad, T.A., Koizumi, O., Nakashima, Y., Grimmelikhuijzen, C.J.P. and Bode, H.R. (1988). Development of the two-part pattern during regeneration of the head in hydra. *Development* 102, 223–235.

Bretschneider, T., Siegert, F. and Weijer, C.J. (1995). 3–dimensional scroll waves of camp could direct cell-movement and gene-expression in *Dictyostelium* slugs. *Proc. Natl. Acad. Sci. USA* 92, 4387–4391.

Bronsvoort, W. and Klok, F. (1985). Ray tracing generalized cylinders. *ACM Transactions on Graphics* 4, 291–303 .

Broun, M., Gee, L., Reinhardt, B. and Bode, H.R. (2005). Formation of the head organizer in hydra involves the canonical wnt pathway. *Development* 132, 2907–2916.

Browne, E.N. (1909). The production of new hydrants in Hydra by insertion of small grafts. *J. Exp. Zool.* 7, 1–23.

Buikema, W.J. and Haselkorn, R. (2001). Expression of the Anabaena hetR gene from a copper-regulated promoter leads to heterocyst differentiation under repressing conditions. *PNAS* 98, 2729–2734.

Butty, A.C., Perrinjaquet, N., Petit, A., Jaquenoud, M., Segall, J.E., Hofmann, K., Zwahlen, C., Peter, M. (2002). A positive feedback loop stabilizes the guaninenucleotide exchange factor Cdc24 at sites of polarization. *EMBO J.* 21, 1565–1576.

Canales, C., Grigg, S. and Tsiantis, M. (2005). The formation and patterning of leaves: recent advances. *Planta* 221, 752–756.

Chant, J. (1999). Cell polarity in yeast. *Annu. Rev. Cell Dev. Biol.* 15, 365–391.

Charest, P.G. and Firtel, R.A. (2006). Feedback signaling controls leading-edge formation during chemotaxis. *Cur. Op. Gen. Dev* 16, 339–347.

Cohn, M.J. and Tickle, C. (1996). Limbs - a model for pattern-formation within the vertebrate body plan. *Trends in Genetics* 12, 253–257.

Comfort, A. (1951). The pigmentation of molluscan shells. *Biological Review* 26, 285–301

Cook, T.A. (1914). Curves of Life. Constable and Company, London, reprinted 1979 by Dover Publications, New York.

Cortie, M.B. (1989). Models for mollusc shell shape. *South African Journal of Science* 85, 454–460.

Coultas, L., Chawengsaksophak, K. and Rossant, J. (2005). Endothelial cells and VEGF in vascular development. *Nature* 438, 937–945.

Coxeter, H.S.M. (1961). Introduction to Geometry. J. Wiley and Sons, New York

Crick, F. (1970). Diffusion in embryogenesis. *Nature* 225, 420–422.

DeRobertis, E.M. (1995). Dismanteling the organizer. *Nature* 374, 407–408.

Devreotes, P.N. (1989). *Dictyostelium discoideum*: A model system for cell-cell interactions in Development. *Science* 245, 1054–1058.

Diaz-Benjumea, J., Gonzales Gaitan, M.A.F. and Garcia-Bellido, A. (1989). Developmental genetics of the wing vein pattern of *Drosophila*. *Genome* 31, 612–619.

Diaz-Benjumea, J., Gonzales-Gaitan, M.A.F. and Garcia-Bellido, A. (1989). Developmental genetics of the wing vein pattern of *Drosophila*. *Genome* 31, 612–619.

do Carmo, M. (1976). Differential Geometry of Curves and Surfaces. Prentice Hall, Englewood Cliffs

Ermentrout, B., Campbell, J. and Oster, G. (1986). A model for shell patterns based on neural activity. *The Veliger* 28, 369–338.

Farrar, N.R. and Spencer, G.E. (2008). Pursuing a ’turning point’ in growth cone research. *Dev. Biol.,* 318, 102–111.

Fassler, P.E. and Sander, K. (1996). Mangold, Hilde (1898-1924) and Spemanns organizer - achievement and tragedy. *Rouxs Archives Dev. Biol.* 205, 323–332.

Feininger, A. and Emerson, W.K. (1972). Shells. The Viking Press

Foley, J.D., van Dam, A., Feiner, S. and Hughes, J. (1990). Computer graphics: Principles and practice. Addison-Wesley, Reading.

Folkman, J. (1976). The vascularization of a tumor. *Sci. American* 234, May, 58–73.

Fowler, D.R., Meinhardt, H. and Prusinkiewicz, P. (1992). Modeling seashells. Proceedings of SIGGRAPH ’92. *Computer Graphics* 26, 379–387.

Gasseling, MT and Saunders, JWJr (1964). Effect of the “Posterior Necrotic Zone” on the early chick wing bud on the pattern and symmetry of limb outgrowth. *Amer. Zool.* 4, 303–304.

Gerisch, G. (1968). Cell aggregation and differentation in *Dictyostelium*. Curr. *Top. Dev. Biol.* 3, 157–232.

Gierer, A. (1977). Biological features and physical concepts of pattern formation exemplified by hydra. *Curr. Top. Dev. Biol.* 11, 17–59.

Gierer, A. (1981). Generation of biological patterns and form: Some physical, mathematical, and logical aspects. *Prog. Biophys. molec. Biol.* 37, 1–47.

Gierer, A. and Meinhardt, H. (1972). A theory of biological pattern formation. Kybernetik 12, 30–39.

Gierer, A., Berking, S., Bode, H., David, C.N., Flick, K., Hansmann, G., Schaller, H. and Trenkner, E. (1972). Regeneration of hydra from reaggregated cells. *Nature New Biology* 239, 98–101.

Giulini, A., Wang, J. and Jackson, D. (2004). Control of phyllotaxy by the cytokinin-inducible response regulator homologue ABPHYL1 1031. *Nature* 430, 1031–1034.

Glass, L. and Mackey, M.C. (1988). From clocks to chaos. Princeton University Press

Goldbeter, A. (1996). Biochemical oscillations and the cellular rhythm: The molecular bases of periodic and chaotic behaviour. Cambridge University Press.

Gordon, N.R. (1990). Seashells: A Photographic Celebration. Friedman group, New York

Gould, A., Itasaki, N. and Krumlauf, R. (1998). Initiation of rhombomeric HoxB4 expression requires induction by somites and a retinoic acid pathway. *Neuron 21, 39–51*.

Grüneberg, H. (1976). Population studies on a polymorphic prosobranch snail (*Clithon (Pictoneritina) oualaniensis* Lesson). *Phil.Transctions Royal Soc. London B* 275, 385–437.

Granero, M.I., Porati, A. and Zanacca, D. (1977). A bifurcation analysis of pattern formation in a diffusion governed morphogenetic field. *J. Math. Biology* 4, 21–27.

Grapin-Botton, A., Bonnin, M.A., Sieweke, M. and Le Douarin, N.M. (1998). Defined concentrations of a posteriorizing signal are critical for MadB/Kreisler segmental expression in the hindbrain. *Development* 125, 1173–1181.

Grieshammer, U., Minowada, G., Pisenti, J.M., Abbott, U.K. and Martin, G.R. (1996). The chick *limbless* mutation causes abnormalities in limb bud dorsal-ventral patterning: implication for apical ridge formation. *Development* 122, 3851–3861.

Gurdon, J.B., Mitchell, A. and Mahony, D. (1995). Direct and continuous assessment by cells of their position in a morphogen gradient. *Nature* 376, 520–521.

Hörstadius, S (1939). The mechanics of sea-urchin development studied by operative methods. *Biol. Rev.* 14, 132–179.

Hale, C.A., Meinhardt, H. and de Boer, P.A.J. (2001). Dynamic localization cycle of the cell division regulator MinE in *Escherichia coli*. *Embo J.* 20, 1563–1572.

Hammer, O. and Bucher, E (1999). Reaction-diffusion processes: application to the morphogenesis of ammonid ornamentation . GeoBios, 32, 841.852

Harfe, B.D., Scherz, P.J., Nissim, S., Tian, F. and McMahon, A.P. (2004). Evidence for an expansion-based temporal Shh gradient in specifying vertebrate digit identities. *CELL* 4, 517–528.

Harris, M.P., Williamson, S., Fallon, J.F., Meinhardt, H. and Prum, R.O. (2005). Molecular evidence for an activator-inhibitor mechanism in development of embryonic feather branching. *PNAS* 102, 11734–11739.

Harrison, R.G. (1918). Experiments on the development of the fore-limb of Amblystoma, a self-differentiating equipotential system. *J. exp. Zool.* 25:413–46.

Hellström, M., Phng, L.K., Hofmann, J.J., Wallgard, E., Coultas, L., Lindblom, P., Alva, J., Nilsson, A.K., Karlsson, L., Gaiano, N., Yoon, K., Rossant, J., Iruela-Arispe, M.L., Kalen, M., Gerhardt, H. and Betsholtz, C. (2007). Dll4 signalling through notch1 regulates formation of tip cells during angiogenesis. *Nature* 445, 776–780.

Herman, G.T. and Liu, W.H. (1973). The dauther of Celia, the french flag and the firing quad: Progress report on a cellular linear iterative-array simulator. *Simulation* 21, 33–41.

Hobmayer, B, Rentzsch, F, Kuhn, K, Happel, CM, Cramer von Laue, C, Snyder, P, Rothbacher, U and Holstein, TW (2000). Wnt signalling molecules act in axis formation in the diploblastic metazoan hydra. *Nature* 407, 186–189.

Hopfield, J.J. and Herz, A.V.M. (1995). Rapid local synchronization of action potentials: Toward computation with coupled integrate-and-fire neurons. *Proc. Natl. Acad. Sci. USA* 92, 6655–6622.

Hu, Z., and Lutkenhaus, J. (1999). Topological regulation of cell division in Escherichia coli involves rapid pole to pole oscillation of the division inhibitor MinC under the control of MinD and MinE. *Mol Microbiol.* 34, 82–90.

Huang, X., Dong, Y. and Zhao, J. (2004). HetR homodimer is a DNA-binding protein required for heterocyst differentiation, and the DNA-binding activity is inhibited by PatS. *PNAS* 2004 101, 4848–4853.

Illert, C. (1989). Formulation and solution of the classical seashell problem. *Il Nuovo Cimento* 11 D, 761–780 .

Izpisua-Belmonte, J.C., De Robertis, E.M., Storey, K.G., Stern, C., (1993). The homeobox gene *goosecoid* and the origin of organizer cells in the early chick blastoderm. *Cell*, 74, 645–659.

Jones, C.M., Arnes, N. and Smith, J.C. (1996). Short-range signalling by Xnr-2 and BMP-4 contrasts with the long-range effects of activin *Current Biology* 6, 1468–1475.

Jung, H.S., Francis-West, P.H., Widelitz, R.B., Jiang, T.X., Ting-Berreth, S., Tickle, C., Wolpert, L. and Chuong, C.M. (1998). Local inhibitory action of *BMP*s and their relationships with activators in feather formation: implications for periodic patterning. *Dev. Biol.* 196, 11–23.

Kaandorp, J. (1994). Fractal modelling: Growth and form in biology. Springer-Verlag, Heidelberg

Kawaguchi, Y. (1982). A morphological study of the form of nature. *Computer Graphics* 16, 223–232.

Kay, R.R., Langridge, P., Traynor, D. and Hoeller, O. (2008). Changing directions in the study of chemotaxis. *Nat. Rev. Mol. Cell Biol.,* 9455–4, 3.

Khaner, O. and Eyal-Giladi, H. (1989). The chick's marginal zone and primitive streak formation. I. Coordinative effect of induction and inhibition. *Dev. Biol.* 134, 206–214.

Kiecker, C. and Lumsden, A. (2005). Compartments and their boundaries in vertebrate brain development. *Nature Rev. Neurosci.* 6, 553–564.

Koch, A.J. and Meinhardt, H. (1994). Biological pattern-formation - from basic mechanisms to complex structures. *Rev. Modern Physics* 66, 1481–1507.

Kuhlemeier, C. (2007). Phyllotaxis. *Trends Plant Sci* 12, 143–150.

Kusch, I. and Markus, M. (1996). Mollusc shell pigmentation: Cellular automaton simulations and evidence for undecidability. *J. theor. Biol.* 178, 333–340.

Ladher, R., Mohun, T.J., Smith, J.C. and Snape, A.M. (1996). Xom - a *Xenopus* homeobox gene that mediates the early effects of BMP-4. *Development* 122, 2385–2394.

Lawrence, P.A., (1992). The making of a fly: The genetics of animal design Blackwell Scientific Publications, Oxford

Lemaire,L., Roeser,T., Izpisua-Belmonte, J.C. and Kessel, M. (1997). Segregating expression domains of two *goosecoid* genes during the transition from gastrulation to neurulation in chick embryos. (Development, in press)

Lenhoff, H. (1991). Ethel Browne, Hans Spemann, and the discovery of the organizer phemenon. *Biol. Bull.* 181, 72–80.

Leptin, M (1991). *twist* and *snail* as positive and negative regulators during *Drosophila* mesoderm development. *Genes Dev.* 5, 1568–1576.

Lindsday, D.T. (1982). A new programmatic basis for shell pigment patterns in the bivalue mollusc *Lioconcha castrensis* L. *Differentiation* 21, 32–36.

Lovtrup, S. and Lovtrup, M. (1988). The morphogenesis of molluscan shells: A mathematical account using biological parameters. *J. Morph.* 197, 53–62.

Lutz, H. (1949). Sur la production experimentale de la polyembryonie et de la monstruosite double ches lez oiseaux. *Arch. d'Anat. Micro. et de Morph.,* 38, 79–144.

Martin, G.R. (1995). Why thumbs are up. *Nature* 374, 410–411.

Meinhardt, H. (1976). Morphogenesis of lines and nets. *Differentiation* 6, 117–123.

Meinhardt, H. (1978). Space-dependent cell determination under the control of a morphogen gradient. *J. theor. Biol.* 74, 307–321.

Meinhardt, H. (1982). Models of Biological Pattern Formation. Academic Press, London (freely available from our website, http://www.eb.tuebingen.mpg.de/meinhardt)

Meinhardt, H. (1983a). A boundary model for pattern formation in vertebrate limbs. *J. Embryol exp. Morphol.* 76, 115–137.

Meinhardt, H. (1983b). Cell determination boundaries as organizing regions for secondary embryonic fields. *Devl. Biol* 96, 375–385.

Meinhardt, H. (1984). Models for positional signalling, the threefold subdivision of segments and the pigmentation pattern of molluscs. *J. Embryol. exp. Morph.* 83, (Supplement) 289–311.

Meinhardt, H. (1993). A model for pattern-formation of hypostome, tentacles, and foot in hydra: how to form structures close to each other, how to form them at a distance. *Dev. Biol.* 157, 321–333.

Meinhardt, H. (1999). Orientation of chemotactic cells and growth cones: Models and mechanisms. *J. Cell Sci.* 112, 2867–2874.

Meinhardt, H. (2002). The radial-symmetric hydra and the evolution of the bilateral body plan: an old body became a young brain. *Bioessays* 24, 185–191.

Meinhardt, H. (2004). Different strategies for midline formation in bilaterians. *Nat Rev Neurosci* 5, 502–510.

Meinhardt, H. (2006). Primary body axes of vertebrates: generation of a near-Cartesian coordinate system and the role of Spemann-type organizer. *Dev Dyn.* 235, 2907–2919.

Meinhardt, H. (2008). Models of biological pattern formation: from elementary steps to the organization of embryonic axes. *Curr. Top. Dev. Biol.* 81, 1–63.

Meinhardt, H. and de Boer, P.A.J. (2001). Pattern formation in *E.coli*: a model for the pole-to-pole oscillations of Min proteins and the localization of the division site. *PNAS* 98, 14202–14207.

Meinhardt, H. and Gierer, A. (1980). Generation and regeneration of sequences of structures during morphogenesis. *J. theor. Biol.* 85, 429–450.

Meinhardt, H. and Klingler, M. (1987). A model for pattern formation on the shells of molluscs. *J. theor. Biol* 126, 63–69.

Mirollo, R.E. and Strogatz, S.H. (1990). Synchronization of pulse-coupled biological oscillators. *SIAM J. On Applied Mathematics* 50, 1645–1662.

Moos, M., Wang, S.W. and Krinks, M. (1995). Anti-dorsalizing morphogenetic protein is a novel TGF-β homolog expressed in the spemann organizer. *Development* 121, 4293–4301.

Murray, J.D. (1989). Mathematical biology. Springer, Heidelberg, New York

Nüsslein-Volhard, C. (1991). Determination of embryonic body axes in the *Drosophila* embryo. *Development Supplement* 1, 1–10.

Nüsslein-Volhard, C. (1996). Gradienten als Organisatoren der Embryonalentwicklung. *Spektrum der Wissenschaften Oktober* 10/1996, pp 38–46.

Nüsslein-Volhard, C. and Wieschaus, E. (1980). Mutations affecting segment number and polarity in *Drosophila*. *Nature* 287, 795–801.

Nellen, D., Burke, R., Struhl, G. and Basler, K. (1996). Direct and long-range action of a dpp morphogen gradient. *Cell* 85, 357–368.

Neumann, D. (1959). Experimentelle Untersuchungen des Farbmusters auf der Schale von *Theodoxus fluviatilis* L. In: Verh. Deutsch. Zoolog. Gesellsch. Münster/Westph. pp.152–156 (Akademische Verlagsgesellschaft Leipzig)

Neumann, D. (1959). Morphologische und experimentelle Untersuchungen über die Variabilität der Farbmuster auf der Schale von *Theodoxus fluviatilis* L. *Z. Morph. Ökol. Tiere* 48, 349–411.

Neumann, D. (1959). Musterumschlag auf der Molluskenschale. *Experienta* 15, 178.

Niehrs, C., Steinbeisser, H. and DeRobertis, E.M. (1994). Mesodermal patterning by a gradient of the vertebrate homeobox gene *goosecoid*. *Science* 263, 817–820.

Oliver, G, Sidell, N, Fiske, W, Heinzmann, C, Mohandas, T, Sparkes, RS and De Robertis, ED (1989). Complementary homeo protein gradients in developing limb buds. *Genes Dev.* 3, 641–650.

Oppenheimer, P. (1986). Real time design and animation of fractal plants and trees. *Computer Graphics* 20, 55–64.

Ozbudak, E.M., Becskei, A. and van Oudenaarden, A. (2005). A System of Counteracting Feedback Loops Regulates Cdc42p Activity during Spontaneous Cell Polarization. *Dev. Cell* 9, 565–571.

Parr, B.A. and McMahon, A.P. (1995). Dorsalizing signal *Wnt-7a* required for normal polarity of D-V and A-P axes of the mouse limb. *Nature* 374, 350–353.

Pickover, C.A. (1989). A short recipe for seashell synthesis. *IEEE Computer Graphics and Applications* 9, 8–11 .

Pickover, C.A. (1991). Computers and the Imagination. St. Martin's Press

Plath, P.J, Schwietering, J. (1992). Improbable event in deterministically growing patterns. In: Fractal Geometry and Computer. Graphics (J. L. Encarnucao, H. O. Peitgen, Skas, G. Englert, Eds.) Springer Verlag.

Prigogine, I. and Lefever, R. (1968). Symmetry breaking instabilities in dissipative systems. II. *J. chem. Phys.* 48, 1695–1700.

Prum, R.O. and Williamson, S. (2002). Reaction-diffusion models of within-feather pigmentation patterning. *Proc. R. Soc. London B* 269, 781–792.

Prusinkiewicz, P. (1994). Visual models of morphogenesis. *Artificial Life* 1, 61–74.

Prusinkiewicz, P. and Streibel, D. (1986). Constraint-based modeling of three-dimensional shapes. In *Proceedings of Graphics Interface '86 - Vision Interface '86*, pp. 158–163.

Psychoyos, D. and Stern, C.D. (1996). Restoration of the organizer after radical ablation of Hensen's node and the anterior primitive streak an the chick embryo. *Development* 122, 3263–3273.

Ptashne, M., Jeffrey, A., Johnson, A.D., Maurer, R., Meyer, B.J., Pabo, C.O., Roberts, T.M. and Sauer, R.T. (1980). How the lambda repressor and Cro work. *Cell* 19, 1–11.

Raskin, D.M. and de Boer, P.A.J. (1999). Rapid pole-to-pole oscillation of a protein required for directing division to the middle of *Escherichia coli*. *PNAS* 96, 4971–4976.

Raup, D.M. (1962). Computer as aid in describing form in gastropod shells. *Science* 138, 150–152 .

Raup, D.M. (1969). Modeling and simulation of morphology by computer. Proceedings of the North American Paleontology Convention

Raup, D.M. and Michelson, A. (1965). Theoretical morphology of the coiled shell. *Science* 147, 1294–1295 .

Regulski, M., Dessain, S., Mcginnis, N. and Mcginnis, W. (1991). High-affinity binding-sites for the deformed protein are required for the function of an autoregulatory enhancer of the deformed gene. *Genes & Development* 5, 278–286.

Risau, W. and Flamme, I. (1995). Vasculogenesis. *Annu. Rev. Cell Dev. Biol.,* 11, 73–91.

Sabelli, B. (1979). Guide to Shells. Simon and Schuster

Sachs, T. (1991). Cell polarity and tissue patterning in plants. *Development Supplement* 1, 83–93.

Saunders, B.W. (1984). *Nautilus* growth and longevity: Evidence from marked and recaptured animals. *Science* 224, 990–992.

Schoute, J.C. (1913). Beiträge zur Blattstellung. *Rec. trav. bot. Neerl.* 10, 153–325.

Segel, L.A. (1984). Modelling dynamic phenomena in molecular and cellular biology. Cambridge University Press, Cambridge

Segel, L.A. and Jackson, J.L. (1972). Dissipative structure: an explanation and an ecological example. *J. theor. Biol.* 37, 545–549.

Seilacher, A. (1972). Divaricate patterns in pelecypod shells. *Lethaia* 5, 325–343.

Seilacher, A. (1973). Fabricational noise in adaptive morphology. *Systematic Zool.* 22, 451–465.

Shapiro, L. and Losick, R. (2000). Dynamic spatial regulation in the bacterial cell. *Cell* 100, 89–98.

Shimizu, H. and Fujisawa, T. (2003). Peduncle of Hydra and the heart of higher organisms share a common ancestral origin. *Genesis* 36, 182–186.

Siegert, F. and Weijer, C.J. (1992). 3–dimensional scroll waves organize *Dictyostelium* slugs. *Proc. Natl. Acad. Sci. USA* 89, 6433–6437.

Simpson-Brose, M, Treisman, J and Desplan, C (1994). Synergy between the *hunchback* and *bicoid* morphogens is required for anterior patterning in *Drosophila*. *Cell* 78, 855–865.

Smith, R.S., Guyomarc'h, S., Mandel, T., Reinhardt, D., Kuhlemeier, C. and Prusinkiewicz, P. (2006). A plausible model of phyllotaxis. *PNAS* 103, 1301–1306.

Spemann, H. and Mangold, H. (1924). Über Induktion von Embryonalanlagen durch Implantation artfremder Organisatoren. *Wilhelm Roux' Arch. Entw. mech. Org.* 100, 599–638.

Starz-Gaiano, M., Melani, M., Wang, X., Meinhardt, H. and Montell, D.J. (2008). Feedback Inhibition of JAK/STAT signaling by apontic is required to limit an invasive cell population. *Dev. Cell* 14, 726–738.

Stewart, I. (1991). All together now. *Nature* 350, 557–557.

Suchting, S., Freitas, C., le Noble, F., Benedito, R., Breant, C., Duarte, A. and Eichmann, A. (2007). The notch ligand delta-like 4 negatively regulates endothelial tip cell formation and vessel branching. *PNAS* 104, 3225–3230.

Technau, U. and Holstein, T.W. (1995). Head formation in hydra is different at apical and basal levels. *Development* 121, 1273–1282.

Thompson, d'Arcy, W. (1952). On Growth and Form. Cambridge

Thompson, d'Arcy, W. (1961). On Growth and Form (Abridged Edition). Cambridge University Press, Cambridge

Tickle, C., Summerbell, D. and Wolpert, L. (1975). Positional signalling and specification of digits in chick limb morphogenesis. *Nature* 254, 199–202.

Trembley, A. (1744). Memoires pour servir a l'histoire d'un genre de polypes d'eau douce. *J. u. H. Verbeek, Leiden.* , .

Turing, A. (1952). The chemical basis of morphogenesis. *Phil. Trans. B.* 237, 37–72.

Udolf, G., Lüer, K., Bossing, T. and Technau, G.M. (1995). Commitment of the CNS progenitors along the dosroventral axis of *Drosophila* neuroectoderm. *Science* 269, 1278–1281.

van Iterson, G. (1907). Mathematische und mikroskopischanatomische Studien über Blattstellungen. *Fischer, Jena.* , .

Vincent, J.P. and Lawrence, P.A. (1994). It takes three to distalize. *Nature* 372, 132–133.

von Rosenhof, R. (1755). Historie der Polypen und anderer kleiner Wasserinsecten. Insektenbelustigung. *Vol. III*. Nürmberg

Wacker, S.A., Jansen, H.J., McNulty, C.L., Houtzager, E. and Durston, A.J. (2004a). Timed interactions between the Hox expressing non-organiser mesoderm and the Spemann organiser generate positional information during vertebrate gastrulation. *Dev. Biol.* 268.207–21, .

Waddington, C. and Cowe, R. (1969). Computer simulation of a molluscan pigmentation pattern. *J. Theor. Biol.* 25, 219–225.

Wanscher, J.H. (1971). Considerations on phase-change and decorations in snail shells. *Hereditas* 71, 75–94.

Ward, P.D. and Chamberlain, J. (1983). Radiographic observation of chamber formation in *Nautilus pompilius*. *Nature* 304, 57–59.

Wilcox, M., Mitchison, G.J. and Smith, R.J. (1973). Pattern formation in the blue-green alga, *Anabaena*. I. Basic mechanisms. *J. Cell Sci.* 12, 707–723.

Willmann, R. (1983). Die Schnecken von Koos. *Spektrum der Wissenschaft (Februar)* pp. 64–76.

Winfree, A.T. (1980). The geometry of biological time. Springer Verlag, New York, Heidelberg, Berlin.

Wolfram, S. (1984). Cellular automata as models of complexity. *Nature* 341, 419–424.

Wolpert, L. (1969). Positional information and the spatial pattern of cellular differentiation. *J. theor. Biol.* 25, 1–47.

Yoon, H.S. and Golden, J.W. (1998). Heterocyst pattern-formation controlled by a diffusible peptide. *Science* 282, 935–938.

Zigmond, S.H. (1977). Ability of polymorphonuclear leukocytes to orient in gradients of chemotactic factors. *J Cell Biol.* 75, 606–617.

Index

Names of mollusks

Other objects