FVS-ACD vs OSM

An independent re-derivation, and a Chapter-11 lever to close the operational gap

Acadian Growth & Yield Meeting | 18 August 2026

Aaron Weiskittel | Center for Research on Sustainable Forests and Cooperative Forestry Research Unit, University of Maine

Honest read: on the operational domain OSM and FVS-ACD are close and OSM is strong. FVS-ACD's edge is biological realism, long horizon, QMD, and transparency.

CONTEXT

Why revisit the 2018 benchmark

2018 (Hennigar & Weiskittel, ECANUSA). OSM trailed only slightly behind FVS-ACD, and beat it on merchantable volume-increment bias and RMSE in ME, NB, and NS. That result set the FVS-ACD to-do list.

2026. Independent re-derivation on fresh data with a common FIA NSVB currency. If OSM's 2018 edge does not replicate, say so with evidence. It largely does not.

1. **Site productivity** ESI holds; height edge is extrapolation
2. **Data harmonization** MagPlot + FIA on one NSVB currency
3. **New calibration approaches** Multi-dimensional scorecard; Ch 11 modifiers
4. **Integrating climate** Current and 2030 ESI as a scenario
5. **AI and productivity** Accelerated the work, caught our own error

His five agenda topics, and what we bring to each

DATA HARMONIZATION

Data harmonized onto one currency

705

ESI-covered in-domain stands, the pooled NSVB scorecard sample

NSVB

FIA volume and biomass as the common currency for every model

ESI v8

30 m climate-and-terrain site driver, cross-border coverage

5

models scored head to head v12.5.0, v12.7.0, v12.9.x, v12.10.1, OSM

Also: recent post-2010 Maine FIA (independent in time from the 1999 to 2005 training FIA), and SILC Seven Islands for the long horizon. Coordinates stay on Cardinal; aggregates only.

OSM leads pooled, v12.9.x closes the gap

Figure 1. Five-way scorecard, level bias percent with bootstrap 95% CIs and green/yellow/red flags, nine NSVB metrics on the operational in-domain set (n = 705)

	v12.5.0	v12.7.0	v12.9.x	v12.10.1	OSM
Basal area (BA)	+2.9% [-0.0, 6.2]	-13.4% [-16.0, -10.5]	-9.9% [-12.7, -6.9]	-9.9% [-12.7, -6.9]	-2.0% [-4.6, 1.0]
Stand density index (SDI)	+5.1% [2.2, 8.3]	-12.1% [-14.7, -9.1]	-9.4% [-12.2, -6.1]	-9.4% [-12.3, -6.3]	-4.0% [-6.8, -1.2]
Quadratic mean diameter (QMD)	-4.9% [-6.9, -3.1]	-0.7% [-2.6, 1.2]	+1.0% [-1.0, 2.9]	+1.0% [-1.0, 2.9]	+11.0% [8.9, 13.1]
Top height (TOPHT)	+4.6% [3.3, 6.1]	+3.7% [2.4, 5.1]	+4.0% [2.7, 5.5]	+4.0% [2.7, 5.4]	+4.4% [3.2, 5.5]
Total volume (TOTVOL)	+9.3% [5.2, 13.5]	-8.3% [-11.6, -4.8]	-3.9% [-7.6, -0.0]	-3.9% [-7.7, 0.2]	+1.6% [-2.1, 5.5]
Merch volume (MERCHVOL)	+8.4% [4.4, 12.4]	-9.1% [-12.3, -5.6]	-4.8% [-8.3, -0.9]	-4.8% [-8.4, -1.0]	+1.0% [-2.4, 4.8]
Sawlog volume (SAWVOL)	+7.8% [3.7, 12.1]	-9.5% [-13.1, -5.7]	-5.2% [-9.1, -0.9]	-5.2% [-8.8, -1.3]	+1.5% [-2.2, 5.8]
Aboveground biomass (BIOMASS)	+9.7% [6.0, 13.8]	-8.1% [-11.3, -4.8]	-3.9% [-7.3, -0.4]	-3.9% [-7.5, -0.2]	+1.0% [-2.2, 4.8]
Stand value (VALUE)	+11.0% [6.5, 16.0]	-6.7% [-10.4, -2.8]	-2.1% [-6.3, 2.4]	-2.1% [-6.3, 2.4]	+2.6% [-1.4, 7.1]


green |bias| < 5% and NSE > 0.6 yellow intermediate red |bias| >= 10% or NSE < 0.4

Corrected merge. An index-join artifact had made OSM look imprecise. With a single keyed merge OSM's modeling efficiency is positive (0.70 to 0.75) and it is the most accurate model on the pooled domain.

OSM and v12.9.x are close. v12.9.x roughly halves the incumbent v12.5.0 volume over-prediction; v12.7.0 over-corrects into a similar-sized under-prediction.


QMD is a durable FVS advantage (about +1 to +2 percent vs OSM +11).

Hardwood is the shared frontier



Softwood is equivalent for every model. Hardwood is the shared frontier: all models over-predict, even OSM (+15). v12.9.x roughly halves the v12.5.0 hardwood bias.

Biological realism: only the self-thinning versions pass all five




The one model that plateaus at the realistic maximum, and it passes all five relations in the seeded-gradient battery; OSM is not scored in that seeded battery.

v12.5.0 never self-thins (basal area runs to 65 with thousands of stems alive), failing yield-density, Reineke, and Sukachev. OSM flags the plateau and Reineke: it does not self-thin on the long horizon.


Honest limit: ingrowth. Recruitment is under-predicted (NB 17.9 vs 39.8, QC 11.7 vs 19.7 stems ha^{-1} yr^{-1}). Direction is correct, it responds to density and management; the level is low. A disclosed limit and a v13 target.

Long horizon: FVS-ACD self-thins, OSM does not



The operational short-interval domain agreement hides a long-horizon divergence driven by self-thinning. This is why FVS-ACD is the choice for carbon, old-growth, and climate work.

Site productivity: keep ESI



Bake-off settled. ESI vs BGI vs CPSI vs legacy CSI, out of sample with spatially blocked CV. ESI wins the dominant-height anchor and ties for diameter growth. BGI and CPSI carry no height signal and lack cross-border coverage.

Why pooled parity is deferred, not hand-waved. We tested a top-height anchor correction and did not ship it. In the shipped config a +4% top-height over-prediction offsets a −10% basal-area under-prediction, so correcting height alone drives volume further negative. True parity needs a joint height-and-basal-area re-anchor. That is v13.0.

NEW CALIBRATION APPROACHES

Only OSM carries a management term

Component	FVS-ACD v12	OSM v2.26.1
Diameter growth	Parametric Kuehne 2022	Random Forest table
Management term	None (verified)	IsManaged column (verified)
Site in DG	Continuous ESI	Binary IsGoodSite
Height calibration	None in base	Local-observation routine
Self-thinning	Composition-aware SDImax	No long-horizon plateau


The verified difference

OSM's diameter-growth table literally carries an **IsManaged** column. FVS-ACD v12 has no management term anywhere.

That is the specific, source-level reason OSM is competitive on the heavily managed benchmark, and exactly what v12.10.1 targets.

OSM ships as a compiled binary; the exposed diameter-growth RF table verifies the management covariate. Mortality and ingrowth management terms are documented, not source-exposed.

The Chapter-11 lever: a fitted release, not a clamp




Direct effect only (Ch 11).
The base equations already carry the indirect structural effect. The modifier isolates the direct effect by the multi-step method (Wang 1990).

Planted = Type 2, sustained.
PCT = Type 1, transient, decaying.

Result: robust re-fit. The direct effect is a fitted additive release (planted +3.8, PCT +2.4 m^2 ha^{-1} BA), not a clamped ratio, so its fold coefficients move with the data. v12.10.1 clearly beats OSM on planted and edges OSM on managed out of sample. The effect is larger than the old 1.6 clamp allowed, which is why the capped version still under-predicted planted.

The modifier closes the managed gap



Robust additive direct-effect release (planted +3.8, PCT +2.4 m^2 ha^{-1} BA), clamp removed; engine carries a sustained (planted) or decaying (PCT) diameter-growth effect bounded by self-thinning. In-sample managed mean absolute bias across nine metrics: v12.9.x 10.2 to v12.10.1 1.3 (OSM 8.3). Cross-validated on the next slide. Gated, default off, so unmanaged stands are untouched.

NEW CALIBRATION APPROACHES

Does the lever hold out of sample?

Mean absolute level bias across nine metrics, lower is better. OSM reference is class-specific.

Cross-validation scheme	Planted	PCT	Managed
Leave-one-installation-out	4.0	3.0	1.3
Leave-one-province-out	5.6	4.1	1.5
Managed split-half (200 reps)	n/a	n/a	5.6
OSM reference	**16.7**	**3.7**	**8.3**

What survives the red team

Managed is a clean out-of-sample win: 1.3 leave-one-installation-out and 1.5 leave-one-province-out vs OSM 8.3.

Planted is a robust win over OSM in every scheme (4.0 vs 16.7) and transfers across the NB to QC border.

PCT now transfers NB to QC (4.1 vs OSM 3.7); the PCT basal-area residual closes. Quebec edge still reported as out of domain, conservatively.

The robust additive re-fit removed the 1.6 clamp; fold coefficients now move with the data (planted basal-area release 3.6 to 4.1 across folds) and the out-of-sample numbers hold. In-sample class bias is near zero by construction, so the cross-validated numbers are the evidence.

INTEGRATING CLIMATE

Climate enters as a scenario driver, not a benchmark

A validated base and a separate scenario driver.

ESI v8 gives current and 2030 climate site surfaces. The projected surface is a forward scenario driver, kept separate from the validated base and never benchmarked on.

For a credible forecast the future step uses a parametric climate-sensitive form (a GAM ratio on the dominant-height anchor), not a random-forest extrapolation. RCP4.5 gives a defensible cold-limited signal of +1.9 m (2030s) to +4.3 m (2090s).

Rule

Validate on observed climate.
Project with a parametric form.
Never benchmark on the projected surface.

AI AND PRODUCTIVITY

AI accelerated the work, humans kept it honest

Accelerated	Weeks of benchmarking, scorecards, and figures compressed into days of iteration.
Kept honest	A human check caught an index-join merge bug that had wrongly made OSM look imprecise; corrected to positive NSE.
Reproducible	Every number traces to a script and a keyed data file on Cardinal; coordinates never leave the cluster.

ROADMAP

v13 is the fall arc: what the data lets us commit to now

1	**v13.0 Height and domain**	Bound the calibrated domain: gate off predictions outside the ESI range so the anchor never extrapolates. The joint re-anchor is now fit. Dominant-height and basal-area increments are each individually unbiased on the 13,894 ME/NH/VT in-domain pairs (increment ratio 1.00 and 1.00), so the volume identity stays unbiased. The v13.0 model variant is built and scoring now on the held-out MagPlot NSVB scorecard. Mature Acadian data still identifies the asymptote for the New Brunswick to Quebec transfer.
2	**v13.1 Hierarchical mortality and ingrowth**	The full Bayesian hierarchical mortality and ingrowth fit is now running. A region random slope on management, standard deviation 1.21 on the log-hazard scale, far larger than the mean effect, so a single capped constant is the wrong object; leave-one-region-out transfer beats it in 5 of 5 held-out regions. This is also where the under-predicted ingrowth is refit (NB 45, QC 59 percent of observed today). Management then enters v13.2 as a growth-scale release, with prediction intervals and climate following.
3	**The ask of the group**	Provincial managed remeasurements and PSP treatment histories, with the provincial site surfaces. This is the data that unblocks the New Brunswick to Quebec management transfer, the one place the managed calibration still leans on thin planted and PCT samples. August is the moment to request a harmonized data-sharing arrangement so v13 is fit on the full Acadian record.

VERDICT & ROADMAP

Close on the operational domain, ahead on realism and transparency

- On the operational inventory domain OSM and FVS-ACD are close and OSM is strong: OSM still leads pooled (mean absolute bias 3.2 vs 4.9, modeling efficiency 0.75 vs 0.71, TOST equivalence 8 of 9 vs 7 of 9). FVS-ACD does not claim pooled parity; its edge is biological realism and transparency.

- It is the one model that plateaus at the realistic maximum, and it passes all five relations in the seeded-gradient battery, where OSM is not scored. Add a durable QMD advantage, a validated climate-ready ESI driver, and full transparency in R. The one honest limit on this slide is ingrowth, under-predicted but in the right direction and a v13 target.

- **August is a v12.10.1 conversation.** Its Chapter-11 direct-effect management modifiers, re-fit as a robust additive release (planted +3.8, PCT +2.4 m^2 ha^{-1} basal area, clamp-free), win the managed cell out of sample (managed 1.3 vs OSM 8.3, planted 4.0 vs 16.7) and transfer PCT NB to QC (4.1 vs 3.7). Gated, default off; unmanaged stands untouched.

- **v13 is the fall arc, and it is now moving:** v13.0's joint height and basal-area re-anchor is fit, individually unbiased in domain, and scoring now against OSM; v13.1's hierarchical mortality and ingrowth refit is running; management enters as a growth-scale release in v13.2, then prediction intervals and climate. The key ask stays the provincial managed remeasurement and PSP treatment data that unblocks the NB to Quebec management transfer.

Questions / Comments

Aaron Weiskittel | aaron.weiskittel@maine.edu | Center for Research on Sustainable Forests and Cooperative Forestry Research Unit, University of Maine

All results reproducible and openly archived; plot coordinates stay on the analysis cluster, aggregates only.

Support: CRSF and CFRU, University of Maine | Maine-FOREST (NSF EPSCoR OIA-1849227) | PERSEUS (USDA NIFA)