



The purpose of this document is to provide an introduction on how to use the Holos model (version 4) and the required vs. optional inputs.

For the purpose of this training, we are going to create a farm that has an annual beef production system, and a feed crop production system. The farm is located in Manitoba near Portage La Prairie.

Launch Holos

Please note that Holos 4 can be installed on a Microsoft Windows PC only. Mac OS will be supported in the next version.

Launch Holos by double-clicking on the Holos desktop icon. If there are no saved farms in the system, Holos will create a new farm and ask the user for a farm name and an optional comment

(Figure 1). If there is already a saved farm in the system, Holos will ask the user to open the existing farm or to create a new farm (Figure 2).

Enter “**Holos 2022**” as a farm name and “**training version**” as the “**comment**”. Ensure to click the “**Advanced Mode**” button so that additional features and functionality will be available. Click “**Ok**” to proceed to the next screen

Ensure “**Metric**” is selected as the unit of measurement type and then click the “**Next**” button at the bottom of the screen (Figure 3).




Figure 1: Entering a name for the new farm.



Figure 2: If a farm has been previously saved, Holos will prompt to re-open that farm.




Figure 3: Select metric as the unit of measurement.

Creating and locating the new beef farm

The beef farm that we will create for this exercise is located in the province of Manitoba. Select “Manitoba” on the “Select a province” screen, and then click the “Next” button.




Figure 4: Select manitoba as the province.

Holos uses **Soil Landscapes of Canada** (SLC), which are a series of GIS coverages that show the major characteristics of soils and land for all of Canada (compiled at a scale of 1:1 million). SLC polygons may contain one or more distinct soil landscape components.

The “Farm Location” screen brings up a map of Canada with the province of Manitoba centered on the screen (Figure 5).

The map contains red colored polygons that can be selected by moving the cursor over the region that contains the location of your farm. You can zoom in or out of the map by using the mouse wheel or by hovering the cursor over the zoom icon at the bottom of the screen.

The beef farm for this example is located between Winnipeg and Portage la Prairie (Portage) with SLC polygon number **851003**.

1. Find and right-click on this polygon to select it on the map. Note that at this point daily climate data will be downloaded from [NASA](#). You can also select a specific polygon by clicking the "Enter a Polygon" button (Figure 6).

Note: Climate data is central to most calculations performed by Holos. For the most accurate estimation of farm emissions, measured climate data should be provided by the user which will override the default data obtained from the NASA weather API.

Holos will use daily precipitation, temperature, and potential evapotranspiration values to model soil carbon change (climate parameter), nitrous oxide emissions, as well as ammonia volatilization.

Farm Location

Select the location of your farm by right-clicking on a region (use mouse wheel to zoom in or out)




Figure 5: Map of the Manitoba province showing the different selectable polygons.

Farm Location

Select the location of your farm by right-clicking on a region (use mouse wheel to zoom in or out)




Figure 6: Specifying a polygon ID using the option button.

Once the farm location is selected, soil information (texture, sand, and clay proportions) for the types of soils found in this region are displayed on the right side of the screen. It's possible that more than one soil type per region will be found and the user is expected to select their soil type from this list or use the default selection. (Figure 7)

For this tutorial, keep the default first selected soil type, and keep the default “Hardiness zone”.

Farm Location

Select the location of your farm by right-clicking on a region (use mouse wheel to zoom in or out)




Figure 7: Multiple soil types might be available for a given region.

Note: Soil data obtained from the user's selected location will be used in the calculation of location-specific N₂O emission factors. Properties such as soil texture, top layer thickness, and soil pH are required for these calculations, and can be overwritten.

Click the “Next” button to proceed to the next step.

Selecting Farm Components

Now that the farm location has been selected, we can move on to the “Component Selection” screen. This is where the user can select different components for their farm. Holos will display all available components on the left side of the screen under the “All Available Components” column (Figure 8). These components are grouped into various categories including Land Management, Beef Production and Dairy Cattle.

If we click on the drop down dropdown button next to a categories' name, we can then see the available components in that category. For this portion of the training section, we will be working with the “Land management” and “Beef production” categories.

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components	My Farm
<input checked="" type="checkbox"/> Land management	
<input checked="" type="checkbox"/> Beef production	
<input checked="" type="checkbox"/> Dairy cattle	
<input checked="" type="checkbox"/> Swine	
<input checked="" type="checkbox"/> Sheep	
<input checked="" type="checkbox"/> Other livestock	
<input checked="" type="checkbox"/> Poultry	
<input checked="" type="checkbox"/> Infrastructure	

Back Next

Canada
851003 (Portage, Manitoba)

Figure 8: The available components screen. Specific components can be chosen here to include in the farm.

The Holos model is designed to define the land management before livestock. This is because we are allowing livestock to be placed onto a specific Pasture (field) for grazing, and that is easier done when a pasture field has been defined already (otherwise the user would have to interrupt the livestock setup to setup a field).

Crop and Hay Production

Now we can add our first component to the farm. Drag a “Field” component from the left side of the screen and drop it on the “My Farm” on the right side (Figure 9). The screen will now update to reflect this new component that you have added to your farm. Holos will label the field as “Field #1”. At this point, we can now enter production information related to the crop being grown on this field.




Figure 9: Adding a component to the farm.

Wheat with Cover Crop

Our first field on the farm will grow continuous wheat with a cover crop of hairy vetch. Change the following elements in the "Field #1" component.

1. Rename the field to "**Wheat & Hairy Vetch**" in the "**Step 1**" section of the screen.
Change the area of the field to **18 ha**.
2. Select "**Wheat**" as the main crop and "**Hairy Vetch**" as the cover crop in "**Step 2**".
3. Under the "**General**" tab:
 - Enter a yield of **3000 kg/ha** (wet weight). The dry weight value will automatically be calculated based on the moisture content of crop value.
 - Select "**Reduced Tillage**" as the tillage type.
 - Enter "**200**" as the amount of irrigation.

- Select '0' as the pesticide passes.
- Leave Harvest method as default selection.

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components	My Farm	Wheat & Hairy Vetch																																												
<input type="checkbox"/> Land management <ul style="list-style-type: none"> <input type="checkbox"/> Crop Rotation <input type="checkbox"/> Shelterbelt <input type="checkbox"/> Field <input type="checkbox"/> Beef production <ul style="list-style-type: none"> <input type="checkbox"/> Dairy cattle <input type="checkbox"/> Swine <input type="checkbox"/> Sheep <input type="checkbox"/> Other livestock <input type="checkbox"/> Poultry <input type="checkbox"/> Infrastructure 	<input type="checkbox"/> Land management <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Wheat & Hairy Vetch 	Step 1: Define the general properties for this field <table border="1"> <tr> <td>Field name :</td> <td>Wheat & Hairy Vetch</td> </tr> <tr> <td>Total area of this field :</td> <td>18 (ha)</td> </tr> <tr> <td>Start year of field history :</td> <td>1985</td> </tr> <tr> <td>End year :</td> <td>2022</td> </tr> </table> Step 2: Click the 'Add Crop' button to describe the crops grown on this field. Enter one crop for continuous fields (i.e. continuous wheat) and the first phase for crop rotations (i.e. wheat, fallow). Holos will use this entry to populate the field history. <table border="1"> <tr> <td>Add Crop</td> </tr> <tr> <td>Year</td> <td>Crop</td> <td>Winter/Cover/Undersown Crop</td> </tr> <tr> <td>></td> <td>2022</td> <td>Wheat</td> </tr> <tr> <td></td> <td></td> <td>Hairy Vetch (<i>Vicia villosa</i> roth)</td> </tr> </table> Step 3: Adjust details of the crop (wheat) grown on this field <table border="1"> <tr> <td>General</td> <td>Fertilizer</td> <td>Manure</td> <td>Winter & Cover Crops</td> <td>Residue</td> <td>Economics</td> </tr> <tr> <td colspan="6"> General Properties <p>Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.</p> <table border="1"> <tr> <td>Yield (wet weight) :</td> <td>3,000 (kg ha⁻¹)</td> </tr> <tr> <td>Yield (dry weight) :</td> <td>2,640 (kg DM ha⁻¹)</td> </tr> <tr> <td>Moisture content of crop :</td> <td>12 (%)</td> </tr> <tr> <td>Tillage :</td> <td>Reduced tillage</td> </tr> <tr> <td>Harvest method :</td> <td>Cash crop</td> </tr> <tr> <td>Amount of irrigation :</td> <td>200 (mm)</td> </tr> <tr> <td>Number of pesticide passes :</td> <td>0</td> </tr> </table> </td> </tr> </table>	Field name :	Wheat & Hairy Vetch	Total area of this field :	18 (ha)	Start year of field history :	1985	End year :	2022	Add Crop	Year	Crop	Winter/Cover/Undersown Crop	>	2022	Wheat			Hairy Vetch (<i>Vicia villosa</i> roth)	General	Fertilizer	Manure	Winter & Cover Crops	Residue	Economics	General Properties <p>Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.</p> <table border="1"> <tr> <td>Yield (wet weight) :</td> <td>3,000 (kg ha⁻¹)</td> </tr> <tr> <td>Yield (dry weight) :</td> <td>2,640 (kg DM ha⁻¹)</td> </tr> <tr> <td>Moisture content of crop :</td> <td>12 (%)</td> </tr> <tr> <td>Tillage :</td> <td>Reduced tillage</td> </tr> <tr> <td>Harvest method :</td> <td>Cash crop</td> </tr> <tr> <td>Amount of irrigation :</td> <td>200 (mm)</td> </tr> <tr> <td>Number of pesticide passes :</td> <td>0</td> </tr> </table>						Yield (wet weight) :	3,000 (kg ha ⁻¹)	Yield (dry weight) :	2,640 (kg DM ha ⁻¹)	Moisture content of crop :	12 (%)	Tillage :	Reduced tillage	Harvest method :	Cash crop	Amount of irrigation :	200 (mm)	Number of pesticide passes :	0
Field name :	Wheat & Hairy Vetch																																													
Total area of this field :	18 (ha)																																													
Start year of field history :	1985																																													
End year :	2022																																													
Add Crop																																														
Year	Crop	Winter/Cover/Undersown Crop																																												
>	2022	Wheat																																												
		Hairy Vetch (<i>Vicia villosa</i> roth)																																												
General	Fertilizer	Manure	Winter & Cover Crops	Residue	Economics																																									
General Properties <p>Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.</p> <table border="1"> <tr> <td>Yield (wet weight) :</td> <td>3,000 (kg ha⁻¹)</td> </tr> <tr> <td>Yield (dry weight) :</td> <td>2,640 (kg DM ha⁻¹)</td> </tr> <tr> <td>Moisture content of crop :</td> <td>12 (%)</td> </tr> <tr> <td>Tillage :</td> <td>Reduced tillage</td> </tr> <tr> <td>Harvest method :</td> <td>Cash crop</td> </tr> <tr> <td>Amount of irrigation :</td> <td>200 (mm)</td> </tr> <tr> <td>Number of pesticide passes :</td> <td>0</td> </tr> </table>						Yield (wet weight) :	3,000 (kg ha ⁻¹)	Yield (dry weight) :	2,640 (kg DM ha ⁻¹)	Moisture content of crop :	12 (%)	Tillage :	Reduced tillage	Harvest method :	Cash crop	Amount of irrigation :	200 (mm)	Number of pesticide passes :	0																											
Yield (wet weight) :	3,000 (kg ha ⁻¹)																																													
Yield (dry weight) :	2,640 (kg DM ha ⁻¹)																																													
Moisture content of crop :	12 (%)																																													
Tillage :	Reduced tillage																																													
Harvest method :	Cash crop																																													
Amount of irrigation :	200 (mm)																																													
Number of pesticide passes :	0																																													

Show Additional Information : Yes No

Back Next

Canada

851003 (Portage, Manitoba)

Figure 10: Field Component of the farm.

4. Select the '**Fertilizer**' tab and click the "**Add Fertilizer Application**" button. Holos has now added a new fertilizer application for this field and will suggest Urea as the fertilizer blend. A default application rate is calculated based on the yield value entered for this field. Details of this fertilizer application can be changed by clicking the "**Show Additional Information**" button (e.g., season of application, different fertilizer blend, etc.).

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

- Land management
 - Crop Rotation
 - Shelterbelt
 - Field

- Beef production
- Dairy cattle
- Swine
- Sheep
- Other livestock
- Poultry
- Infrastructure

My Farm

Land management

- Wheat & Hairy Vetch

Wheat & Hairy Vetch

Step 1: Define the general properties for this field

Field name :	<input type="text" value="Wheat & Hairy Vetch"/>
Total area of this field :	1 (ha)
Start year of field history :	1985
End year :	2022

Step 2: Click the 'Add Crop' button to describe the crops grown on this field. Enter one crop for continuous fields (i.e. continuous wheat) and the first phase for crop rotations (i.e. wheat, fallow). Holos will use this entry to populate the field history.

Year	Crop	Winter/Cover/Undersown Crop
2022	Wheat	Hairy Vetch (<i>Vicia villosa</i> roth)

Step 3: Adjust details of the crop (wheat) grown on this field

Show Additional Information : Yes No

General Fertilizer Manure Winter & Cover Crops Residue Economics

2022 Wheat - Fertilizer Management

Multiple fertilizer applications can be made to this field throughout the year. To add a fertilizer application, click 'Add Fertilizer Application' to specify the details.

Season of application	Blend	Application rate (kg ha ⁻¹)
Spring	Urea	52.5

A suggested fertilizer application rate of 52.5 (kg ha⁻¹) for wheat with a yield of 3000 (kg ha⁻¹) has been set. This value can be adjusted as needed. X

731014 (Daysland Plain, Alberta) | Total emissions: 505 (kg CO₂e) | Loss : \$ 711.60 | Multi-Year Mode

Figure 11: Adding fertilizer to a field.

Note: At a minimum, Holos requires the area of the field, type of crop grown, and a field-specific fertilizer application rate to calculate direct and indirect nitrous oxide emissions.

Residue management of each crop (and cover crop) can be adjusted in Holos (see 'Residue' tab). Holos provides default values depending on the type of crop being grown and will set a value for percentage of product returned to soil, percentage of straw returned to soil, etc. These residue input settings will have an impact on the final soil carbon change estimates.

Furthermore, biomass fractions and N concentrations can be overwritten by the user, and in this way 'custom' crops can be added that are currently not available.

Page 11 of 34

Native Grasslands Information

The cow-calf operation (defined later on) relies on native pasture for the summer months (May through October).

1. Drag a new “**Field**” tab component to your list of components. Enter the name “**Native Grassland**” in the ‘Field name’ input box.
2. Enter “**100**” as the total area of the field.
3. Select “**Rangeland (Native)**” from the crop list under ‘Crop’ column (**step 2**). Please note that Holos auto populates the ‘**Winter/Cover/Undersown Crop**’ area when a perennial crop is selected.
4. Keep ‘**0**’ as the amount of irrigation and pesticide passes.
5. No fertilizer is used for this crop.

The screenshot shows the 'Native Grassland' field setup in the Holos software. The 'My Farm' section on the left lists 'Native Grassland' and 'Wheat & Hairy Vetch'. The main screen shows 'Native Grassland' selected. Step 1 details the field's general properties: name 'Native Grassland', area '1 (ha)', start year '1985', end year '2022', and history entry at 'Yes'. Step 2 shows the 'Add Crop' button and a table for crop history with entries for 'Year' (2022), 'Crop' ('Rangeland (Native)'), and 'Winter/Cover/Undersown Crop' ('Rangeland (Native)'). Step 3 allows adjusting crop details like irrigation (0 mm) and pesticide passes (0). Additional information includes fuel energy (1.43 GJ ha^-1), herbicide energy (0.46 GJ ha^-1), rate constant (0.0187), and maximum carbon produced (3,036 g m^-2). A note states: 'It is assumed that 30% of root biomass is turned over annually before harvest, and then 100% on harvest for perennials. This can be changed on the 'Residue Tab''. Navigation buttons 'Back' and 'Next' are at the bottom, along with a Canada logo and '851003 (Portage, Manitoba)'.

Figure 12: Native Grasslands information.

Barley Grain and Mixed Hay Rotation

To demonstrate the crop rotation component (as opposed to using individual field components), we

will assume that barley grain and mixed hay are grown in rotation, with the mixed hay under seeded to the barley so that it can be harvested in both main years (example derived from University of Alberta's Breton plots).

When using the "Crop Rotation" component, any sequence of crops that are input into this components will be applied to each individual field that is part of the rotation setup. This means one field is added for each rotation phase, and the rotation shifts so that each rotation phase is present on one field. Since each field can have a different historical management, soil carbon algorithms will run for each field.

For this example, we assume that the farm requires **70 ha** of barley grain and mixed hay, which are grown in rotation. We will need to setup three fields where barley grain is rotated in each field every two years (Figure 13). When using the crop rotation component, the crop management input of a specific crop is repeated on each field in the rotation where the crop is grown.

To setup the rotation:

1. Add one "**Crop Rotation**" component from the available components.
2. To expand the horizontal space available in Holos, click on "**View**" from the top menu bar and select "**Hide List of Available Components**".
3. The rotation of this field **begins in 1985 and ends in 2022**. Under step 1, please ensure that these two values are set as the start and end year respectively.
4. Enter "**70**" ha as the **total area** of this field.
5. Under **Step 2** change the crop to **Barley**. The **year** for this crop should be 2022.
 - Under the **General Tab** enter **3000** kg/ha (wet weight) as the **yield** for this crop.
 - Change the **tillage type** to **Reduced Tillage**.
 - Keep **0** as the amount of irrigation and number of pesticide passes.
6. Now add another crop to this rotation. Click on "**Add Crop**" under "**Step 2**" to add a second crop to the rotation. Note that Holos sets the year for this new crop to 2021 or one before the previous crop's year. This means that Holos is expecting the user to enter crops that have been grown in reverse order back to 1985.

Note: It is not necessary to enter a crop for each individual year going back to 1985, only enough crops to describe a single phase of the rotation will need to be entered by the user. Holos will then copy the phase information and back-populate the field history (e.g., Holos will copy the rotation back to 1985 on behalf of the user).

7. For this newly added crop select "**Tame Mixed(grass/legume)**" as the crop type.
8. Click on the "**Add crop**" button one more time. For this third crop, select "**Tame Mixed(grass/legume)**" once again as the crop type.

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components	My Farm
<input type="checkbox"/> Land management Crop Rotation <input type="checkbox"/> Shelterbelt <input type="checkbox"/> Field <input checked="" type="checkbox"/> Beef production <input checked="" type="checkbox"/> Dairy cattle <input checked="" type="checkbox"/> Swine <input checked="" type="checkbox"/> Sheep <input checked="" type="checkbox"/> Other livestock <input checked="" type="checkbox"/> Poultry <input checked="" type="checkbox"/> Infrastructure	Crop rotation #1 <input type="checkbox"/> Land management Crop rotation #1 <input type="checkbox"/> Native Grassland <input type="checkbox"/> Wheat & Hairy Vetch

Crop rotation #1

Step 1: Starting with one field in the rotation, enter the start and end year of this rotation and the size of the field

Start year :	1985	My crop history will be entered beginning at the start year :	Yes	No
End year :	2022	My rotation uses a left shift over the fields :	Yes	No
Total area of this field :	70 (ha)			

Step 2: Click the 'Add Crop' button to add crops to this rotation. Keep adding crops until the first Holos will add a new field to your farm for each crop in the rotation. phase of this rotation has been listed.

Add Crop

Year	Crop	Winter/Cover/Undersown Crop			
2020	Tame mixed (grass/legume)	Tame mixed (grass/legume)	X	▼	
2021	Tame mixed (grass/legume)	Tame mixed (grass/legume)	X	▲	▼
2022	Barley	None	X	▲	

Crop rotation #1 [Field #1] - Barley

2022 - Barley
2021 - Tame mixed (grass/legume)
2020 - Tame mixed (grass/legume)

Crop rotation #1 [Field #3] - Tame mixed (grass/legume)

2022 - Tame mixed (grass/legume)
2021 - Tame mixed (grass/legume)
2020 - Barley

Crop rotation #1 [Field #2] - Tame mixed (grass/legume)

2022 - Tame mixed (grass/legume)
2021 - Barley
2020 - Tame mixed (grass/legume)

Step 3: Adjust the details of barley. These adjustments will be copied to the other fields in the rotation when barley was grown

Show Additional Information : Yes No

General	Fertilizer	Manure	Winter & Cover Crops	Residue	Economics
General Properties <p>Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.</p> <p>Yield (wet weight) : 3,000 (kg ha⁻¹) Yield (dry weight) : 2,640 (kg DM ha⁻¹) Moisture content of crop : 12 (%) Tillage : Reduced tillage Harvest method : Cash crop Amount of irrigation : 0 (mm) Number of pesticide passes : 0</p>					
Additional Information <p>Fuel energy : 2.39 (GJ ha⁻¹) Herbicide energy : 0.23 (GJ ha⁻¹)</p>					

Back Next

Canada

851003 (Portage, Manitoba)

Figure 13: An example of a crop rotation of three crops.

- Now add harvest data to each of the tame mixed crops. You will need to select each **tame mixed** crop and add the harvest data to that specific crop. So select the first tame mixed crop (2021) and then:
 - Go under the **Harvest Tab** and click **Add Harvest Date** button to create a new harvest.
 - Select a Harvest date of "August 31, 2021", assuming the harvest is done on the same day every year.
 - Select **Mid** for **Forage growth stage**.
 - Enter **5** as the total number of bales.
 - Enter **500** as the **Wet bale weight**.
- Repeat** the above steps (step 9) **for the second** tame mixed crop.

If the tame mixed field is harvested more than once, the "Add Harvest Date" button can be used to add subsequent harvests.

Cow-Calf Operation

Click the view menu item again and uncheck **Hide List of Available Components** option so that we can see all of the available components again.

Adding animal components follows the exact same approach that was used for land management components. Under the **Beef Production** category in the available components, drag and drop one **Beef Cow-Calf** component to the **My farm** section on the right. **Replacement heifers** will not be used in this example, so we will remove this group by clicking the **X** icon right next to its entry under **Step 1**.

The screenshot shows the 'Component Selection' screen for adding a 'Beef Cow-Calf' component. The 'Available Components' sidebar lists categories like Land management, Beef production, Dairy cattle, etc., with 'Beef Cow-Calf' selected. In the 'My Farm' section, 'Beef Cow-Calf' is listed with an 'X' icon. Step 1: 'Define the animal groups for this cow-calf component' shows a table with 'Bulls', 'Cows', and 'Calves' rows, each with an 'X' icon. Step 2: 'Define the management periods for the selected animal group (bulls)' shows a table with three rows: 'Winter feeding' (01-01-2021 - 04-30-2021), 'Summer grazing' (05-01-2021 - 10-31-2021), and 'Extended fall grazing' (11-01-2021 - 12-31-2021). Step 3: 'Define the management of bulls during this selected period of time (2021-01-01 - 2021-04-30)' includes sections for General Management (Number of animals: 4, Daily gain: 0, Start weight: 900.00, End weight: 900.00) and Additional Information (Gain coefficient of bulls: 1.20). Navigation buttons 'Back' and 'Next' are at the bottom.

Figure 14: The Beef Cow-Calf Component.

Entering Beef Cows, Calves and Bulls Information.

Beef Cows - Winter Feeding

Following the annual feeding cycle, the beef farm we are working with is **divided into three management (production) periods**. We can now enter production and management data corresponding to these three management periods.

1. Under the animal groups section in ‘**Step 1**’, make sure that the “**Cows**” row is selected in order to enter the associated management information for that group.
2. Click the management period named “**Winter Feeding**” in ‘**Step 2**’ to activate that management period.
3. Ensure “**January 1, 2021**” is set as the ‘**Start date**’ and that “**April 30, 2021**” is set as the ‘**End date**’.

Next, we can enter data related to the number of animals, diet, manure system, and housing type.

- **General Tab:**

- Enter **150** as the number of animals.
- Keep the remaining entries at their default values.

Note: *The number of animals, average daily gain, and feed quality are the minimum required inputs for calculating methane and nitrous oxide emissions. Length of management periods (i.e., duration of grazing) also will be needed. Housing and manure management information are also important inputs but are relatively more impactful on the emissions of monogastrics.*

- **Diet Tab:**

We are going to create a custom diet for our group of cows during the “**Winter feeding**” management period. (Holos incorporates feed ingredient information from the recently published Nutrient Requirements of Beef Cattle book (2016).

Click on the ‘**Diet**’ tab. Since we are going to create our own custom diet, we will click on the ‘**Custom Diet Creator**’ button. Note that Holos provides a default set of animal diets that can be used.

- **Custom-Diet Creator:**

- Click the “**Add Custom Diet**” button in the “**Step 1**” section of the screen to create a new custom diet.
- Rename this diet to “**My Custom Cow Diet**” then press the Enter key to save the name.
- To add ingredients to our new diet, select “**Alfalfa hay**” from the ingredient list, and then click the “**Add Selected Ingredient to Diet**” button.
- We will add one more ingredient to our diet. Select ‘**Barley Hay**’ from the ingredient list, and then click the “**Add Selected Ingredient to Diet**” button.
- Enter **50%** for ‘**Barley Hay**’ and **50%** for ‘**Alfalfa Hay**’ in “**Step 3**”. Note that Holos now reports the diet being complete since all ingredients total up to 100%.
- Click the “**OK**” button to save the new custom diet

- Select the “**My Custom Cow Diet**” from the drop down-down menu in front of **Diet Type**.

Note: Diet quality information such as crude protein, total digestible nutrient, and fat are required inputs so that Holos can estimate enteric methane emissions from an animal group.

Custom Diet Creator

Step 1: To begin creating a diet press "Add Custom Diet"

Diet name	Forage (% DM)	CP (% DM)	TDN (Ix, %)	Fat (% DM)	NDF (% DM)	ME (Mcal kg^-1)
My Custom Diet	100	15.35	57.7	1.95	49.3	

Step 3: Diet is complete

Ingredient	Percentage in diet (% DM)		
Alfalfa hay	50	X	
Barley hay	50	X	

Total %: 100

Step 2: Add ingredients to your diet by selecting the ingredient and then click “Add Selected Ingredient to Diet”

Full Text Search							
Ingredient	DM (% AF)	Forage (% DM)	CP (% DM)	TDN (% DM)	Starch (% DM)	Fat (% DM)	ME (Mcal kg^-1)
Barley grain	89.7	0	12.8	84.1	56.7	2.2	3
Barley grain flaked	81.1	0	12.5	84	59.3	2	3
Barley hay	88	100	10.9	60.2	5.7	2.4	2.2
Barley silage	33.6	100	12	60.6	9.2	3.5	2.2
Barley straw	85.1	100	6.1	48.3	14.9	1	1.7
Beet pulp, dry	91.5	0	9.1	66.6	0.9	1.1	2.4
Beet pulp, wet	21.9	0	9.5	66.6	1.7	0.9	2.4
Bermudagrass fresh	34.9	100	15.2	57.3	1.8	2.8	2.1

OK

Figure 15: Custom diet creator for Cows animal group.

- Housing Tab:
 - Select **Confined no barn** as the housing type.
- Manure Tab:
 - Select **Deep Bedding** as the manure handling system.

Beef Cows - Summer Grazing

Click on the management period named **Summer Grazing**. Ensure that the start date is set to May 1st, 2021 and the end date is set to October 31st, 2021.

- **General Tab:**
 - Enter **150** as the number of animals.
- **Diet Tab:**
 - Select **High energy protein** as the diet type.

- **Housing Tab:**
 - Select **Pasture** as the housing type.
 - Select **Native Grassland** as the pasture location.
- **Manure Tab:**
 - Select **Pasture** as the manure handling system.

Beef Cows - Extended Fall Grazing

Click on the management period named **Extended Fall Grazing**. Ensure that the start date is set to November 1st, 2021 and the end date is set to December 31st, 2021.

- **General Tab:**
 - Enter **150** as the number of animals.
- **Diet Tab:**
 - Select **Medium energy protein** as the diet type.
- **Housing Tab:**
 - Select **Pasture** as the housing type.
 - Select **Native Grassland** as the pasture location.
- **Manure Tab:**
 - Select **Pasture** as the manure handling system.

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components	My Farm	Step 1: Define the animal groups for this cow-calf component			Step 2: Define the management periods for the selected animal group (cows)																			
<input checked="" type="checkbox"/> Land management	<input checked="" type="checkbox"/> Beef production	<input checked="" type="checkbox"/> Beef Cow-Calf <input type="button" value="Add Group"/> <input checked="" type="checkbox"/> Land management <input checked="" type="checkbox"/> Crop rotation #1 <input checked="" type="checkbox"/> Native Grassland <input checked="" type="checkbox"/> Wheat & Hairy Vetch			<input type="button" value="Add Management Period"/> <table border="1"> <thead> <tr> <th>Management period name</th> <th>Start date</th> <th>End date</th> <th>Number of days</th> </tr> </thead> <tbody> <tr> <td>Winter feeding</td> <td>01-01-2021</td> <td>04-30-2021</td> <td>119</td> </tr> <tr> <td>Summer grazing</td> <td>05-01-2021</td> <td>10-31-2021</td> <td>183</td> </tr> <tr> <td>Extended fall grazing</td> <td>11-01-2021</td> <td>12-31-2021</td> <td>60</td> </tr> </tbody> </table>				Management period name	Start date	End date	Number of days	Winter feeding	01-01-2021	04-30-2021	119	Summer grazing	05-01-2021	10-31-2021	183	Extended fall grazing	11-01-2021	12-31-2021	60
Management period name	Start date	End date	Number of days																					
Winter feeding	01-01-2021	04-30-2021	119																					
Summer grazing	05-01-2021	10-31-2021	183																					
Extended fall grazing	11-01-2021	12-31-2021	60																					
<input checked="" type="checkbox"/> Beef production	<input checked="" type="checkbox"/> Beef Cow-Calf	<input checked="" type="checkbox"/> Beef Cow-Calf <input type="button" value="Add Group"/> <input checked="" type="checkbox"/> Land management <input checked="" type="checkbox"/> Crop rotation #1 <input checked="" type="checkbox"/> Native Grassland <input checked="" type="checkbox"/> Wheat & Hairy Vetch			<input type="button" value="Add Management Period"/> <table border="1"> <thead> <tr> <th>Management period name</th> <th>Start date</th> <th>End date</th> <th>Number of days</th> </tr> </thead> <tbody> <tr> <td>Winter feeding</td> <td>01-01-2021</td> <td>04-30-2021</td> <td>119</td> </tr> <tr> <td>Summer grazing</td> <td>05-01-2021</td> <td>10-31-2021</td> <td>183</td> </tr> <tr> <td>Extended fall grazing</td> <td>11-01-2021</td> <td>12-31-2021</td> <td>60</td> </tr> </tbody> </table>				Management period name	Start date	End date	Number of days	Winter feeding	01-01-2021	04-30-2021	119	Summer grazing	05-01-2021	10-31-2021	183	Extended fall grazing	11-01-2021	12-31-2021	60
Management period name	Start date	End date	Number of days																					
Winter feeding	01-01-2021	04-30-2021	119																					
Summer grazing	05-01-2021	10-31-2021	183																					
Extended fall grazing	11-01-2021	12-31-2021	60																					
<input checked="" type="checkbox"/> Dairy cattle	<input checked="" type="checkbox"/> Beef Stockers & Backgrounder	<input checked="" type="checkbox"/> Beef Stockers & Backgrounder <input checked="" type="checkbox"/> Beef Finisher			<input type="button" value="Add Management Period"/> <table border="1"> <thead> <tr> <th>Management period name</th> <th>Start date</th> <th>End date</th> <th>Number of days</th> </tr> </thead> <tbody> <tr> <td>Winter feeding</td> <td>01-01-2021</td> <td>04-30-2021</td> <td>119</td> </tr> <tr> <td>Summer grazing</td> <td>05-01-2021</td> <td>10-31-2021</td> <td>183</td> </tr> <tr> <td>Extended fall grazing</td> <td>11-01-2021</td> <td>12-31-2021</td> <td>60</td> </tr> </tbody> </table>				Management period name	Start date	End date	Number of days	Winter feeding	01-01-2021	04-30-2021	119	Summer grazing	05-01-2021	10-31-2021	183	Extended fall grazing	11-01-2021	12-31-2021	60
Management period name	Start date	End date	Number of days																					
Winter feeding	01-01-2021	04-30-2021	119																					
Summer grazing	05-01-2021	10-31-2021	183																					
Extended fall grazing	11-01-2021	12-31-2021	60																					
<input checked="" type="checkbox"/> Swine	<input checked="" type="checkbox"/> Sheep	<input checked="" type="checkbox"/> Sheep <input checked="" type="checkbox"/> Other livestock <input checked="" type="checkbox"/> Poultry <input checked="" type="checkbox"/> Infrastructure			<input type="button" value="Add Management Period"/> <table border="1"> <thead> <tr> <th>Management period name</th> <th>Start date</th> <th>End date</th> <th>Number of days</th> </tr> </thead> <tbody> <tr> <td>Winter feeding</td> <td>01-01-2021</td> <td>04-30-2021</td> <td>119</td> </tr> <tr> <td>Summer grazing</td> <td>05-01-2021</td> <td>10-31-2021</td> <td>183</td> </tr> <tr> <td>Extended fall grazing</td> <td>11-01-2021</td> <td>12-31-2021</td> <td>60</td> </tr> </tbody> </table>				Management period name	Start date	End date	Number of days	Winter feeding	01-01-2021	04-30-2021	119	Summer grazing	05-01-2021	10-31-2021	183	Extended fall grazing	11-01-2021	12-31-2021	60
Management period name	Start date	End date	Number of days																					
Winter feeding	01-01-2021	04-30-2021	119																					
Summer grazing	05-01-2021	10-31-2021	183																					
Extended fall grazing	11-01-2021	12-31-2021	60																					

Step 3: Define the management of cows during this selected period of time (2021-01-01 - 2021-04-30) Show Additional Information : Yes No

General Management		Additional Information	
Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period. Number of animals : 150 Daily gain : 0 ($kg\ day^{-1}$) Start weight : 610.00 (kg) End weight : 610.00 (kg) Milk production : 8.00 ($kg\ day^{-1}$)		Milk fat content : 4.00 (%) Milk protein : 3.38 (%) Gain coefficient of cow : 0.80	

General Diet Housing Manure

Canada 851003 (Portage, Manitoba)

Figure 16: Beef Cow-Calf, Cow group

Bulls

Click on the **Bulls** row in the animal group section **Step 1**. Information related to diet, housing and manure management is identical to the cows group.

- Right click on the **Bulls** animal group. A menu will appear allowing you to select the option to copy management periods from another animal group. Since the management for the bulls is similar to the management for the cows, click the **Copy Management From -> Cows** sub-menu item.
- Adjust the **number of bulls** for each of the three management periods to **4**.

Component Selection
Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components	My Farm
Land management	Beef production
Crop Rotation	Beef Cow-Calf
Shelterbelt	
Field	
Beef production	Land management
Beef Cow-Calf	Crop rotation #1
Beef Stockers & Backgrounders	
Beef Finisher	Native Grassland
Dairy cattle	Wheat & Hairy Vetch
Swine	
Sheep	
Other livestock	
Poultry	
Infrastructure	

Beef Cow-Calf

Step 1: Define the animal groups for this cow-calf component

Step 2: Define the management periods for the selected animal group (bulls)

Add Management Period

Management period name	Start date	End date	Number of days
Winter feeding	01-01-2021	04-30-2021	119
Summer grazing	05-01-2021	10-31-2021	183
Extended fall grazing	11-01-2021	12-31-2021	60

Step 3: Define the management of bulls during this selected period of time (2021-01-01 - 2021-04-30)

Show Additional Information : Yes No

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Additional Information

Gain coefficient of bulls : 1.20

Number of animals : 4
Daily gain : 0 (kg day^-1)
Start weight : 900.00 (kg)
End weight : 900.00 (kg)

Back Next

Canada

851003 (Portage, Manitoba)

Figure 17: Copying data from another animal group

Beef Calves

Calves on our farm are born on March 1 and weaned on September 30 at the age of seven months. Using a final weaning rate of 85%, we will have 110 calves from March to September. Following the cows, calves will be in confinement for the months of March and April and will be grazing on pasture from May to September. This will result in two separate management periods.

Click on the **Calves** row in the animal group section **Step 1** to activate the calf group. The first management period will span from **March 1, 2021 to April 30, 2021** and the second management period will span from **May 1, 2021 to September 30, 2021**.

Management Period # 1:

Rename this period from Management Period # 1 to **Confinement**.

- **General Tab:**
 - Enter **110** as the number of animals.
 - All other options as default.
- **Diet Tab:**
 - Select **Medium energy protein** as the diet type.
- **Housing Tab:**
 - Select **Confined no barn** as the housing type.
- Manure Tab:
 - Select **Deep bedding** as the manure handling system.

Management Period # 2:

Rename this period from Management Period # 2 to **Grazing**.

- **General Tab:**
 - Enter **110** as the number of animals.
- **Diet Tab:**
 - Select **High energy protein** as the diet type.
- **Housing Tab:**
 - Select **Pasture** as the housing type.
 - Select **Native Grassland** as the pasture location.
- Manure Tab:
 - Select **Pasture** as the manure handling system.

Beef Stocker & Backgrounder operation

To enter information on backgrounder and stocker animals, we will add a new “**Beef Stockers & Backgrounders**” component to our farm.

- Drag and drop a new **Beef Stockers & Backgrounders** component. The beef farm manages 200 backgrounders (100 steers and 100 heifers).
- Click on the **Heifers** group to activate it and to enter management data for this group.

- For “Management period #1”, enter “**October 1, 2021**” as the ‘**Start date**’ and “**January 18, 2022**” as the ‘**End date**’.
- **General Tab:**
 - Enter **100** as the number of animals.
 - Enter **1.1** kg/day as the daily gain.
 - Enter **240** kg as the start weight.
- **Diet Tab:**
 - Select **Medium Growth** as the diet type.
 - Select **None** as the diet additive.
- **Housing Tab:**
 - Select **Confined no barn** as the housing type.
- **Manure Tab:**
 - Select **Deep bedding** as the manure handling system.
- The management data for the **Steers** group is the same as **Heifers**. Right click on the Steers group to activate the right-click menu and select **Copy Management From -> Heifers**

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the “Available Components” column to the “My Farm” column. Click on an item under the “My Farm” column to enter details.

Available Components	My Farm
<input checked="" type="checkbox"/> Land management	Beef production
<input checked="" type="checkbox"/> Beef production	Beef Cow-Calf
<input checked="" type="checkbox"/> Dairy cattle	Beef Finisher
<input checked="" type="checkbox"/> Swine	
<input checked="" type="checkbox"/> Sheep	
<input checked="" type="checkbox"/> Other livestock	
<input checked="" type="checkbox"/> Poultry	Beef Stockers & Backgrounder
<input checked="" type="checkbox"/> Infrastructure	
	Poultry
	Chicken Meat Production
	Land management
	Crop rotation #1
	Native Grassland
	Wheat & Hairy Vetch

Beef Stockers & Backgrounder

Step 1: Define the animal groups for this backgrounding component

Add Group

Group name	Heifers	X
	Steers	X

Step 2: Define the management periods for the selected animal group (heifers)

Add Management Period

Management period name	Start date	End date	Number of days	
Management period #1	10-01-2021	01-18-2022	109	X

Step 3: Define the management of heifers during this selected period of time (2021-10-01 - 2022-01-18)

Show Additional Information : Yes No

General Diet Housing Manure

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Number of animals : 110
Daily gain : 1.10 (kg day^-1)
Start weight : 240 (kg)
End weight : 360 (kg)

Please review the selected diet information, body weight, and daily gain to ensure animals are being provided with their daily energy requirements. X

[Back](#) [Next](#)

Canada 
851003 (Portage, Manitoba)

Figure 18: Beef stocker & backgrounder operation

Finishing Feedlot Operation

We will now **repeat the steps used for Beef Stockers & Backgrounders** to enter the beef feedlot management data. Drag a new **Beef Finisher** component from **All components** to your list of components.

- The beef farm manages 200 backgrounders (100 steers and 100 heifers) in a feedlot operation for **170 days**.
- Click on the **Heifers** group to activate it and to enter management data for this group.
 - For “Management period #1”, enter “**January 19, 2022**” as the **Start date** and “**July 7th, 2022**” as the **End date**.
 - **General Tab:**
 - Enter **100** as the number of animals.
 - Enter **1.2** kg/day as the daily gain.
 - Enter **350** kg as the start weight.
 - **Diet Tab:**
 - Select **Barley grain based diet** as the diet type.
 - Select **None** as the diet additive.
 - **Housing Tab:**
 - Select **Confined no barn** as the housing type.
 - Manure Tab:
 - Select **Deep bedding** as the manure handling system.
- The management data for the **Steers** group is the same as **Heifers**. Right click on the Steers group to activate the right-click menu and select **Copy Management From -> Heifers**

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components	My Farm
<input checked="" type="checkbox"/> Land management	<input checked="" type="checkbox"/> Beef production
Crop Rotation	Beef Finisher
Shelterbelt	<input checked="" type="checkbox"/> Beef Stockers & Backgrounders
Field	Beef Cow-Calf
<input checked="" type="checkbox"/> Beef production	<input checked="" type="checkbox"/> Land management
Beef Cow-Calf	Crop rotation #1
Beef Stockers & Backgrounders	Native Grassland
Beef Finisher	Wheat & Hairy Vetch
<input checked="" type="checkbox"/> Dairy cattle	
Dairy cattle	
<input checked="" type="checkbox"/> Swine	
<input checked="" type="checkbox"/> Sheep	
<input checked="" type="checkbox"/> Other livestock	
<input checked="" type="checkbox"/> Poultry	
<input checked="" type="checkbox"/> Infrastructure	

Beef Finisher

Step 1: Define the animal groups for this finishing component

Add Group

Group name	Start date	End date	Number of days
Heifers	01-19-2022	07-07-2022	170
Steers			

Step 2: Define the management periods for the selected animal group (steers)

Add Management Period

Step 3: Define the management of steers during this selected period of time (2022-01-19 - 2022-07-07)

Show Additional Information : Yes No

General Diet Housing Manure

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Number of animals : 100
Daily gain : 1.2 (kg day⁻¹)
Start weight : 350 (kg)
End weight : 554 (kg)

Back Next

Canada

851003 (Portage, Manitoba)

Figure 19: Beef Finisher component

Adding a Manure Application to the Wheat Field

Holos has the ability to add manure applications from manure that is sourced from livestock on the current farm or from imported manure (off-site). Since we have now defined our animal components, we can apply manure to any field on our farm.

1. Select the **Wheat & hairy vetch** field from the list of components added to our farm.
2. Click on the **Manure tab** and then click the **Add Manure Application** button.
 - Select **Beef cattle** as the **Manure type**.
 - Select **Livestock** as the **Origin of manure**.
 - Select **Deep Bedding** as the **Manure handling system**.
 - Enter **200 kg/ha** as the amount of manure applied to this field.
3. Note that both chemical fertilizers and manure applications can be made on the same field

Adding supplemental hay/forage for grazing animals

We can also add additional hay/forage for animals that are grazing on a particular field. Since we have now placed a group of animals on the “Native Grassland” field component, and we have also

provided harvest information for our mixed hay crops on the crop rotation component, we can then add an additional forage supplement for these grazing animals.

1. Select the **Native Grassland** field component we created earlier.
2. Click on the **Grazing tab**.
 - Click the **Add Supplemental Hay** button to add additional forage for the animals on this field.
 - Enter "**On-farm**" as the **Sources of bales**.
 - Choose **Crop rotation #1 [Field #2] - Tame Mixed (grass/legume)** under **Field** to select the source of the supplemental hay.
 - Change the **Number of bales** to 1
 - Enter **500** as the wet bale weight.
 - Keep the moisture content as the default value.

Note: It is not recommended to mix different species of grasses together. Here, we are only demonstrating the ability of Holos to add supplemental hay to a field that has grazing animals

The screenshot shows the Holos software interface. At the top, there's a navigation bar with 'File', 'Settings', 'Tools', 'View', and 'Help'. Below it is a 'Component Selection' section with a title 'Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.' A 'My Farm' sidebar on the left lists components: Beef production (Beef Finisher, Beef Stockers & Backgrounders, Beef Cow-Calf), Land management (Crop rotation #1, Native Grassland, Wheat & Hairy Vetch). The main area is titled 'Step 3: Adjust details of the crop (rangeland (native)) grown on this field'. It shows '2022 Rangeland (Native) - Grazing Details'. Under 'Grazing Animal Groups', it lists groups like '4 bulls grazing from 1 May to 31 October (183 days)', '4 bulls grazing from 1 November to 31 December (60 days)', etc., each with an 'Edit Group' link. Below this is a section for 'Add Supplemental Hay/Forage for Grazing Animals'. It includes a table with columns: Date, Source of bales, Field, Number of bales, Wet bale weight (kg), and Moisture content. A single row is selected: '10-17-2022, On-farm, Crop rotation #1 [Field #2] - Tame mixed (grass/legume), 1, 500'. To the right of the table is a vertical scale for moisture content from 0 to 5. At the bottom, there are 'Back' and 'Next' buttons, and the Canada logo with the text '851003 (Portage, Manitoba)'.

Figure 20 - Adding supplemental hay/forage for grazing animals.

Pullet Farm Operation:

We will add one last animal component to our farm. In addition to the beef cattle operations of this farm, we will also be adding a “**Chicken Meat Production**” component to our farm. If you hover your mouse cursor over the “Chicken Meat Production” component under the “Poultry” category, Holos will display a tooltip that gives a brief description of a chicken meat production operation:

“Chicks arriving in the operation from a multiplier hatchery are raised to market weight (1-4 kg)”

1. Drag one “**Chicken Meat Production**” component to the farm. For each group (Pullets and Cockerels), each management period for that group will consist of **400** animals. This means numbers of animals will be consistent throughout the management periods / year.
2. Select the **Pullets** group. The start and end dates for each management period will be:
 - **Brooding Stage:** Start: January 1st, 2022 - End: January 22nd, 2022.
 - **Rearing Stage:** Start: January 23rd, 2022 - End: June 26th, 2022.
 - **Rearing Stage:** Start: June 27th, 2022 - End: November 28th, 2022.
3. For each management period, set number of animals to **400**.
4. Leave the entries in Housing and Manure as default.
 - The management data for the **Cockerels** group is the same as **Pullets**. Right click on the Cockerels group to activate the right click menu and select **Copy Management From -> Pullets**

Timeline Screen

We are now finishing the process of defining our farm. Click the **Next** button to go forward to the timeline screen.

The timeline screen provides a visual layout of all the fields from 1985 to the specified end year for each field. This screen also allows the user to add historical and projected production systems.

The **Add Historical Production System** button enables the user to add a different cropping history to individual fields whereas the **Add Projected Production System** button enables the user to add a future (projected) cropping system to individual fields.

Adding a historical production system

We will assume that the barley grain and mixed hay rotation fields were previously in a continuous

wheat cropping system between **1985 and 2000**.

1. To add a new historical cropping system, select one of the fields that are in the barley grain and mixed hay rotation. To select an item, click on the timeline bar to activate that field. We will select the first field in this rotation (i.e., the field with the name of “**Crop rotation #1 [Field #1] - Barley**”)
2. Click on the **Add Historical Production System** button which will add a new row to the table under the “**Step 1**” section in the upper left section of the screen. Notice that this new entry has the words “**Historical management practice**” added.
3. We will set the end year of this historical management practice to the year **2000**. To adjust this we use the numeric up/down buttons within the cell.
4. Select the newly added **Historical management practice** and then click the “**Edit Selected**” button. This will open a new screen that allows us to adjust the crops grown and the management during this period.
5. Click on the “**Barley**” crop under the “**Step 2**” section. Change the crop type to ‘**Wheat**’ and on the ‘**General**’ tab change the yield to **3,500 kg/ha**. We will keep the other settings unchanged.
6. We also need to remove the “**Tame mixed**” crops from this historical period. Click the ‘x’ icon beside each of the “**Tame mixed**” crops under the “**Step 2**” section. Clicking the ‘x’ icon will remove these crops from the rotation for this period of time.
7. Click “**Ok**” to save adjustments we just made to this field.
8. Repeat these same steps so that the other fields in this rotation also have continuous wheat from **1985 to 2000** using the same steps we used for the first field.

Timeline

Use this screen to adjust the starting and ending dates of your management practices

Fields

Step 1: Select a field from the list or timeline below to activate it

System	Start year	End year	Edit	Remove
Crop rotation #1 [Field #1] - Barley (Historical management practice)	1985	2000		
Crop rotation #1 [Field #2] - Tame mixed (grass/legume) (Historical management practice)	1985	2000		
Crop rotation #1 [Field #3] - Tame mixed (grass/legume)	2001	2022		
Crop rotation #1 [Field #2] - Tame mixed (grass/legume)	2001	2022		
Crop rotation #1 [Field #3] - Tame mixed (grass/legume) (Historical management practice)	1985	2000		
Crop rotation #1 [Field #1] - Barley	2001	2022		
Native Grassland	1985	2022		
Wheat & Hairy Vetch	1985	2022		

Step 2: Now that you have selected your field, add a historical or projected production system or edit the selected production system.

[Add Historical Production System](#)
[Add Projected Production System](#)
[Edit Selected](#)

1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022

Crop rotation #1 [Field #1] - Barley

Crop rotation #1 [Field #1] - Barley (Historical management practice) Wheat 1985 - 2000	Crop rotation #1 [Field #1] - Barley Tame mixed (grass/legume), Tame mixed (grass/legume), Barley 2001 - 2022
---	---

Crop rotation #1 [Field #2] - Tame mixed (grass/legume)

Crop rotation #1 [Field #2] - Tame mixed (grass/legume) (Historical management practice) Wheat 1985 - 2000	Crop rotation #1 [Field #2] - Tame mixed (grass/legume) Tame mixed (grass/legume), Barley, Tame mixed (grass/legume) 2001 - 2022
--	--

Crop rotation #1 [Field #3] - Tame mixed (grass/legume)

Crop rotation #1 [Field #3] - Tame mixed (grass/legume) (Historical management practice) Wheat 1985 - 2000	Crop rotation #1 [Field #3] - Tame mixed (grass/legume) Barley, Tame mixed (grass/legume), Tame mixed (grass/legume) 2001 - 2022
--	--

Native Grassland

Native Grassland Rangeland (Native) 1985 - 2022

[Back](#) [Next](#)

Canada

851003 (Portage, Manitoba)

Figure 21: Customized Timeline Screen**Timeline**

Use this screen to adjust the starting and ending dates of your management practices

Step 1: Select a field from the list or timeline below to activate it

System	Start year	End year	Edit
Wheat & Hairy vetch	1985	2022	
Native Grassland	1985	2022	
Crop rotation #1 [Field #1] - Barley	2001	2022	
Crop rotation #1 [Field #1] - Barley (Historical management practice)	1985	2000	
Crop rotation #1 [Field #2] - Hay mixed (Historical management practice)	2001	2022	
Crop rotation #1 [Field #3] - Hay mixed (Historical management practice)	1985	2000	
Crop rotation #1 [Field #1] - Barley (Historical management practice)	2001	2022	
Crop rotation #1 [Field #2] - Hay mixed (Historical management practice)	1985	2000	
Crop rotation #1 [Field #3] - Hay mixed (Historical management practice)	2001	2022	

85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 00 01 02 03 04 05

Wheat & Hairy vetch

Wheat & Hairy vetch Wheat 1985 - 2022

Figure 22: Adjusted start and end year for production systems on the timeline screen.

Fields

Crop rotation #1 [Field #1] - Barley (Historical management practice)

Step 1: Define the general properties for this field

Total area of this field :	70	(ha)
Start year of field history :	1985	
End year :	2000	

Step 2: Click the 'Add Crop' button to describe the crops grown on this field. Enter one crop for continuous fields (i.e. continuous wheat) and the first phase for crop rotations (i.e. wheat, fallow). Holos will use this entry to populate the field history.

Add Crop

	Crop	Winter/Cover/Undersown Crop
>	Wheat	None

Step 3: Adjust details of the crop (wheat) grown on this field

Show Additional Information : Yes No

General Fertilizer Manure Winter & Cover Crops Residue Economics

General Properties

Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.

Yield (wet weight) :	3,500	(kg ha ⁻¹)
Yield (dry weight) :	3,080	(kg DM ha ⁻¹)
Moisture content of crop :	12	(%)
Tillage :	Intensive tillage	
Harvest method :	Cash crop	
Amount of irrigation :	0	(mm)
Number of pesticide passes :	0	

OK

Figure 23: Editing crops in a historical period of the rotation.

Details Screen

Click the “**Next**” button to go forward to the details screen.

To avoid the requirement that a user needs to provide crop yields going back to 1985 for each field on the farm, the model will use Stats Canada reported crop yields as defaults (where available). The model allows the user to calculate how changes in crop type, yield, tillage, residue management, manure, irrigation or fallow will result in changes to soil carbon.

We will adjust this grid so that we can view the above ground and below ground carbon inputs for our wheat field and then we will adjust the crop yield for one specific year.

1. We will set a filter on the first column named ‘**Field name**’ so that we only display information for our **wheat and hairy vetch field**. Beside the column heading, click the ‘**funnel**’ icon to set a filter. Check the box beside **Wheat & hairy vetch**.
2. On the far left of this screen, click the “**Enable Columns**” sidebar (located near the “Field name” column).
3. Place a check beside **Above ground carbon input** to show the column and remove the check beside the **Notes** column to hide it.
4. Click the **Enable Columns** sidebar again to collapse it.
5. We can now (optionally) adjust the yields for our wheat field for any given year if actual measured yields are available.
6. Adjust the yield for **1987** to be **4,100 kg/ha**.
7. Note that Holos has updated the above ground carbon inputs for this.

Cropping System						
	Reload Data From Previous Screen		Yield Assignment Method : Average Yield			
	Field name	Time period	Year	Crop	Yield (kg ha ⁻¹)	Above ground carbon input (kg C ha ⁻¹)
>	Wheat & Hairy Vetch	Current	1985	Wheat	3000.00	2628.596
	Wheat & Hairy Vetch	Current	1985	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1986	Wheat	3000.00	2270.755
	Wheat & Hairy Vetch	Current	1986	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1987	Wheat	4100.00	3806.290
	Wheat & Hairy Vetch	Current	1987	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1988	Wheat	3000.00	1133.995
	Wheat & Hairy Vetch	Current	1988	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1989	Wheat	3000.00	2361.158
	Wheat & Hairy Vetch	Current	1989	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1990	Wheat	3000.00	2716.506
	Wheat & Hairy Vetch	Current	1990	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1991	Wheat	3000.00	2450.928
	Wheat & Hairy Vetch	Current	1991	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1992	Wheat	3000.00	3147.074
	Wheat & Hairy Vetch	Current	1992	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1993	Wheat	3000.00	1995.680
	Wheat & Hairy Vetch	Current	1993	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1994	Wheat	3000.00	2540.072
	Wheat & Hairy Vetch	Current	1994	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1995	Wheat	3000.00	2270.755
	Wheat & Hairy Vetch	Current	1995	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000
	Wheat & Hairy Vetch	Current	1996	Wheat	3000.00	2803.809
	Wheat & Hairy Vetch	Current	1996	Hairy Vetch (<i>Vicia villosa</i> roth)	3000.00	450.000

Back Next



851003 (Portage, Manitoba)

Figure 24: Details screen

Discover results

Click the “**Next**” button to move to the final results report. Results will now be displayed in a variety of reports and charts.

1. Click on the tab named **Emissions Pie Chart**

Starting with the **Emissions pie chart** we can see an overall breakdown of the enteric CH4, manure CH4, direct and indirect N2O. We are also able to see a detailed breakdown of the sources of these emissions.

2. Click the “Yes” button beside **Show details**

We can see that the biggest source of emissions from our farm is the cow-calf component. If you hover your mouse pointer over any slice of this chart you can get an isolated look at the different emission sources.

3. Click on the tab named **Detailed Emission Report**

The **Detailed Emission Report** will display a monthly or annual GHG emission report. The detailed emission report will report on enteric methane, manure methane, direct & indirect N2O, and CO2 emissions from the farm.

Click the **Report Format (Monthly)** button to switch to a monthly report. Now we can see a monthly breakdown of all emissions from the farm and the emission source.

In the **Unit of measurements** drop-down menu, you can choose to have the results displayed as CO₂ equivalents (CO₂e) or as unconverted greenhouse gas (GHG), and you can also choose the unit of measurement as either tonnes or kilograms.

The **Estimates of Production** report provides total harvest yields, amount of land applied manure, and estimates of milk production for dairy components.

The **Feed Estimate** report provides an estimate of dry matter intake based on energy requirements of the animal and the energy in the feed.

File Settings Tools View Help

Results

The results screen displays various reports about your farming operation. Each tab reports on a specific aspect of your emission footprint

Multyear Carbon Modelling	Estimates of Production	Feed Estimate Report	Manure Management	Economics	Emissions Pie Chart	Overall Emissions	Component Emissions	Detailed Emission Report		
Report Format :	Monthly	Annual	Units of measurement :	Kg CO ₂ e	<input type="button" value="Export to Excel"/>	Show Additional Columns :	Yes	No		
	Farm	Component name	Emission source	Enteric CH ₄ (kg CO ₂ e)	Manure CH ₄ (kg CO ₂ e)	Direct N ₂ O (kg CO ₂ e)	Indirect N ₂ O (kg CO ₂ e)	Farm Energy CO ₂ (kg CO ₂ e)	Upstream CO ₂ (kg CO ₂ e)	Subtotal (kg CO ₂ e)
▲ Backgounding										
Farm #1	Beef Stockers & Backgrounders	Heifers	67020.51	49669.13	8305.72	4606.64	5.61	0.00	129607.62	
Farm #1	Beef Stockers & Backgrounders	Steers	67020.51	49669.13	8305.72	4606.64	5.61	0.00	129607.62	
			134041.02	99338.26	16611.44	9213.28	11.22	0.00	259215.23	
▲ Chicken Meat Production										
Farm #1	Chicken Meat Production	Pullets	N/A	60.77	58.24	81.53	2.72	0.00	203.27	
Farm #1	Chicken Meat Production	Cockerels	N/A	60.77	90.20	81.53	2.72	0.00	235.23	
			0.00	121.55	148.45	163.05	5.45	0.00	438.50	
▲ Cow-calf										
Farm #1	Beef Cow-Calf	Bulls	8090.24	2467.46	494.01	337.39	0.22	0.00	11389.33	
Farm #1	Beef Cow-Calf	Cows	362226.22	110251.94	20786.34	14488.64	8.35	0.00	507761.49	
Farm #1	Beef Cow-Calf	Calves	58735.82	9106.95	1217.47	1033.95	3.09	0.00	70097.29	
			429052.29	121826.36	22497.82	15859.98	11.67	0.00	589248.11	
▲ Crops										
Farm #1	Wheat & Hairy Vetch	Wheat, 18 (ha)	N/A	N/A	4473.01	3901.16	12671.64	3965.19	21045.81	
▲ Finishing										
Farm #1	Beef Finisher	Heifers	47771.36	41686.51	11360.11	6365.08	7.96	0.00	107191.02	
		Totals	658636.04	304659.19	75814.85	63801.56	48450.89	3965.19	1151362.53	

Negative values indicate carbon storage. Positive values indicate greenhouse gas emissions.

Uncertainty

- Enteric CH₄ : 20 % (+/-)
- Manure CH₄ : 20 % (+/-)
- Direct N₂O : 40 % (+/-)
- Indirect N₂O : 60 % (+/-)
- Energy CO₂ : 40 % (+/-)

Emissions are reported on a monthly basis. In some cases, prorating of yearly emissions occurs. Pro-rating occurs with soil/cropping N₂O emissions - direct and indirect - as per user entered breakdown. Soil carbon, tree planting carbon, energy CO₂ emissions to crop and spread manure, and poultry and other animals emissions are allocated equally throughout the year.

Canada 851003 (Portage, Manitoba)

Figure 25: Detailed emissions report.




Figure 26: Estimates of production report.




Figure 27: Feed estimate report.

Soil carbon modelling results

On the results screen we can see the change in soil carbon over time by clicking the “Multiyear Carbon Modelling” tab. This tab displays a graph showing the change in soil carbon over time for each one of our fields.

For each field on the graph, you can hover your mouse over the series to get more information for each historical year of the field.

If we click on one of these points, we can then view a more detailed breakdown of these results. We can also export this data by clicking the “Export to Excel” button.




Figure 28: Carbon report section. Allows switching between graph and table format.

Finally...

Whole-systems approach

An ecosystem consists of not only the organisms and the environment they live in but also the interactions within and between. A whole systems approach seeks to describe and understand the entire system as an integrated whole, rather than as individual components. This holistic approach can be very complex and describing the process can be difficult. One method to conceptualize a whole system is with a mathematical model.

The whole-systems approach ensures the effects of management changes are transferred throughout the entire system to the resulting net farm emissions. In some cases, reducing one GHG will actually increase the emissions of another. The whole-systems approach avoids potentially ill-advised practices based on preoccupation with one individual GHG.

To download Holos, for more information, or to access a recent list of Holos related publications, visit:
www.agr.gc.ca

To contact us, email:

aafc.holos.acc@canada.ca