

# Expected Values Estimated via Mean-Field Approximation are $1/N$ -Accurate

Nicolas Gast

Inria, Grenoble, France

Sigmetrics 2017, Urbana-Champaign

# What is mean-field approximation ?

We study a population of  $N$  interchangeable objects.

$\chi^{(N)}$  denotes the empirical measure.

$\chi_i^{(N)}(t) =$  fraction of objects in state  $i$

Idea of mean-field: Some models simplify as  $N \rightarrow \infty$

Theorem (Kurtz 70,...)


$$X^{(N)}(t) \approx x(t)$$

# Idea of mean-field: Some models simplify as $N \rightarrow \infty$


Theorem (Kurtz 70,...)

$$X^{(N)}(t) \approx x(t)$$

Example:  $N$  servers



Randomly choose two, and select one




# Idea of mean-field: Some models simplify as $N \rightarrow \infty$


Theorem (Kurtz 70,...)

$$X^{(N)}(t) \approx x(t)$$

Example:  $N$  servers



Randomly choose two, and select one



# Mean-field approximation is widely used in our community

A few examples of SIGMETRICS papers...

- 2016 **Asymptotics of Insensitive Load Balancing and Blocking Phases** – Jonckheere – Prabhu
- 2016 **On the Approximation Error of Mean-Field Models** – Ying
- 2015 **Power of d Choices for Large-Scale Bin Packing: A Loss Model** – Xie et al
- 2015 **Transient and Steady-state Regime of a Family of List-based Cache Replacement Algorithms** – Gast, Van Houdt
- 2013 **Queueing system topologies with limited flexibility.** – Tsitsiklis, Xu
- 2013 **A mean field model for a class of garbage collection algorithms in flash-based solid state drives.** – Van Houdt
- 2012 **Fluid limit of an asynchronous optical packet switch with shared per link full range wavelength conversion.** – Van Houdt, Bortolussi
- 2011 **On the power of (even a little) centralization in distributed processing.** –
- 2010 **Randomized load balancing with general service time distributions.** – Bramson et al.
- 2010 **Incentivizing peer-assisted services: a fluid shapley value approach.** – Misra et al
- 2010 **A mean field model of work stealing in large-scale systems.** – Gast, Gaujal
- 2009 **The age of gossip: spatial mean field regime.** – Chaintreau et al.


How accurate is mean-field approximation?

# How accurate is mean-field approximation?

Theorem (Kurtz 70... Ying 16)

$$X^{(N)}(t) \approx x(t) + \frac{1}{\sqrt{N}} G_t$$

$X_3(t)$  – Fraction of servers with 3 jobs






# How accurate is mean-field approximation?

Theorem (Kurtz 70... Ying 16)


$$X^{(N)}(t) \approx x(t) + \frac{1}{\sqrt{N}} G_t$$



# How accurate is mean-field approximation?

Theorem (Kurtz 70... Ying 16)

$$X^{(N)}(t) \approx x(t) + \frac{1}{\sqrt{N}} G_t$$



# How accurate is mean-field approximation?

Theorem (Kurtz 70... Ying 16)

$$X^{(N)}(t) \approx x(t) + \frac{1}{\sqrt{N}} G_t$$



# How accurate is mean-field for expected values?

Can we use the approximation for  $N = 1000$ ?  $N = 100$ ?  $N = 10$ ?

$N$	10	100	1000	$\infty$
Average queue length (simulation)	2.8040	2.3931	2.3567	2.3527

Table: Two-choice model with  $\rho = 0.9$

# How accurate is mean-field for expected values?

Can we use the approximation for  $N = 1000$ ?  $N = 100$ ?  $N = 10$ ?

$N$	10	100	1000	$\infty$
Average queue length (simulation)	2.8040	2.3931	2.3567	2.3527
Error of mean-field	0.4513	0.0404	0.0040	0

Table: Two-choice model with  $\rho = 0.9$

## How accurate is mean-field for expected values?

Can we use the approximation for  $N = 1000$ ?  $N = 100$ ?  $N = 10$ ?

$N$	10	100	1000	$\infty$
Average queue length (simulation)	2.8040	2.3931	2.3567	2.3527
Error of mean-field	0.4513	0.0404	0.0040	0

Table: Two-choice model with  $\rho = 0.9$

Contributions :

- 1 We show that under very general conditions, the error is in  $O(1/N)$
- 2 In addition to previous work, the drift needs to be twice-differentiable.
- 3 We study numerically the power-of-two choice.

# Outline

- 1 The Kurtz's Population Model : Classical Convergence Results
- 2 The  $O(1/N)$ -Accuracy of Mean-Field Approximation
- 3 Example: two-choice model
- 4 Recap and Discussion

# Outline

- 1 The Kurtz's Population Model : Classical Convergence Results
- 2 The  $O(1/N)$ -Accuracy of Mean-Field Approximation
- 3 Example: two-choice model
- 4 Recap and Discussion



# CTMC

A continuous-time Markov chain (CTMC) with state-space  $\mathbf{E}$  is given by an initial state  $x_0$  and its transitions ( $\ell \in \mathcal{L}$ ):

$$X \mapsto X + \ell \quad \text{at rate} \quad \beta_\ell(X).$$

The drift is  $f(x) = \sum_{\ell} \ell \beta_\ell(x)$ .

---

<sup>1</sup>We assume  $(\mathbf{E}, \|\cdot\|)$  is a Banach space, not necessarily  $\mathbb{R}^d$ .

# Population CTMC

Density dependent population process (70s)

A population process is a sequence of CTMC  $\mathbf{X}^N$ , indexed by the population size  $N$ , with state spaces  $\mathbf{E}^N \subset \mathbf{E}$ , with initial state  $x_0$  and with transitions (for  $\ell \in \mathcal{L}$ ):

$$X \mapsto X + \frac{\ell}{N} \quad \text{at rate } N\beta_\ell(X).$$

The drift is  $f(x) = \sum_{\ell} \ell \beta_\ell(x)$ .

---

<sup>1</sup>We assume  $(\mathbf{E}, \|\cdot\|)$  is a Banach space, not necessarily  $\mathbb{R}^d$ .

## Transient regime

Let  $\Phi_t$  denotes the (unique) solution of the ODE :

$$\Phi_t x = x + \int_0^t f(\Phi_s x) ds.$$

## Transient regime


Let  $\Phi_t$  denotes the (unique) solution of the ODE :

$$\Phi_t x = x + \int_0^t f(\Phi_s x) ds.$$

### Theorem (Kurtz 70s)

If  $f$  is Lipschitz-continuous with constant  $L$ , then for any fixed  $T$ :

$$\sup_{t < T} \|X^N(t) - \Phi_t X^N(0)\| = O(1/\sqrt{N}) \quad \xrightarrow{N \rightarrow \infty} 0.$$



## Stationary regime

If the ODE  $\dot{x} = f(x)$  has a unique fixed point  $x^*$  that is exponentially stable, then:

### Theorem (Ying 2016)

If  $f$  is Lipschitz-continuous with constant  $L$ , then for any fixed  $T$ :


$$\mathbb{E} \left[ \left\| X^N - x^* \right\| \right] = O(1/\sqrt{N}) \quad \xrightarrow{N \rightarrow \infty} 0.$$

(the uniqueness of the fixed point is not sufficient, see Benaim-Le Boudec 2008).

# Outline


- 1 The Kurtz's Population Model : Classical Convergence Results
- 2 The  $O(1/N)$ -Accuracy of Mean-Field Approximation
- 3 Example: two-choice model
- 4 Recap and Discussion

$1/\sqrt{N}$  or  $1/N$ ?



$N$	10	100	1000	$+\infty$
Average queue length ( $m^N$ )	2.81	2.39	2.36	2.35
Error ( $m^N - m^\infty$ )	0.46	0.039	0.004	0

$1/\sqrt{N}$  or  $1/N$ ?



$N$	10	100	1000	$+\infty$
Average queue length ( $m^N$ )	2.81	2.39	2.36	2.35
Error ( $m^N - m^\infty$ )	0.46	0.039	0.004	0



## Steady-state analysis

$\dot{x} = f(x)$  has an exponentially stable attractor  $x^*$  if for any solution:

$$\|x(t) - x^*\| \leq Ce^{-\alpha t} \|x(0) - x^*\|.$$

# Steady-state analysis

$\dot{x} = f(x)$  has an exponentially stable attractor  $x^*$  if for any solution:

$$\|x(t) - x^*\| \leq Ce^{-\alpha t} \|x(0) - x^*\|.$$

## Theorem

*If  $f$  is twice differentiable, if the ODE has an exponentially stable attractor  $x^*$  and if there exists a bounded set  $\mathcal{B}$  such that  $\mathbf{P}[X^N \notin \mathcal{B}] = O(1/N^2)$ , then for any bounded function  $h$ , there exists a constant  $K$  such that:*

$$\limsup_{N \rightarrow \infty} N \left| \mathbb{E}[h(X^N)] - h(x^*) \right| \leq K.$$

- Note: A similar result holds for the transient behavior.

# Main ideas of the proof

## 1. Comparison of the generators:

$$(L^{(N)}h)(x) = \sum_{\ell \in \mathcal{L}} N\beta_{\ell}(x) \left( h\left(x + \frac{\ell}{N}\right) - h(x) \right)$$

$$(\Lambda h)(x) = \sum_{\ell \in \mathcal{L}} \beta_{\ell}(x) Dh(x) \cdot \ell = Dh(x) \cdot f(x)$$

## 2. Stein's method :

$$\mathbb{E} \left[ h(X^N) - h(x^*) \right] = \mathbb{E} \left[ (\Lambda - L^{(N)})(Gh)X^N \right],$$


where  $Gh(x) = \int_0^{\infty} (h(\Phi_t x) - h(x^*)) dt$  satisfies  $(x) - h(x^*) = \Lambda(Gh)x$ .

## 3. Perturbation theory shows that $D^2(Gh)$ is twice-differentiable.

# Outline

- 1 The Kurtz's Population Model : Classical Convergence Results
- 2 The  $O(1/N)$ -Accuracy of Mean-Field Approximation
- 3 Example: two-choice model
- 4 Recap and Discussion

# The two choice model<sup>2</sup>



Randomly choose two, and select one

Infinite state-space:

$$X_0(t), X_1(t), \dots$$

where

$X_i(t)$  = fraction with  $i$  or more jobs.

---

<sup>2</sup>This model or variants have been heavily studied (Vvedenskaya 96, Mitzenmacher 98 ... Tsitsiklis et al. 2016,...).

# Does this model satisfies our assumptions?

$(\mathbf{E}, \|\cdot\|)$  is the set of infinite sequences such that  $\|x\|_w = \sum_{i=1}^{\infty} w_i |x_i| < \infty$ .

- Transitions : easy
- Regularity of the drift : easy
- Unique attractor : mitzenmacher 98
- Stationary measure concentrates on a bounded set : coupling argument : 2-choice  $\ll$  1-choice.

## The power of two-choice

Our theory guarantees that the average queue length satisfies:

$$m^N(\rho) = m^\infty(\rho) + O(1/N),$$

where  $m^\infty(\rho) \approx \log_2 \frac{1}{1-\rho}$  (as  $\rho \rightarrow 1$ ).

## The power of two-choice

Our theory guarantees that the average queue length satisfies:

$$m^N(\rho) = m^\infty(\rho) + O(1/N),$$

where  $m^\infty(\rho) \approx \log_2 \frac{1}{1-\rho}$  (as  $\rho \rightarrow 1$ ).

By simulation, we observe that  $N(m^N(\rho) - m^\infty) = d(\rho) \approx \frac{\rho^2}{2(1-\rho)}$

$N$	10	20	30	50	$+\infty$
$m^N(\rho)$	2.804	2.566	2.491	2.434	—
$m^\infty(\rho) + \frac{\rho^2}{2N(1-\rho)}$	2.758	2.555	2.488	2.434	2.353

**Table:** Average queue length in the two-choice model ( $\rho = 0.9$ ).



The quality of the approximation degrades as  $\rho$  goes to 1

Simulation results suggest that:

$$m^N(\rho) \approx \underbrace{m^\infty(\rho)}_{\approx \log_2 \frac{1}{1-\rho}} + \frac{1}{N} \underbrace{d(\rho)}_{\approx \frac{\rho^2}{2(1-\rho)}} + O\left(\frac{1}{N^2}\right)$$

Conjecture: the power of two-choice holds if  $N = \Omega\left(\frac{1}{1-\rho}\right)$

# Outline


- 1 The Kurtz's Population Model : Classical Convergence Results
- 2 The  $O(1/N)$ -Accuracy of Mean-Field Approximation
- 3 Example: two-choice model
- 4 Recap and Discussion

# Recap

① Convergence of mean-field model is  $O(1/N)$ .

- ▶ Works for transient and steady-state
- ▶ Works for infinite-dimensional state space.

② Our approach is to focus on the expected values



## In practice

For many mean-field models :

$$\mathbb{E} \left[ X^N \right] \approx x + \frac{C}{N},$$

- We can estimate  $C$  by simulating one value of  $N$  (e.g.  $N = 100$ ).
- We use this  $C$  to deduce the performance for other values of  $N$ .

This provides a new light for the two-choice.

# Does it always work?

- Works for the model of Kurtz
- Also works for the model of *Benaim-Le Boudec 08* by using uniformization

But: it requires the drift to be **twice-differentiable**.

- (see counter-example on the paper)

# Extension and open questions

- Heavy-traffic regime
- Multiple stable equilibria.
- Non-homogeneous population.
  - ▶ e.g., caching

Paper, simulations (and slides) are **reproducible**:

<https://github.com/ngast/meanFieldAccuracy>

# Thank you!

<http://mescal.imag.fr/membres/nicolas.gast>

[nicolas.gast@inria.fr](mailto:nicolas.gast@inria.fr)

## Mean-field and decoupling

Benaïm,  
Le Boudec 08

*A class of mean field interaction models for computer and communication systems*, M.Benaïm and J.Y. Le Boudec., Performance evaluation, 2008.

Le Boudec 10

*The stationary behaviour of fluid limits of reversible processes is concentrated on stationary points.*, J.-Y. L. Boudec. , Arxiv:1009.5021, 2010

Darling Norris 08

*R. W. R. Darling and J. R. Norris*, Differential equation approximations for Markov chains, Probability Surveys 2008

G. 16

*Construction of Lyapunov functions via relative entropy with application to caching*, Gast, N., ACM MAMA 2016

G. 16

*Expected Values Estimated via Mean-field approximation are  $1/N$  accurate*, Gast, N., SIGMETRICS 2017

Budhiraja et al. 15

*Limits of relative entropies associated with weakly interacting particle systems.*, A. S. Budhiraja, P. Dupuis, M. Fischer, and K. Ramanan. , Electronic journal of probability, 20, 2015.

# References (continued)

## Optimal control and mean-field games:

- G.,Gaujal Le Boudec 12 *Mean field for Markov decision processes: from discrete to continuous optimization*, N.Gast,B.Gaujal,J.Y.Le Boudec, IEEE TAC, 2012
- G. Gaujal 12 *Markov chains with discontinuous drifts have differential inclusion limits.*, Gast N. and Gaujal B., Performance Evaluation, 2012
- Lasry Lions *Mean field games*, J.-M. Lasry and P.-L. Lions, Japanese Journal of Mathematics, 2007.
- Tembine et al 09 *Mean field asymptotics of markov decision evolutionary games and teams*, H. Tembine, J.-Y. L. Boudec, R. El-Azouzi, and E. Altman., GameNets 00

## Applications: caches

- Don and Towsley *An approximate analysis of the LRU and FIFO buffer replacement schemes*, A. Dan and D. Towsley., SIGMETRICS 1990
- G. Van Houdt 15 *Transient and Steady-state Regime of a Family of List-based Cache Replacement Algorithms.*, Gast, Van Houdt., ACM Sigmetrics 2015