

Multi-dimensional High-Order FEM Methods for Electron Kinetics in ALE Hydrodynamics

SIAM Conference on Computational Science and Engineering (CSE19) – Spokane, Washington, USA



Milan Holec, Ben S. Southworth

February 25 - March 1, 2019



LLNL-PRES-768124

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344. Lawrence Livermore National Security, LLC

Lawrence Livermore
National Laboratory

1 Motivation - Nonlocal Magneto-Hydrodynamic model (Nonlocal-MHD)

2 S_N discontinuous Galerkin finite element approach to kinetics

3 P_N mixed finite element approach to kinetics

4 Conclusions

Classical MHD

HYDRODYNAMICS *local* $\rightarrow \mathbf{q}_e = -\kappa_{SH} T_e^{2.5} \nabla T_e$

$$\begin{aligned}\frac{d\rho}{dt} &= -\rho \nabla \cdot \mathbf{u}, \\ \rho \frac{d\mathbf{u}}{dt} &= -\nabla(p_i + p_e) + \frac{c}{4\pi} \nabla \times \mathbf{B} \times \mathbf{B}, \\ \rho \left(\frac{\partial \varepsilon_i}{\partial T_i} \frac{dT_i}{dt} + \frac{\partial \varepsilon_i}{\partial \rho} \frac{d\rho}{dt} \right) &= -p_i \nabla \cdot \mathbf{u} - G(T_i - T_e), \\ \rho \left(\frac{\partial \varepsilon_e}{\partial T_e} \frac{dT_e}{dt} + \frac{\partial \varepsilon_e}{\partial \rho} \frac{d\rho}{dt} \right) &= -p_e \nabla \cdot \mathbf{u} + \nabla \cdot (\kappa_{SH} T_e^{2.5} \nabla T_e) + \sigma \mathbf{E} \cdot \mathbf{E} + G(T_i - T_e) + Q_{IB}(\mathbf{E}_L),\end{aligned}$$

MAXWELL EQUATIONS *resistive* $\rightarrow \mathbf{j} = \sigma \mathbf{E}$

$$\begin{aligned}\nabla \times \mathbf{E} &= -\frac{1}{c} \frac{d\mathbf{B}}{dt}, \\ \nabla \times \mathbf{B} &= \frac{4\pi}{c} \sigma \mathbf{E},\end{aligned}$$

KINETICS OF ELECTRONS *Landau – Fokker – Planck*

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla f + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{v}} f = \Gamma \int \nabla_{\mathbf{v}} \nabla_{\mathbf{v}} (\mathbf{v} - \tilde{\mathbf{v}}) \cdot (f \nabla_{\tilde{\mathbf{v}}} f - f \nabla_{\mathbf{v}} f) d\tilde{\mathbf{v}} + \frac{\nu_{ei}}{2} \frac{\partial^2 f}{\partial \Omega^2}.$$

Nonlocal-MHD

HYDRODYNAMICS

4D

$$\begin{aligned}
 \frac{d\rho}{dt} &= -\rho \nabla \cdot \mathbf{u}, \\
 \rho \frac{d\mathbf{u}}{dt} &= -\nabla(p_i + p_e) + \mathbf{j}(f, \mathbf{E}) \times \mathbf{B}, \\
 \rho \left(\frac{\partial \varepsilon_i}{\partial T_i} \frac{dT_i}{dt} + \frac{\partial \varepsilon_i}{\partial \rho} \frac{d\rho}{dt} \right) &= -p_i \nabla \cdot \mathbf{u} - G(T_i - T_e), \\
 \rho \left(\frac{\partial \varepsilon_e}{\partial T_e} \frac{dT_e}{dt} + \frac{\partial \varepsilon_e}{\partial \rho} \frac{d\rho}{dt} \right) &= -p_e \nabla \cdot \mathbf{u} - \nabla \cdot \mathbf{q}_e(f) + \mathbf{j}(f, \mathbf{E}) \cdot \mathbf{E} + G(T_i - T_e) + Q_{IB}(\mathbf{E}_L),
 \end{aligned}$$

MAXWELL EQUATIONS

4D

$$\begin{aligned}
 \nabla \times \mathbf{E} &= -\frac{1}{c} \frac{d\mathbf{B}}{dt}, \\
 \nabla \times \mathbf{B} &= \frac{4\pi}{c} \mathbf{j}(f, \mathbf{E}),
 \end{aligned}$$

KINETICS OF ELECTRONS

6D

$$\mathbf{v} \cdot \nabla f + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_v f = v \tilde{\nu}_e \frac{\partial}{\partial v} (f - f_{MB}(T_e)) + \nu_{scat} (f_0 - f).$$

3D velocity space FEM discretizations

AWBS electron kinetic model 7D

$$\begin{aligned} C_V \frac{d T_e}{dt} &= -\nabla \cdot \mathbf{q}_e(f) + \mathbf{j}(f) \cdot \mathbf{E} + S_H, \\ \mathbf{v} \cdot \nabla f + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{v}} f &= v \tilde{\nu}_e \frac{\partial}{\partial v} (f - f_{MB}(T_e)) + \nu_{scat} (f_0 - f). \end{aligned}$$

- C7 *miniapp* (S_N - discontinuous Galerkin FEM) 3D velocity space in spherical coordinates (v, ϕ, θ)

$$\mathbf{n} \cdot \nabla f + \frac{\mathbf{E} \cdot \mathbf{n}}{v} \frac{\partial f}{\partial v} + \frac{E_\phi - v B_\theta}{v^2} \frac{\partial f}{\partial \phi} + \frac{E_\theta + v B_\phi}{v^2 \sin(\phi)} \frac{\partial f}{\partial \theta} = \tilde{\nu}_e \frac{\partial}{\partial v} (f - f_M) + \frac{\nu_{scat}}{v} (f_0 - f).$$


- AP1 *miniapp* (P_N (VEF $\xi = 1/3$) - continuous Galerkin mixed FEM) *Physicists like it!*

$$\begin{aligned} \xi \nabla \cdot \mathbf{f}_1 + \xi \frac{q_e}{m_e v} \mathbf{E} \cdot \left(\frac{\partial \mathbf{f}_1}{\partial v} + \frac{2}{v} \mathbf{f}_1 \right) &= \tilde{\nu}_e \frac{\partial}{\partial v} (f_0 - f_M), \\ \nabla f_0 + \frac{q_e}{m_e v} \mathbf{E} \frac{\partial f_0}{\partial v} + \frac{q_e \mathbf{B}}{m_e c v} \times \mathbf{f}_1 &= \tilde{\nu}_e \frac{\partial \mathbf{f}_1}{\partial v} - \frac{\nu_{scat}}{v} \mathbf{f}_1, \end{aligned}$$


S_N upwind DG , Multigrid Approximate-Ideal-Relaxation (AIR) solver

$$\mathbf{M}_{(\tilde{\nu}_e - \frac{E \cdot \mathbf{n}_d}{v})} \cdot \frac{\Delta \mathbf{f}_d}{\Delta v} = (\mathbf{n}_d \cdot \mathbf{G} + \mathbf{F}_d) \cdot (\tilde{\mathbf{f}}_d + \Delta \mathbf{f}_d) + \mathbf{M}_{(\frac{\nu_{scat}}{v})} \cdot (\tilde{\mathbf{f}}_d + \Delta \mathbf{f}_d) + \mathbf{S}_{(\nu_{scat}, E, B)} \cdot \tilde{\mathbf{f}} + \mathbf{S}_{(\tilde{\nu}_e \frac{\partial f_M}{\partial v})}$$



Velocity groups	32	64	128	256	order
Backward Euler	1.595e-1	8.618e-2	4.456e-2	2.264e-2	0.98
SDIRK2	3.217e-2	8.888e-3	2.322e-3	5.924e-3	1.97
SDIRK3	2.455e-2	3.639e-3	4.887e-4	6.317e-05	2.96
Hydro temperature	Kinetic temperature (50 groups)				



Local hydro temperature





Nonlocal kinetic temperature




AP1 formulation, fixed P1 angular discretization PCG(AMG)

$$\begin{aligned} \mathbf{M}_{(\nu_e)}^{L_2} \cdot \frac{d\mathbf{f}_0}{dv} - \mathbf{V}_{\left(\frac{\xi q_e E}{m_e v}\right)}^{L_2} \cdot \frac{d\mathbf{f}_1}{dv} &= \mathbf{D}_{(\xi)}^{L_2} \cdot \mathbf{f}_1 + \mathbf{M}_{\left(\frac{\xi^2 q_e E}{m_e v^2}\right)}^{L_2} \cdot \mathbf{f}_1 + \mathbf{S}_{\left(\nu_e \frac{\partial f_M}{\partial v}\right)}^{L_2}, \\ \mathbf{M}_{(\nu_e)}^{H_1} \cdot \frac{d\mathbf{f}_1}{dv} - \mathbf{V}_{\left(\frac{q_e E}{m_e v}\right)}^{H_1} \cdot \frac{d\mathbf{f}_0}{dv} &= \mathbf{G}^{H_1} \cdot \mathbf{f}_0 + \mathbf{M}_{\left(\frac{\nu_{scat}}{v}\right)}^{H_1} \cdot \mathbf{f}_1 + \mathbf{C}_{\left(\frac{q_e B}{m_e c v}\right)}^{H_1} \cdot \mathbf{f}_1, \end{aligned}$$



Electron velocity limit - friction vs. E stopping

$$\left(\tilde{\nu}_e - \frac{\mathbf{E} \cdot \mathbf{n}}{v}\right) \frac{\partial f}{\partial v} = \mathbf{n} \cdot \nabla f + \frac{\nu_{scat}}{v} (f - f_0) + \frac{E_\phi - v B_\theta}{v^2} \frac{\partial f}{\partial \phi} + \frac{E_\theta + v B_\phi}{v^2 \sin(\phi)} \frac{\partial f}{\partial \theta} + \tilde{\nu}_e \frac{\partial f_M}{\partial v}.$$



E stopping overtakes collisions for $Kn > 0.1$.

$Kn = \frac{\lambda}{L}$	10^{-4}	10^{-3}	10^{-2}	10^{-1}	1
v_{lim}/v_{th}	70.8	22.4	7.3	3.1	1.8

Adaptive DSA preconditioner for Hydro-Kinetics coupling

AWBS electron kinetic model 7D

$$\begin{aligned}
 C_V \frac{d T_e}{dt} &= -\nabla \cdot \mathbf{q}_e(f) + \mathbf{j}(f) \cdot \mathbf{E} + S_H, \\
 &= \int_{4\pi} \int_v v \tilde{\nu}_e \frac{\partial f}{\partial v} v^4 dv d\mathbf{n} - \sigma T_e^{-0.5} + S_H, \\
 \mathbf{v} \cdot \nabla f + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_v f &= v \tilde{\nu}_e \frac{\partial}{\partial v} (f - f_{MB}(T_e)) + \nu_{scat} (f_0 - f).
 \end{aligned}$$

Continuum analysis of local ($\text{Kn} \ll 1$) transport regime → DIFFUSION

$$C_V \frac{d T_e}{dt} = \nabla \cdot \lambda(T_e^{2.5}) \nabla T_e + O(\text{Kn}^2)$$

Precodnitioned fixed-point iteration $\mathbf{E}(\Delta \mathbf{T}) \equiv (\mathbf{I} - \mathbf{C}) \cdot \mathbf{M}_\sigma(\tilde{\mathbf{T}} + \Delta \mathbf{T})^\alpha + \mathbf{C} \cdot \mathbf{D}_\lambda(\tilde{\mathbf{T}} + \Delta \mathbf{T})^\alpha$

$$\mathbf{M}_{C_V} \cdot \frac{\Delta \mathbf{T}^{k+1}}{\Delta t} + \mathbf{E}(\Delta \mathbf{T}^{k+1}) - \mathbf{E}(\Delta \mathbf{T}^k) = \mathbf{K}(f^k) - \mathbf{M}_\sigma(\tilde{\mathbf{T}} + \Delta \mathbf{T}^k)^\alpha.$$




Finally, we get an unconditionally stable backward Euler (SDIRK) fast iterating scheme

$$\mathbf{M}_{C_V} \cdot \frac{\Delta \mathbf{T}^{k+1}}{\Delta t} + (\mathbf{I} - \mathbf{C}) \cdot \mathbf{M}_\sigma(\tilde{\mathbf{T}} + \Delta \mathbf{T}^{k+1})^\alpha + \mathbf{C} \cdot \mathbf{D}_\lambda(\tilde{\mathbf{T}} + \Delta \mathbf{T}^{k+1})^\alpha = \mathbf{C} \cdot \mathbf{D}_\lambda(\tilde{\mathbf{T}} + \Delta \mathbf{T}^k)^\alpha + \mathbf{K}(f^k) - \mathbf{C} \cdot \mathbf{M}_\sigma(\tilde{\mathbf{T}} + \Delta \mathbf{T}^k)^\alpha,$$

where adaptive coefficient diffusion $\mathbf{C} \xrightarrow{\text{Kn} \ll 1} 1$ and nonlocal transport $\mathbf{C} \xrightarrow{\text{Kn} > 1} 0$ and $\mathbf{C} \in (1, 0)$ in between.

Conclusions

- 7D microscopic world of electrons in hydro simulations.
- S_N high-order DG finite element approach.
- P_1 high-order mixed finite element approach.
- E field dominated stopping (P_1 fails).
- Adaptive DSA preconditioner for Hydro-Kinetics coupling (deep learning on \mathbf{C}).
- Algebraic Multigrid solver pAIR scales $\log(P)^{1.22}$.



MS85 - Developments in Algebraic Multigrid for Nonsymmetric and Hyperbolic Problems - Ben S. Southworth
<https://github.com/CEED/Laghos/tree/master/amr>
<https://computation.llnl.gov/projects/blast>
<https://mfem.org>
holec1@llnl.gov

Thank you for your attention. Any questions?



CASC

Center for Applied
Scientific Computing

Disclaimer: This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees make any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represent that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and options of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.