




AWBS kinetic modeling of electrons with nonlocal Ohms law in plasmas relevant to inertial confinement fusion

Milan Holec, P. Loiseau, J. P. Brodrick, D. Del Sorbo, A. Debayle, V. Tikhonchuk, J.-L. Feugeas,
Ph. Nicolai, B. Dubroca, C. P. Ridgers, R. J. Kingham



LLNL - ClubMED,
Livermore, CA,
Jan 7, 2019

Hydrodynamic model of plasma

Fokker-Planck/Boltzmann transport equation

$$\frac{\partial \mathbf{f}}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} \mathbf{f} + \frac{q_e}{m_e} \left(\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B} \right) \cdot \nabla_{\mathbf{v}} \mathbf{f} = \sigma \nabla_{\mathbf{v}} \cdot \int \frac{|\mathbf{v} - \mathbf{v}'|^2 \mathbf{l} - (\mathbf{v} - \mathbf{v}') \otimes \mathbf{v} - \mathbf{v}'}{|\mathbf{v} - \mathbf{v}'|^3} (\nabla_{\mathbf{v}} \mathbf{f}(\mathbf{v}) f(\mathbf{v}') - \mathbf{f}(\mathbf{v}) \nabla_{\mathbf{v}} f(\mathbf{v}')) d\mathbf{v}'. \quad (1)$$

Fluid equations in Lagrangian frame (velocity space moments of (1))

$$\begin{aligned}\frac{d\rho}{dt} &= -\rho \nabla \cdot \mathbf{U}, \\ \rho \frac{d\mathbf{U}}{dt} &= \nabla \cdot \sigma + \mathbf{j} \times \mathbf{B}, \\ \rho \frac{d\varepsilon}{dt} &= \sigma : \nabla \mathbf{U} - \nabla \cdot \mathbf{q}_e.\end{aligned}$$

Microscopic closure

$$\begin{aligned}\sigma &= -\rho \int (\mathbf{v} - \mathbf{U}) \otimes (\mathbf{v} - \mathbf{U}) \mathbf{f} d\mathbf{v}^3 \approx -\mathbf{l}\rho + \tilde{\sigma}(\nabla \mathbf{U}), \\ \mathbf{q}_e &= \frac{m_e}{2} \int |\mathbf{v} - \mathbf{U}|^2 (\mathbf{v} - \mathbf{U}) \mathbf{f} d\mathbf{v}^3 \approx \frac{m_e}{2} \int_{4\pi} \mathbf{n} \int_0^\infty |\mathbf{v}|^5 \mathbf{f} d|\mathbf{v}| d\mathbf{n} \approx -\kappa(T^{2.5}) \nabla T, \\ \mathbf{j} &= q_e \int (\mathbf{v} - \mathbf{U}) \mathbf{f} d\mathbf{v}^3.\end{aligned}$$

Local diffusive transport regime


Collision operators

$$C_{FP} = \Gamma_{ee} \int \nabla_v \nabla_v (\mathbf{v} - \tilde{\mathbf{v}}) \cdot (f \nabla_{\tilde{\mathbf{v}}} f - f \nabla_v f) d\tilde{\mathbf{v}} + \frac{\nu_{ei}}{2} \frac{\partial}{\partial \mu} \left((1 - \mu^2) \frac{\partial f}{\partial \mu} \right), \quad (2)$$

$$C_{BGK} = \nu_e (f_M - f) + \frac{Z + 4.2}{Z + 0.24} \frac{\nu_{ei}}{2} \frac{\partial}{\partial \mu} \left((1 - \mu^2) \frac{\partial f}{\partial \mu} \right), \quad (3)$$







$$C_{FP0} = \nu \nu_e \frac{\partial}{\partial v} \left[C(f_0) f_0 + D(f_0) \frac{\partial f_0}{\partial v} \right] + \frac{Z + 4.2}{Z + 0.24} \frac{\nu_{ei}}{2} \frac{\partial}{\partial \mu} \left((1 - \mu^2) \frac{\partial f}{\partial \mu} \right), \quad (4)$$

$$C_{AWBS} = \nu \frac{\nu_e}{2} \frac{\partial}{\partial v} (f - f_M) + \frac{\nu_{ei} + \frac{\nu_e}{2}}{2} \frac{\partial}{\partial \mu} \left((1 - \mu^2) \frac{\partial f}{\partial \mu} \right). \quad (5)$$



	$Z = 1$	$Z = 2$	$Z = 4$	$Z = 16$	$Z = 116$
Δq_{AWBS}	0.057	0.004	0.037	0.021	0.004


Nonlocal electron transport regime - PIC/VFP0/AWBS



Electron velocity limit - friction vs. \mathbf{E} stopping


$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla f + \frac{q_e}{m_e} \left(\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B} \right) \cdot \frac{\partial f}{\partial \Omega} = \left(v \frac{\nu_e}{2} - \frac{q_e}{m_e} \mathbf{v} \cdot \mathbf{E} \right) \frac{\partial f}{\partial v} - v \frac{\nu_e}{2} \frac{\partial f_M}{\partial v} + \frac{\nu_{ei} + \frac{\nu_e}{2}}{2} \frac{\partial^2 f}{\partial \Omega^2}.$$

Kn^e	10^{-4}	10^{-3}	10^{-2}	10^{-1}	1
v_{lim}/v_{th}	70.8	22.4	7.3	3.1	1.8



\mathbf{E} stopping overtakes collisions for $\text{Kn}>0.1$.

NIF hohlraum - plasma profiles provided by HYDRA



SNB approach

$$\tilde{f} = f_M + \delta f_0 + \Omega \cdot (\mathbf{f}_{1M} + \delta \mathbf{f}_1),$$

$$\mathbf{f}_{1M} = -\frac{Z+0.24}{Z+4.2} \lambda_{ei} f_M \left(\frac{\nabla n_e}{n_e} + \left(\frac{v^2}{2v_{th}^2} - \frac{3}{2} \right) \frac{\nabla T_e}{T_e} - \frac{q_e \mathbf{E}_L}{m_e v_{th}^2} \right), \quad \mathbf{E}_L = \frac{m_e v_{th}^2}{q_e} \left(\frac{\nabla n_e}{n_e} + \frac{5}{2} \frac{\nabla T_e}{T_e} \right).$$

Nonlocal Ohm's law

We found that the quality of the electric field evaluation is essential for a correct plasma modeling

Nonlocal current in plasma

$$j(f, \mathbf{E}) = e \int v \mathbf{f}_1 v^2 dv = e \int v \frac{\nu_{ei}^2 \mathbf{E}^* + \omega_B \omega_B \cdot \mathbf{E}^* + \nu_{ei} \omega_B \times \mathbf{E}^*}{\nu_{ei}(\omega_B^2 + \nu_{ei}^2)} v^2 dv, \quad (6)$$

where $\mathbf{E}^* = v \frac{\nu_e}{2} \frac{\partial \mathbf{f}_1}{\partial v} - v \nabla f_0 - \frac{q_e}{m_e} \mathbf{E} \frac{\partial f_0}{\partial v}$ is an effective electric field in plasma. A comparison to the *Generalized Ohm's law* shows a correct local behavior of (6), especially that $\nabla \times \nabla f_0 \sim \nabla \times \frac{\nabla p_e}{n_e} \sim \nabla n_e \times \nabla T_e$

Generalized Ohm's law vs. nonlocal Ohm's law

$$\begin{aligned} \mathbf{E} &= (\mathbf{R}_T - \nabla p_e) + \frac{\mathbf{j}}{\sigma} + \mathbf{j} \times \mathbf{B}, \\ \mathbf{E} v \frac{q_e}{m_e} \frac{\partial f_0}{\partial v} &= v^2 \nabla f_0 + v^2 \frac{\nu_e}{2} \frac{\partial \mathbf{f}_1}{\partial v} - v \nu_{ei} \mathbf{f}_1 + v \mathbf{f}_1 \times \frac{q_e}{m_e c} \mathbf{B}. \end{aligned}$$

Nonlocal-MHD

HYDRODYNAMICS

$$\begin{aligned}\frac{d\rho}{dt} &= -\rho \nabla \cdot \mathbf{u}, \\ \rho \frac{d\mathbf{u}}{dt} &= -\nabla(p_i + p_e) + \mathbf{j}(f, \mathbf{E}) \times \mathbf{B}, \\ \rho \left(\frac{\partial \varepsilon_i}{\partial T_i} \frac{dT_i}{dt} + \frac{\partial \varepsilon_i}{\partial \rho} \frac{d\rho}{dt} \right) &= -p_i \nabla \cdot \mathbf{u} - G(T_i - T_e), \\ \rho \left(\frac{\partial \varepsilon_e}{\partial T_e} \frac{dT_e}{dt} + \frac{\partial \varepsilon_e}{\partial \rho} \frac{d\rho}{dt} \right) &= -p_e \nabla \cdot \mathbf{u} - \nabla \cdot \mathbf{q}_e(f) + G(T_i - T_e) + Q_{IB}(\mathbf{E}_L),\end{aligned}$$

MAXWELL EQUATIONS

$$\begin{aligned}\nabla \times \mathbf{E} &= -\frac{1}{c} \frac{d\mathbf{B}}{dt}, \\ \nabla \times \mathbf{B} &= \frac{4\pi}{c} \mathbf{j}(f, \mathbf{E}),\end{aligned}$$

AWBS KINETICS OF ELECTRONS

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla f + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{v}} f = \mathbf{v} \frac{\nu_e}{2} \frac{\partial}{\partial \mathbf{v}} (f - f_{MB}(T_e)) + \frac{\nu_{ei} + \frac{\nu_e}{2}}{2} \frac{\partial^2 f}{\partial \Omega^2}.$$

Nonlocal-MHD \iff aiming HIGH



Height 250 m, overhang 40 m, Visera, Riglos, Spain.

Nonlocal-MHD \iff aiming HIGH



Height 250 m, overhang 40 m, Visera, Riglos, Spain.

Nonlocal-MHD \iff aiming HIGH



Height 250 m, overhang 40 m, Visera, Riglos, Spain.