

# AWBS kinetic modeling of electrons with nonlocal Ohm's law in plasmas relevant to inertial confinement fusion

M. Holec\*

*Center for Applied Scientific Computing, Lawrence Livermore National Laboratory,  
P.O. Box 808, L-561, Livermore, CA 94551, USA. and  
Centre Lasers Intenses et Applications,  
Universite de Bordeaux-CNRS-CEA,  
UMR 5107, F-33405 Talence, France.*

P. Loiseau and A. Debayle  
*CEA, DAM, DIF, F-91297 Arpajon Cedex, France.*

J. P. Brodrick, D. Del Sorbo, and C. P. Ridgers  
*York Plasma Institute, Department of Physics, University of York,  
Heslington, York, YO10 5DD, UK.*

V. Tikhonchuk, J.-L. Feugeas, Ph. Nicolai, and B. Dubroca  
*Centre Lasers Intenses et Applications,  
Universite de Bordeaux-CNRS-CEA,  
UMR 5107, F-33405 Talence, France.*

R. J. Kingham  
*Plasma Physics Group, Blackett Laboratory, Imperial College,  
London SW7 2BW, United Kingdom.*  
(Dated: December 5, 2018)

The interaction of lasers with plasmas very often leads to nonlocal transport conditions, where the classical hydrodynamic model fails to describe important microscopic physics related to highly mobile particles. In this study we analyze and further propose a modification of the Albritton-Williams-Bernstein-Swartz collision operator [1] previously used in [2] for the nonlocal electron transport under conditions relevant to ICF. The electron distribution function provided by this modification exhibits some very desirable properties when compared to the full Fokker-Planck operator in the local diffusive regime, and also performs very well when benchmarked against Vlasov-Fokker-Planck and collisional PIC codes in the nonlocal transport regime, where we find that the effect of the electric field via the nonlocal Ohm's law is an essential ingredient in order to capture the electron kinetics properly.

## I. INTRODUCTION

The first modern attempts at kinetic modeling of plasma can be traced back to the fifties, when Cohen, Spitzer, and Routly (CSR) [3] demonstrated that the effect of Coulomb collisions between electrons and ions in the ionized gas predominantly results from frequently occurring events of cumulative small deflections rather than occasional close encounters. This effect was originally described by Jeans in [4] and Chandrasekhar [5] proposed to use the diffusion equation model of the Vlasov-Fokker-Planck type (VFP) [6].

A classical paper by Spitzer and Harm (SH) [7] provides the computation of the electron distribution function (EDF) in a plasma (from low to high  $Z$ ) with a temperature gradient accounting for e-e and e-i collisions. The resulting expressions for current and heat flux are

widely used in plasma hydrodynamic models. The distribution function based on the spherical harmonics method in its first approximation (P1) [8] is of the form  $f^0 + \mu f^1$ , where  $f^0$  and  $f^1$  are isotropic and  $\mu$ , is the direction cosine between the particle trajectory and some preferred direction in space. It should be emphasized that the SH solution expresses a small perturbation of equilibrium, i.e. that  $f^0$  is the Maxwell-Boltzmann distribution and  $\mu f^1$  represents a very small anisotropic deviation. This approximation holds for  $L_T \gg \lambda_e$ , a condition which is often invalid in laser plasmas, where  $L_T$  is the temperature length scale and  $\lambda_e$  the mean free path of electrons. It is worth mentioning, that electrons having 3 to 4 times the thermal velocity are dominantly responsible for heat-flow and that those faster than 6 times the thermal velocity can be completely neglected in this local theory.

The actual cornerstone of the modern VFP simulations was set in place by Rosenbluth [9], when he derived a simplified form of the VFP equation for a finite expansion of the distribution function, where all the terms are computed according to plasma conditions, includ-

---

\* holec1@llnl.gov

ing  $f^0$ , which of course needs to tend to the Maxwell-Boltzmann distribution. Consequently, the pioneering work on numerical solution of the VFP equation [10, 11] revealed the importance of the nonlocal electron transport in laser-heated plasmas. In particular, that the heat flow down steep temperature gradients in unmagnetised plasma cannot be described by the classical, local fluid description of transport [7, 12]. This is due to the classical  $f^1$  is not a small deviation (especially for electrons having 3 to 4 times the thermal velocity), i.e.  $f^0 \sim f^1$  characterized by  $L_T \sim \lambda_e$ . It was also shown that a thermal transport inhibition [10] around the peak of the temperature gradient, and a nonlocal preheat ahead of the main heat wave front, naturally appear. These effects are attributed to significant deviations of  $f^0$  from Maxwellian distribution.

Nevertheless, numerical solution of the VFP equation even in the Rosenbluth formalism remains very challenging computationally, because the e-e collision integral is nonlinear. More simple linear forms of e-e collision operator are needed. Although some VFP simulations on experimentally relevant timescales have been performed (for recent examples see [13–19], an extensive review has been conducted by Thomas et al. [20]), their relative computational inefficiency severely limits the range of simulations that can be performed.

It is the purpose of this paper to present an efficient alternative to a full solution of the VFP equation to accurately calculate nonlocal transport, based on the Albritton-Williams-Bernstein-Swartz collision operator (AWBS) [1]. In Section II we propose a modified form of the AWBS collision operator, where its important properties are further presented in Section III with the emphasis on its comparison to the full VFP solution in local diffusive regime. In Section IV we define a full model of electron kinetics and the way of discretizing the electron phase-space and also the coupling of the kinetic model to magneto-hydrodynamics. Section V focuses on the performance of the AWBS transport equation model compared to modern kinetic codes including VFP codes Aladin and Impact [21], and PIC code Calder [22], where the cases related to real laser generated plasma conditions are studied. Finally, the most important outcomes of our research are concluded in Section VI.

## II. THE AWBS KINETIC MODEL

The electrons in plasma can be modeled by the deterministic Vlasov model of charged particles

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f + \frac{q_e}{m_e} \left( \mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B} \right) \cdot \nabla_{\mathbf{v}} f = C_{ee}(f) + C_{ei}(f), \quad (1)$$

where  $f(t, \mathbf{x}, \mathbf{v})$  represents the density function of electrons at time  $t$ , spatial point  $\mathbf{x}$ , and velocity  $\mathbf{v}$ ,  $\mathbf{E}$  and  $\mathbf{B}$  are the electric and magnetic fields in plasma,  $q_e$  and  $m_e$  being the charge and mass of electron.

The general form of the e-e collision operator  $C_{ee}$  is the Fokker-Planck form published by Landau [23]

$$C_{FP}(f) = \Gamma \int \nabla_{\mathbf{v}} \nabla_{\mathbf{v}'} (\mathbf{v} - \mathbf{v}') \cdot (f \nabla_{\mathbf{v}'} f - f' \nabla_{\mathbf{v}} f) d\mathbf{v}', \quad (2)$$

where  $\Gamma = \frac{4\pi q_e^4 \ln \Lambda}{m_e^2}$  and  $\ln \Lambda$  is the Coulomb logarithm. The e-i collision operator  $C_{ei}$  could be expressed in a simpler form since massive ions are considered to be motionless compared to electrons during a collision. The scattering operator accounts for the change of electron velocity without change in the velocity magnitude, i.e. angular scattering. It is expressed in spherical coordinates as

$$C_{ei}(f) = \frac{\nu_{ei}}{2} \left( \frac{\partial}{\partial \mu} \left( (1 - \mu^2) \frac{\partial f}{\partial \mu} \right) + \frac{1}{1 - \mu^2} \frac{\partial^2 f}{\partial \theta^2} \right), \quad (3)$$

where  $\mu = \cos \phi$ ,  $\phi$  and  $\theta$  are the polar and azimuthal angles, and  $\nu_{ei} = \frac{Z n_e \Gamma}{v^3}$  is the e-i collision frequency.

The e-e collision operator needs to be linearized for efficient computation. Fisch introduced a linear form of  $C_{ee}$  in [24] in the high-velocity limit ( $v \gg v_{th}$ ) electron collision operator

$$C_H(f) = \nu_e \frac{\partial}{\partial v} \left( f + \frac{v_{th}^2}{v} \frac{\partial f}{\partial v} \right) + \frac{\nu_e}{2} \left( 1 - \frac{v_{th}^2}{2v^2} \right) \left( \frac{\partial}{\partial \mu} \left( (1 - \mu^2) \frac{\partial f}{\partial \mu} \right) + \frac{1}{1 - \mu^2} \frac{\partial^2 f}{\partial \theta^2} \right), \quad (4)$$

where  $\nu_e = \frac{n_e \Gamma}{v^3}$  is the e-e collision frequency and  $v_{th} = \sqrt{\frac{k_B T_e}{m_e}}$  is the electron thermal velocity. The linear form of  $C_H$  arises from an assumption that the fast electrons predominantly interact with the thermal (slow) electrons, which is an important simplification to the form (2). However the diffusion term in the e-e collision operator (4) still presents numerical difficulties.

A yet simpler form of the collision operator of electrons was proposed in [2]

$$C_{AWBS}(f) = \nu_e^* \frac{\partial}{\partial v} (f - f_M) + \frac{\nu_{ei} + \nu_e^*}{2} \left( \frac{\partial}{\partial \mu} \left( (1 - \mu^2) \frac{\partial f}{\partial \mu} \right) + \frac{1}{1 - \mu^2} \frac{\partial^2 f}{\partial \theta^2} \right), \quad (5)$$

where  $f_M = \frac{n_e}{(2\pi)^{3/2} v_{th}^3} \exp\left(-\frac{v^2}{2v_{th}^2}\right)$  is the Maxwell-Boltzmann equilibrium distribution. Here, the first term representing the AWBS operator [1] accounts for relaxation to equilibrium due to the e-e collisions, and the second term accounts for the e-i and e-e collisions contribution to scattering.

A method of angular momenta for the solution of the electron kinetic equation with the collision operator (5) was introduced in [2, 25].

In (5) we have introduced a modified e-e collision frequency  $\nu_e^*$  in order to address a proper behavior with respect to  $Z$ , which is further analyzed in Section III and promising results compared to the full FP operator are presented.

### III. BGK, AWBS, AND FOKKER-PLANCK MODELS IN LOCAL DIFFUSIVE REGIME

An approximate solution to the so-called *local diffusive regime* of electron transport can be found, since the *diffusive regime* refers to a low anisotropy given by the projection  $\mu$ . i.e. modeled by the P1 form of EDF

$$\tilde{f}(z, v, \mu) = f^0(z, v) + \mu f^1(z, v), \quad (6)$$

where  $z$  is the spatial coordinate along the axis  $z$ ,  $v$  the magnitude of transport velocity.

The approximate transport solution is then obtained when analyzing the action of the time-steady form of (1) in 1D on the approximation (6) as

$$\mu \left( \frac{\partial \tilde{f}}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial \tilde{f}}{\partial v} \right) + \frac{q_e E_z}{m_e} \frac{(1 - \mu^2)}{v^2} \frac{\partial \tilde{f}}{\partial \mu} = \frac{1}{v} C(\tilde{f}), \quad (7)$$

where  $C$  is a given collision operator including both e-e and e-i collisions.

The locality of transport is the best expressed in terms of the Knudsen number  $Kn = \frac{\lambda}{L}$ , where  $\lambda$  is the mean free path of electron and  $L$  the characteristic length scale of plasma. Consequently, plasma conditions characterized by  $Kn \ll 1$  exhibit a local transport regime. This measure then play a very important role in our analysis, where we use the electron-electron and electron-ion mean free paths  $\lambda_e = Z\lambda_{ei} = \frac{v}{\nu_e}$ , and the density and temperature plasma scale lengths  $L_{n_e} = n_e / \frac{\partial n_e}{\partial z}$  and  $L_{T_e} = T_e / \frac{\partial T_e}{\partial z}$ .

In practice, the Knudsen number of thermal electrons is often used as a measure of the locality of transport corresponding to given plasma conditions, where  $Kn(v_{th}) < 0.001$  is considered the limit of validity of the local transport theory [26].

#### A. The BGK local diffusive electron transport

Bhatnagar, Gross, and Krook introduced a very simple form of a collision operator [27]

$$C_{BGK}(\tilde{f}) = \nu_e(f_M - \tilde{f}) + \frac{\nu_{ei} + \nu_e}{2} \frac{\partial}{\partial \mu} (1 - \mu^2) \frac{\partial \tilde{f}}{\partial \mu}. \quad (8)$$

In spite of its simple form, BGK collision operator (8) serves as a useful model providing a relevant kinetic response, yet only qualitative with respect to the FP collision operator (2). In particular, the conservation of kinetic energy, momentum, and number of particles is often violated [28].

However, the form of (8) provides a simple analytical treatment of local diffusive transport regime, when used in (7). As a result, one finds a simple form of the BGK isotropic and anisotropic terms of (6) to be

$$f^0 = f_M, \quad (9)$$

$$f^1 = -\frac{\lambda_e}{Z+2} \left( \frac{\partial f_M}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial f_M}{\partial v} \right), \quad (10)$$

where a detailed derivation of (9) and (10) can be found in Appendix A. When the quasi-neutrality constraint imposed by  $\mathbf{E}_L$  (A8) is used, one finally obtains the analytical BGK form of the anisotropic term

$$f^1 = -\mu \left( \frac{v^2}{2v_{th}^2} - 4 \right) \frac{1}{Z+2} \frac{\lambda_e}{L_{T_e}} f_M. \quad (11)$$

The details about the BGK distribution function compared to other collision operators can be found in Section III D.

#### B. The AWBS local diffusive electron transport

Similarly to the BGK model, the AWBS collision operator 5 explicitly uses equilibration to the Maxwell-Boltzmann distribution  $f_M$ . On the other hand, AWBS originates from  $C_H$ , which is derived from the full FP operator (2). This makes the AWBS operator to be superior to the BGK operator, which is considered a purely phenomenological model.

If (5) is used in (7), one obtains the following equations governing the AWBS isotropic and anisotropic terms of (6)

$$\frac{\partial f^0}{\partial v} = \frac{\partial f_M}{\partial v}, \quad (12)$$

$$\frac{\partial f^1}{\partial v} - \frac{Z + r_A}{v r_A} f^1 = \frac{\lambda_e}{v r_A} \left( \frac{\partial f_M}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial f_M}{\partial v} \right), \quad (13)$$

where  $r_A$  represents a scaling parameter defining the modified e-e collision frequency as  $\nu_e^* = r_A \nu_e$ . A detailed derivation of (12) and (13) can be found in Appendix A. Consequently, one finds the AWBS model equation for  $f^1$  in local diffusive regime to be

$$\begin{aligned} \frac{\partial f_{AWBS}^1}{\partial v} + \frac{Z + r_A}{v r_A} f_{AWBS}^1 = \\ \frac{\lambda_e}{v r_A} \left( \frac{1}{L_{n_e}} + \left( \frac{v^2}{2v_{th}^2} - \frac{3}{2} \right) \frac{1}{L_{T_e}} - \frac{q_e E_z}{m_e v_{th}^2} \right) f_M. \end{aligned} \quad (14)$$

which can be solved analytically by using quadratures

$$\text{Vladimir's analytical formula.} \quad (15)$$

The details about the AWBS distribution function compared to other collision operators and a proper evaluation of the scaling parameter  $r_A$  can be found in Section III D.

#### C. The Fokker-Planck local diffusive electron transport

The solution to the 1D transport equation (7) using the Fokker-Planck collision operator (2) is very ambitious, as demonstrated in [3, 5, 9], fortunately, one can

use the explicit evaluation of the electron distribution function published in [7], which takes the following form

$$f_{SH}^1 = \frac{v_{2th}^4}{\Gamma Z n_e} \left( 2\tilde{D}_T \left( \frac{v}{v_{2th}} \right) + \frac{3}{2} \frac{\gamma_T}{\gamma_E} \tilde{D}_E \left( \frac{v}{v_{2th}} \right) \right) \frac{f_M}{T} \frac{\partial T_e}{\partial z}, \quad (16)$$

where  $\tilde{D}_T(x) = Z D_T(x)/B$ ,  $\tilde{D}_E(x) = Z D_E(x)/A$ ,  $\gamma_T$ , and  $\gamma_E$  are numerical values in TABLE I, TABLE II, and TABLE III in [7], and  $v_{2th} = \sqrt{\frac{k_B T_e}{2m_e}}$ .

One should be aware, that the solution of (7) equipped with the full FP collision operator reveals the importance of e-e Coulomb collisions, which is emphasized in the  $Z$  dependence of the distribution function, current, heat flux, electric field, etc. In particular, the latter exhibits the following dependence [7]

$$\mathbf{E} = \frac{m_e v_{th}^2}{q_e} \left( \frac{\nabla n_e}{n_e} + \left( 1 + \frac{3}{2} \frac{Z + 0.477}{Z + 2.15} \right) \frac{\nabla T_e}{T_e} \right), \quad (17)$$

which for  $Z \gg 1$  corresponds to the classical Lorentz electric field (A8).

#### D. Summary of the BGK, AWBS, and Fokker-Planck local diffusive transport

Ever since the SH paper [7], the effect of microscopic electron transport on the current  $\int q_e \mathbf{v} \tilde{f} d\mathbf{v}$  and the heat flux  $\int \frac{m_e |\mathbf{v}|^2}{2} \mathbf{v} \tilde{f} d\mathbf{v}$  in plasmas under local diffusive conditions has been understood. By overcoming some delicate aspects of the numerical solution to (2) presented in the CSR paper [3], the effect of electron-electron collisions was properly quantified and the correct dependence on  $Z$  of the heat flux  $\mathbf{q}$  was approximated as [7]

$$\mathbf{q} = \xi(Z) \mathbf{q}_L = \frac{Z + 0.24}{Z + 4.2} \mathbf{q}_L, \quad (18)$$

where  $\xi$  is the  $Z$ -dependence [29] approximation and  $\mathbf{q}_L$  is the heat flux given by the Lorentz gas model [30]

$$f_{Lorentz}^1 = -\mu \left( \frac{v^2}{2v_{th}^2} - 4 \right) \frac{\lambda_{ei}}{L_{Te}} f_M. \quad (19)$$

In the case of BGK the collision operator (8) needs to be corrected in order to provide a same local behavior as (18), i.e. a correct dependence on  $Z$ . Consequently, we define a scaling formula

$$r_B(Z) = \frac{\zeta Z}{\xi(Z + 2\zeta)}, \quad (20)$$

based on comparison of the formula (11) to  $\xi f_{Lorentz}^1$  and we write a consistent local diffusion version of BGK

$$C_{BGK}(\tilde{f}) = r_B \nu_e (f_M - \tilde{f}) + \frac{r_B}{\zeta} \frac{\nu_{ei} + \zeta \nu_e}{2} \frac{\partial}{\partial \mu} (1 - \mu^2) \frac{\partial \tilde{f}}{\partial \mu}, \quad (21)$$

	$Z = 1$	$Z = 2$	$Z = 4$	$Z = 16$	$Z = 116$
$\bar{\Delta} \mathbf{q}_{AWBS}$	0.057	0.004	0.037	0.021	0.004
$\phi(Z)$	-0.037	-0.003	0.04	0.058	0.065

TABLE I. Relative error  $\bar{\Delta} \mathbf{q}_{AWBS} = |\mathbf{q}_{AWBS} - \mathbf{q}_{SH}|/|\mathbf{q}_{SH}|$  of the  $\nu_e^* = \frac{\nu_e}{2}$  scaling used in the AWBS model (5) showing the discrepancy (maximum 6%) with respect to the original solution of the heat flux given by numerical solution in Spitzer and Harm [7]. The values of  $\phi(Z)$  (a weak dependence (23)) are also shown.

where the constant  $\zeta$  can be set arbitrarily, because it does not affect the local EDF of (21)

$$f_{BGK}^1 = -\mu \left( \frac{v^2}{2v_{th}^2} - 4 \right) \frac{\zeta}{r_B(Z)} \frac{1}{Z + 2\zeta} \frac{\lambda_e}{L_{Te}} f_M, \quad (22)$$

which is identical to  $\xi f_{Lorentz}^1$  for any value of  $\zeta$ . One should notice that  $r_B(Z \gg 1) = \zeta$ , i.e.  $\zeta$  can be adjusted appropriately for example to better address the transport in nonlocal regime.

We have performed an extensive analysis in the case of the AWBS operator in order to obtain the heat flux behavior while varying  $Z$ . As expected, the heat flux magnitude did not match exactly the  $Z$ -dependence (18), e.g. for  $Z = 1$  the AWBS heat flux was about 60% less than the SH calculation, while there was a perfect match in the case of  $Z \gg 1$ . By assuming that the e-e collisions are responsible for this inadequacy, we searched for a scaling of  $\nu_e$  in (5). Interestingly, we found an almost constant scaling  $r_A$ , i.e. with a very weak dependence on  $Z$  as

$$\nu_e^* = r_A(Z) \nu_e = \left( \frac{1}{2} + \phi(Z) \right) \nu_e \approx \frac{\nu_e}{2}, \quad (23)$$

where can be approximated as  $\phi(Z) = \frac{0.59Z - 1.11}{8.37Z + 5.15} \ll \frac{1}{2}$  for any  $Z$ , i.e. we decide to use  $r_A = \frac{1}{2}$ . Indeed, TABLE I shows  $\phi(Z)$  and corresponding relative error (maximum 6%) of the heat flux modeled by (5) vs. SH results represented by (18). It should be noted that the error is calculated with respect to original values presented in TABLE III in [7]. It is worth mentioning, that the zero current condition was imposed in all AWBS calculations, where the consistent electric field yield an equal value to the classical Lorentz value (A8) for any  $Z$ .

Nevertheless, the electron-electron collisions scaling [29] represented by (18) provides only an integrated information about the heat flux magnitude. If one takes a closer look at the distribution function itself, the conformity of the modified AWBS collision operator is even more emphasized as can be seen in FIG. 1 showing

the flux moment in spherical coordinates of velocity

$$q_1 = \frac{m_e v^2}{2} v f^1 v^2. \quad (24)$$

In the case of the high  $Z$  plasma ( $Z = 116$ ), AWBS exactly aligns with the Lorentz gas limit (19). In the opposite case of the low  $Z$  Hydrogen plasma ( $Z = 1$ ), the AWBS distribution function approaches closely the numerical SH solution (16). BGK (21) takes the Lorentz gas distribution function for any  $Z$  only scaled by  $\xi$ . The AWBS collision operator (5) (red dashed line) represented by a derivative with respect to  $v$  provides a significant improvement with respect to the SH (Fokker-Planck) solution (16) (solid black line) compared to the simplest BGK model (22) (dashed-dot blue line) in FIG. 1.




FIG. 1. The flux velocity moment of the anisotropic part of the electron distribution function in low  $Z = 1$  and high  $Z = 116$  plasmas in diffusive regime. In the case of  $Z = 1$  the AWBS model matches very well the reference solution given by the SH calculation [7] in comparison to the BGK model. In the case of  $Z = 116$  the AWBS model aligns exactly with the Lorentz gas approximation as expected. The BGK and the SH curves are not shown for  $Z = 116$ , but also correspond to the Lorentz gas distribution function.

#### IV. THE AWBS NONLOCAL TRANSPORT MODEL OF ELECTRONS

In order to define a nonlocal transport model of electrons, we use the AWBS collision operator and the P1 angular discretization of the electron distribution function

$$\tilde{f}(\mathbf{x}, \mathbf{n}, v) = f_0(\mathbf{x}, v) + \mathbf{n} \cdot \mathbf{f}_1(\mathbf{x}, v), \quad (25)$$

consisting of the isotropic part represented by the zeroth angular moment  $f_0 = \frac{1}{4\pi} \int_{4\pi} \tilde{f} d\mathbf{n}$  and the directional part represented by the first angular moment  $\mathbf{f}_1 = \frac{3}{4\pi} \int_{4\pi} \mathbf{n} \tilde{f} d\mathbf{n}$ , where  $\mathbf{n}$  is the transport direction. Then,

the first two angular moments [28] applied to the stationary form of (1) with collision operator (5) (extended by (23)) lead to the model equations

$$v \frac{\nu_e}{2} \frac{\partial}{\partial v} (f_0 - f_M) = \frac{v}{3} \nabla \cdot \mathbf{f}_1 + \frac{q_e}{m_e} \frac{\mathbf{E}}{3} \cdot \left( \frac{\partial \mathbf{f}_1}{\partial v} + \frac{2}{v} \mathbf{f}_1 \right), \quad (26)$$

$$v \frac{\nu_e}{2} \frac{\partial \mathbf{f}_1}{\partial v} - \nu_{scat} \mathbf{f}_1 = v \nabla f_0 + \frac{q_e}{m_e} \mathbf{E} \frac{\partial f_0}{\partial v} + \frac{q_e \mathbf{B}}{m_e c} \times \mathbf{f}_1, \quad (27)$$

where  $\nu_{scat} = \nu_{ei} + \frac{\nu_e}{2}$ . The system of equations (26) and (27) is called the **AP1 model** (AWBS + P1).

The AP1 model gives us information about the electron distribution function providing a bridge between kinetic and fluid description of plasma. For example the *flux* quantities as electric current and heat flux due to the motion of electrons

$$\mathbf{j} = \frac{4\pi}{3} q_e \int v \mathbf{f}_1 v^2 dv, \quad \mathbf{q}_h = \frac{4\pi}{3} \frac{m_e}{2} \int v^3 \mathbf{f}_1 v^2 dv,$$

where  $d\tilde{v} = v^2 dv$  the spherical coordinates metric, are based on corresponding velocity moments (integrals) of the first angular moment of EDF. Consequently, the explicit formula for the first angular moment from (27) proves to be extremely useful

$$\mathbf{f}_1 = \frac{\nu_{scat}^2 \mathbf{F}^* + \boldsymbol{\omega}_B \cdot \mathbf{F}^* - \nu_{scat} \boldsymbol{\omega}_B \times \mathbf{F}^*}{\nu_{scat}(\boldsymbol{\omega}_B^2 + \nu_{scat}^2)}, \quad (28)$$

because it provides a valuable information about the dependence of macroscopic *flux* quantities on electric and magnetic fields in plasma, where  $\boldsymbol{\omega}_B = \frac{q_e \mathbf{B}}{m_e c}$  is the electron gyro-frequency and  $\mathbf{F}^* = v \frac{\nu_e}{2} \frac{\partial \mathbf{f}_1}{\partial v} - v \nabla f_0 - \frac{q_e}{m_e} \mathbf{E} \frac{\partial f_0}{\partial v}$ .

#### A. Nonlocal Ohm's Law

The expression (28) becomes extremely useful when used to describe the electron fluid momentum, i.e. the current velocity moment can be written as

$$\mathbf{j}_{(f,E,B)} = \mathbf{J}_{Ohm} \frac{\partial f_0}{\partial v} \mathbf{E} + \frac{m_e}{q_e} \mathbf{J}_{Ohm} \left( v \nabla f_0 - v \frac{\nu_e}{2} \frac{\partial \mathbf{f}_1}{\partial v} \right), \quad (29)$$

where we used the following notation  $\mathbf{J}_{Ohm} \mathbf{g} = -\frac{4\pi q_e^2}{3m_e} \int v \frac{\nu_{scat}^2 \mathbf{g} + \boldsymbol{\omega}_B \cdot \mathbf{g} - \nu_{scat} \boldsymbol{\omega}_B \times \mathbf{g}}{\nu_{scat}(\boldsymbol{\omega}_B^2 + \nu_{scat}^2)} v^2 dv$  showing how the operator  $\mathbf{J}_{Ohm}$  acts on a general vector field  $\mathbf{g}$ . We refer to (29) as to the **nonlocal Ohm's law**. The need for a nonlocal Ohm's law to accurately capture magnetic field advection by the Nernst effect has previously been demonstrated [14, 31, 32], although a full investigation of how well our new Ohm's law captures this is beyond the scope of this article. The proper high  $Z$  ( $\nu_e \ll \nu_{ei}$ ) local asymptotic to the standard Ohm's law can be found when  $f_0 \rightarrow f_M$  and weak magnetization ( $\boldsymbol{\omega}_B \ll \nu_{ei}$ ) is

considered. Then (29) simplifies to

$$\mathbf{j} = -\frac{q_e^2}{m_e} \int \frac{v^3}{\nu_{ei}} \left( \mathbf{E} \frac{\partial f_M}{\partial v} + \frac{m_e}{q_e} v \nabla f_M \right) dv = \frac{16\sqrt{\frac{2}{\pi}} q_e^2 k_B^{\frac{3}{2}} T_e^{\frac{3}{2}}}{m_e^{\frac{5}{2}} \Gamma Z} \left[ \mathbf{E} - \frac{\frac{5}{2} n_e k_B \nabla T_e + \nabla n_e k_B T_e}{q_e n_e} \right], \quad (30)$$

which can be directly compared to the local fluid theory

$$\mathbf{E} = \sigma(f_0)^{-1} \mathbf{j} - \frac{\nabla P(f_0)}{q_e n_e} \xrightarrow{f_0 \rightarrow f_M} \mathbf{E}_l = \frac{\mathbf{j}}{\sigma_l} + \frac{\nabla p_e - \mathbf{R}_{T_e}}{q_e n_e}, \quad (31)$$

while addressing properly the local electric field  $\mathbf{E}_l$  given by the pressure  $p_e = n_e k_B T_e$ , the thermal force  $\mathbf{R}_{T_e} = -\frac{3}{2} n_e k_B \nabla T_e$  and the local electrical conductivity  $\sigma_l = 16\sqrt{\frac{2}{\pi}} q_e^2 k_B^{\frac{3}{2}} T_e^{\frac{3}{2}} / m_e^{\frac{5}{2}} \Gamma Z$  [12]. In (31) we defined the non-local electrical tensor conductivity

$$\sigma = \mathbf{J}_{Ohm} \frac{\partial f_0}{\partial v}, \quad (32)$$

and the nonlocal microscopic force

$$\nabla P = \sigma^{-1} m_e n_e \mathbf{J}_{Ohm} v \nabla f_0, \quad (33)$$

based on (29).

The local dependence of the AP1 current (30) on electric field and gradients of  $n_e$  and  $T_e$  clearly demonstrates, that (31) is a local version of (29). This also implies that (29) provides a very important physics related to the magnetic field source in terms of nonlocal Biermann battery, since the curl on the electric field (31) gives

$$\nabla \times \frac{\nabla P}{q_e n_e} \xrightarrow{f_0 \rightarrow f_M} \nabla \times \frac{\nabla p_e - \mathbf{R}_{T_e}}{q_e n_e} = \frac{k_B}{q_e n_e} \nabla T_e \times \nabla n_e. \quad (34)$$

The nonlocal Biermann battery equivalent (34) can lead to a spontaneous magnetic field generation under uniform density plasma profile as has been shown in [33].

A local version of the **nonlocal Ohm's law** (29) compared to the *generalized Ohm's law* (31) when a magnetic field is applied would require a much more delicate analysis and we leave it as a future complementary work.

## B. AWBS Nonlocal Magneto-Hydrodynamics

The *AWBS nonlocal magneto-hydrodynamic model* (Nonlocal-MHD) refers to two temperature single-fluid hydrodynamic model extended by a kinetic model of electrons using the AWBS transport equation, which provides a direct coupling between hydrodynamics and Maxwell equations.

Mass, momentum density, and total energy  $\rho$ ,  $\rho \mathbf{u}$ , and  $E = \frac{1}{2} \rho \mathbf{u} \cdot \mathbf{u} + \rho \varepsilon_i + \rho \varepsilon_e$ , where  $\rho$  is the density of plasma,  $\mathbf{u}$  the plasma fluid velocity,  $\varepsilon_i$  the specific internal ion energy density, and  $\varepsilon_e$  the specific internal electron energy

density, are modeled by the Euler equations in the Lagrangian frame [34, 35]

$$\frac{d\rho}{dt} = -\rho \nabla \cdot \mathbf{u}, \quad (35)$$

$$\rho \frac{d\mathbf{u}}{dt} = -\nabla(p_i + p_e) + \mathbf{j}_{(f, \mathbf{E}, \mathbf{B})} \times \mathbf{B}, \quad (36)$$

$$\rho C_{V_i} \frac{dT_i}{dt} = (\rho^2 C_{T_i} - p_i) \nabla \cdot \mathbf{u} - G(T_i - T_e), \quad (37)$$

$$\rho C_{V_e} \frac{dT_e}{dt} = (\rho^2 C_{T_e} - p_e) \nabla \cdot \mathbf{u} + G(T_i - T_e) - \nabla \cdot \mathbf{q}_{h(f, \mathbf{E}, \mathbf{B})} + Q_{IB}, \quad (38)$$

where  $T_i$  is the temperature of ions,  $T_e$  the temperature of electrons,  $p_i$  the ion pressure,  $p_e$  the electron pressure,  $\mathbf{q}_h$  the heat flux,  $Q_{IB}$  the inverse-bremsstrahlung laser absorption (which can also distort the distribution function away from a Maxwellian [36], strongly modifying the transport [37], an effect which will not be considered further here) and  $G = \rho C_{V_e} \nu_{ei}$  is the ion-electron energy exchange rate. The thermodynamic closure terms  $p_e$ ,  $p_i$ ,  $C_{V_i} = \frac{\partial \varepsilon_i}{\partial T_i}$ ,  $C_{T_i} = \frac{\partial \varepsilon_i}{\partial \rho}$ ,  $C_{V_e} = \frac{\partial \varepsilon_e}{\partial T_e}$ ,  $C_{T_e} = \frac{\partial \varepsilon_e}{\partial \rho}$  are obtained from an equation of state (EOS), e.g. the SESAME equation of state tables [38, 39].

The magnetic and electric fields are modeled by Maxwell equations

$$\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}, \quad (39)$$

$$\frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} = \nabla \times \mathbf{B} - \frac{4\pi}{c} \mathbf{j}_{(f, \mathbf{E}, \mathbf{B})}, \quad (40)$$

where the initial state of  $\mathbf{B}$  and  $\mathbf{E}$  obeys the Gauss's law.

We have explicitly written the current and heat flux as dependent on electron kinetics, represented by the electron distribution function  $f$ , and electric and magnetic fields. In principal,  $\mathbf{j}_{(f, \mathbf{E}, \mathbf{B})}$  and  $\mathbf{q}_{h(f, \mathbf{E}, \mathbf{B})}$  can be referred to as the *kinetic closure* and is provided by the **AP1 model** (26) and (27).

## C. Numerical Implementation of the AWBS Electron Kinetics

It is well known, that as the electron transport exhibits a quasi-steady behavior, the same holds for the electric field in (40) on the time scale of the fluid. Consequently, Ampere's law (40) takes a quasi-steady form  $\frac{4\pi}{c} \mathbf{j}_{(f, \mathbf{E}, \mathbf{B})} = \nabla \times \mathbf{B}$  when used in hydrodynamics. Proceeding further, one can make use of the **nonlocal Ohm's law** (29) to write a *fully kinetic form of Ampere's law* governing the electric field  $\mathbf{E}$

$$\mathbf{J}_{Ohm} \frac{\partial f_0}{\partial v} \mathbf{E} + \frac{m_e}{q_e} \mathbf{J}_{Ohm} v \nabla f_0 = \frac{c}{4\pi} \nabla \times \mathbf{B}. \quad (41)$$

In order to solve the kinetics of electrons, we adopt a high-order finite element discretization [40, 41] of

the model equations (26), (27), (41)

$$\mathbf{M}_{(\frac{v\nu_e}{2})}^{L_2} \cdot \frac{d\mathbf{f}_0}{dv} - \mathbf{V}_{(\frac{q_e\mathbf{E}}{3m_e})}^{L_2} \cdot \frac{d\mathbf{f}_1}{dv} = \mathbf{D}_{(\frac{\mathbf{v}}{3})}^{L_2} \cdot \mathbf{f}_1 + \mathbf{M}_{(\frac{2q_e\mathbf{E}}{3m_e v})}^{L_2} \cdot \mathbf{f}_1 + \mathbf{b}_{(\frac{v\nu_e}{2} \frac{\partial f_M}{\partial v})}^{L_2}, \quad (42)$$

$$\mathbf{M}_{(\frac{v\nu_e}{2})}^{H_1} \cdot \frac{d\mathbf{f}_1}{dv} - \mathbf{V}_{(\frac{q_e\mathbf{E}}{m_e})}^{H_1} \cdot \frac{d\mathbf{f}_0}{dv} = \mathbf{G}_{(v)}^{H_1} \cdot \mathbf{f}_0 + \mathbf{M}_{(\nu_{scat})}^{H_1} \cdot \mathbf{f}_1 + \mathbf{C}_{(\frac{q_e\mathbf{B}}{m_e c} \times)}^{H_1} \cdot \mathbf{f}_1, \quad (43)$$

$$\mathbf{J}_{(\frac{\partial f_0}{\partial v})}^{ND} \cdot \mathbf{E} = \mathbf{J}\mathbf{G}_{(\frac{m_e v}{q_e})}^{ND} \cdot \mathbf{f}_0 + \mathbf{b}_{(\frac{c}{4\pi} \nabla \times \mathbf{B})}^{ND}, \quad (44)$$

where the continuous differential operators are represented by standard discrete analogs (matrices of bilinear forms)  $\mathbf{M}, \mathbf{G}, \mathbf{D}, \mathbf{V}, \mathbf{C}$ , i.e. mass, gradient, divergence, vector field dot product, and vector field curl, and by  $\mathbf{J}, \mathbf{J}\mathbf{G}$  matrices specific to **nonlocal Ohm's law** (29). The linear form  $\mathbf{b}$  represents sources, i.e. temperature  $T_e$  via  $\frac{\partial f_M}{\partial v}$  and the curl of the magnetic field  $\mathbf{B}$ . These finite element discrete analogs are defined on piece-wise continuous  $L_2$  finite element space (domain of  $\mathbf{f}_0$ ), continuous  $H_1$  finite element space (domain of  $\mathbf{f}_1$ ) [40], and Nedelec finite element space (domain of  $\mathbf{E}$ ). We do not show their definitions since it is out of the scope of this article.

The strategy of solving (42) and (43) resides in integrating  $\frac{d\mathbf{f}_0}{dv}$  and  $\frac{d\mathbf{f}_1}{dv}$  along the velocity magnitude. This is done by starting the integration from the maximum velocity ( $v = 7v_{th}^{max}$  is a sufficiently high limit) to zero velocity using the Implicit Runge-Kutta method. The value  $v_{th}^{max}$  equals the electron thermal velocity corresponding to the maximum electron temperature in the current profile of plasma. It should be noted, that the backward integration concept is crucial for the model, since it corresponds to the deceleration of electrons due to collisions [42]. Consequently, we refer to *decelerating* AP1 model, which however, leads to the limitation of the electric field described in Appendix B.

## V. BENCHMARKING THE AWBS NONLOCAL TRANSPORT MODEL

Having shown several encouraging properties of the AWBS transport equation defined by (5) under local diffusive conditions in Section III, this section focuses on analyzing its behavior under nonlocal plasma conditions, extensively investigated in numerous publications [2, 10, 26, 43–46]. A variety of tests suitable for benchmarking the nonlocal electron transport models have been published [2, 25, 29, 47–49], we focus on conditions relevant to inertial confinement fusion plasmas generated by lasers.

We show results of our implementation of the AP1 nonlocal transport model presented in Section IV benchmarked against simulation results provided by a rather

complete set of kinetic models with varying complexity. The most reliable models represents a collisional Particle-In-Cell code Calder [22, 50] resolving the plasma frequency time scale, and a standard VFP codes Aladin and Impact [21]. In addition, we compare the SNB non-local transport model [51] used in hydrodynamic codes. That is a first time when a collisional PIC code is used for benchmarking of nonlocal electron transport models.

### Calder PIC code

The particle evolution in the phase-space, including small angle binary collisions, is described with the Maxwell equations (39), (40) coupled with the ion and electron Vlasov equations with the Landau-Beliaev-Budker collisions integral (LBB) [23, 52]

$$\frac{\partial f_\alpha}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f_\alpha + q_\alpha (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{p}} f_\alpha = C_{LBB}(f_\alpha, f_\alpha) + \sum_{\beta} C_{LBB}(f_\alpha, f_\beta). \quad (45)$$

The LBB collision integral takes the form

$$C_{LBB}(f_\alpha, f_\beta) = -\frac{\partial}{\partial \mathbf{p}} \cdot \frac{\Gamma_{\alpha\beta}}{2} \left[ \int \mathbf{U}(\mathbf{p}, \mathbf{p}') \cdot (f_\alpha \nabla_{\mathbf{p}'} f_\beta' - f_\beta' \nabla_{\mathbf{p}} f_\alpha) \right] d^3 \mathbf{p}', \quad (46)$$

where its relativistic kernel reads  $\mathbf{U}(\mathbf{p}, \mathbf{p}') = \frac{r^2/\gamma\gamma'}{(r^2-1)^{3/2}} [(r^2-1)\mathbf{I} - \mathbf{p} \otimes \mathbf{p} - \mathbf{p}' \otimes \mathbf{p}' + r(\mathbf{p} \otimes \mathbf{p}' + \mathbf{p}' \otimes \mathbf{p})]$  with  $\gamma = \sqrt{1+\mathbf{p}^2}$ ,  $\gamma' = \sqrt{1+\mathbf{p}'^2}$  and  $r = \gamma\gamma' - \mathbf{p} \cdot \mathbf{p}'$ . The momentum  $\mathbf{p}_\alpha$  ( $\mathbf{p}_\beta$ ) is normalized to  $m_\alpha c$  (resp.  $m_\beta c$ ). The collision operator (46) tends to (2) in the non-relativistic limit. The aforementioned model is solved in 3D by the PIC code CALDER. [22, 50].

### Impact and Aladin VFP codes

PIC simulations are extremely expensive as the collisions require description of the velocity space in 3 dimensions. Yet, a reduction of dimensions can be done by developing the distribution function in a Cartesian tensor series, equivalent to expansion in the spherical harmonics [53]. The first order form corresponds to the P1 approximation (25) and coupled with the Landau-Fokker-Planck collisional operator (2) leads to the P1-VFP model [21, 53]:

$$\frac{\partial f_0}{\partial t} + \frac{v}{3} \nabla \cdot \mathbf{f}_1 + \frac{q_e}{3m_e v^2} \frac{\partial}{\partial v} (v^2 \mathbf{E} \cdot \mathbf{f}_1) = C_{ee}^0(f_0), \quad (47)$$

$$\frac{\partial \mathbf{f}_1}{\partial t} + v \nabla f_0 + \frac{q_e \mathbf{E}}{m_e} \frac{\partial f_0}{\partial v} + \frac{q_e \mathbf{B}}{m_e} \times \mathbf{f}_1 = -\nu_{ei} \mathbf{f}_1. \quad (48)$$

where only the isotropic part of the distribution function in in the e-e collision integral (2) is used as the following

$$C_{ee}^0(f_0) = \frac{\Gamma}{v^2} \frac{\partial}{\partial v} \left[ C(f_0) f_0 + D(f_0) \frac{\partial f_0}{\partial v} \right], \quad (49)$$

$$C(f_0(v)) = 4\pi \int_0^v f_0(u) u^2 du,$$

$$D(f_0(v)) = \frac{4\pi}{v} \int_0^v u^2 \int_u^\infty w f_0(w) dw du.$$

The codes Impact and Aladin solve the system (47) and (48) with the Maxwell equations (39) and (40) in two spatial dimensions, assuming immobile ions.

The model AP1 uses similar equations as Aladin and Impact with the difference, that AP1 describes the steady-state electron distribution function with respect to the ions, and is using a simplified (linear) collision operator inherently coupled to ions via the hydrodynamic equations.

### SNB approach

Now considered as a standard nonlocal electron transport models in hydrodynamic codes, SNB [51] represents an efficient P1 method based on the velocity dependent form of the collision BGK operator. It uses EDF approximation representing deviation from the local BGK theory

$$\tilde{f} = f_M + \delta f_0 + \mathbf{n} \cdot (\mathbf{f}_{1M} + \delta \mathbf{f}_1). \quad (50)$$

Equations for the zero and first angular moments follow from the electron transport equation with scaled collision operator (21) according to the SNB approximation (50) (similar to (26) and (27))

$$r_B \delta f_0 = -\frac{v}{3} \nabla \cdot \delta \mathbf{f}_1 - \frac{v}{3} \nabla \cdot \mathbf{f}_{1M} - \frac{q_e \mathbf{E}}{m_e} \cdot \left( \frac{\partial \mathbf{f}_{1M}}{\partial v} + \frac{\partial \delta \mathbf{f}_1}{\partial v} + \frac{2}{v} (\mathbf{f}_{1M} + \delta \mathbf{f}_1) \right), \quad (51)$$

$$\frac{\nu_{ei}}{\xi} \delta \mathbf{f}_1 = -v \nabla \delta f_0 - \frac{q_e \mathbf{E}}{m_e} \cdot \frac{\partial \delta \mathbf{f}_1}{\partial v} - \underbrace{\frac{\nu_{ei}}{\xi} \mathbf{f}_{1M} - v \nabla f_M - \frac{q_e \mathbf{E}}{m_e} \frac{\partial f_M}{\partial v}}_{= 0 \text{ defines } \mathbf{f}_{1M}}, \quad (52)$$

where the magnetic field was neglected, the under-braced part of (52) when  $\delta f_0$  and  $\delta \mathbf{f}_1$  are zero defines the local anisotropic term

$$\mathbf{f}_{1M} = -\xi \lambda_{ei} f_M \left( \frac{\nabla n_e}{n_e} + \left( \frac{v^2}{2v_{th}^2} - \frac{3}{2} \right) \frac{\nabla T_e}{T_e} - \frac{q_e \mathbf{E}}{m_e v_{th}^2} \right), \quad (53)$$

and the efficiency of SNB resides in omitting the electric field effect (crossed out terms in (51) and (52)), which leads to a simple diffusion equation for the correction to the isotropic part of the distribution function

$$\frac{1}{\lambda_e^{SNB}} \delta f_0 - \nabla \cdot \frac{\lambda_{ei}^{SNB}}{3} \nabla \delta f_0 = \nabla \cdot \frac{\xi \lambda_{ei}}{3} f_M \frac{\nabla T_e}{T_e}, \quad (54)$$

where  $\frac{1}{\lambda_{ei}^{SNB}} = \frac{\nu_{ei}}{\xi v} + \frac{|q_e \mathbf{E}|}{\frac{1}{2} m_e v^2}$  and  $\lambda_e^{SNB} = \frac{v}{r_B \nu_e}$ , and the source term based on  $\mathbf{f}_{1M}$  simplifies by avoiding the electric field effect, density gradient and the  $v$ -dependent bracket in (53). The missing effect of  $\mathbf{E}$  in (54) is accounted for by an isotropic scattering in definition of  $\lambda_{ei}^{SNB}$  [51]. Consequently, the effect of the electric field in SNB is accounted for only via  $\mathbf{f}_{1M}$ , where the electric field is fixed to  $\mathbf{E}_L$ .

As shown previously, the BGK collision operator (21) provides one free parameter  $\zeta$ . We propose to use  $\zeta = 2$  giving  $r_B(Z \gg 1) = 2$  which agrees with  $r = 2$  in SNB formulation proposed in [49] for the case of ICF relevant plasma. We also have  $r_B(Z = 1) = 1.677$ , which means that our pure kinetic derivation of SNB varies just slightly from a constant value  $r_B = 2$  [49], yet it provides slightly better results. The explicit form of the anisotropic part of EDF then reads  $\mathbf{f}_1 = \mathbf{f}_{1M} - \lambda_{ei}^{SNB} \nabla \delta f_0$ .

### A. Heat-bath problem

AP1 is compared to Calder, Aladin, Impact, and SNB by calculating the heat flow in the case of a homogeneous plasma with a large temperature variation

$$T_e(z) = 0.575 - 0.425 \tanh((z - 450)s), \quad (55)$$

which exhibits a steep gradient at the point  $450 \mu\text{m}$  connecting a hot bath ( $T_e = 1 \text{ keV}$ ) and cold bath ( $T_e = 0.17 \text{ keV}$ ) and  $s$  is the parameter of steepness. This test is referred to as a simple non-linear heat-bath problem and originally was introduced in [47] and further investigated in [2, 25, 48, 49].

The total computational box size is  $700 \mu\text{m}$ . We performed Aladin, Impact, and Calder simulations showing an evolution of temperature starting from the initial profile (55). Due to the initial distribution function being approximated by a Maxwellian, the first phase of the simulation exhibits a transient behavior of the heat flux. After several ps the distribution adjusts properly to its asymptotic form and the heat flux profiles can be usefully compared. We then take the temperature profiles from Aladin/Impact/Calder and compare with AP1 and SNB models which calculate a stationary heat flow for a given temperature profile. For all heat-bath simulations the electron density, Coulomb logarithm and ionisation were kept constant and uniform. The Coulomb logarithm was held fixed throughout,  $\ln \Lambda = 7.09$ .

We show AP1 results for two ionization states, namely  $Z = 1, 10$  in FIG. 2 and FIG. 3, respectively, corresponding to a moderate nonlocality ( $\text{Kn}^e \sim 10^{-2}$ ) leading to a roughly 40 % inhibition compared to the local SH heat flux maximum. The original temperature profile steepness  $s = 1/50 \mu\text{m}$ . It is preferable to use  $\text{Kn}^e = \frac{\lambda_e}{\sqrt{Z+1} L_{Te}}$  instead of  $\text{Kn} = \frac{\lambda_{ei}}{L_{Te}}$ , because  $\sqrt{Z+1}$  provides a better scaling of nonlocality with respect to ionization [26], i.e. the flux inhibition and  $\text{Kn}^e$  are kept approximately the same when varying  $Z$  in FIG. 2 and FIG. 3. In addition to the heat flux profiles, we also show the distribution function details related to the approximate point of the heat flux maximum ( $460 \mu\text{m}$ ) and to the point of the nonlocal preheat effect ( $580 \mu\text{m}$ ) in the form of the flux moment of EDFs anisotropic part (24). The non-local preheat effect shows a very good agreement with previous results published in [48].






FIG. 2. AP1 performance in a low- $Z$  heat-bath problem compared to the VFP code Aladin (left) and the collisional PIC code Calder (right). The heat flux and temperature profiles at 20 ps are shown in the top plots also for AP1 and SNB. Middle and bottom plots show a kinetic detail of the anisotropic part of EDF (its flux velocity moment) at two different spatial points. The results of the local Lorentz gas theory scaled by the SH correction are also shown for reference. An excellent agreement in EDF between AP1 (AWBS collision operator (5)) and Calder (full Landau-Fokker-Planck collision operator (2)) is observed.

The top left plot of FIG. 2 shows heat flux profiles computed by Aladin, AP1, and SNB corresponding to the temperature  $T_e$  profile computed by Aladin and the top right plot of FIG. 2 shows heat flux profiles computed by Calder, AP1, and SNB corresponding to the temperature  $T_e$  profile computed by Calder. Both kinetic simulations by Aladin and Calder evolved up to 20 ps for  $Z = 1$ . The anisotropic part of EDF, in particular, the heat flux velocity moment  $q_1$ , at the heat flux maximum (triangle point) and at the nonlocal preheat region (circle point) computed by AP1 and SNB for the temperature profiles by Aladin and Calder, can be

used as a detailed comparison of four conceptually different models: the full anisotropy form (2) of the FP collision operator (Calder); the isotropic form (49) of the FP collision operator (Aladin); the simplified linear form (5) of the FP collision operator (AWBS in AP1); the nonlocal electron transport model (54) (SNB). Excellent match of  $q_1$  can be seen between AP1 and Calder at the both spatial points. On the other hand, the AP1 profiles of EDF provide a reasonable match to Aladin too, however, one observes a deviation which resembles to the low  $Z$  trend shown in FIG. 1, where AP1 corresponds to AWBS and Aladin to BGK curves. To summarize, various FP-

like codes are compared in detail, in particular collisional PIC for the first time, and all show a very good match. Furthermore, the effect of the anisotropy in the collision model, captured by AP1 and neglected by Aladin and Impact, proves to be important in the low- $Z$  plasma.

In the case  $Z = 10$ , we show heat flux profiles computed by Aladin, AP1, and SNB corresponding to the temperature  $T_e$  profile computed by Aladin up to 12 ps in the top plot of FIG. 3. Corresponding profiles of a self-consistently calculated electric fields by Aladin and AP1 (using the nonlocal Ohm's law) are shown in the higher middle plot. Also the local theory based electric field  $\mathbf{E}_L$  used by SNB is shown. EDF at the point of the approximate heat flux maximum (triangle) of the temperature profile is shown in the lower middle plot, where a very precise match between AP1 and Aladin can be observed, and  $q_1$  at the preheat point (circle) of the temperature profile is shown in the bottom plot. In the latter case AP1 shows a very similar properties as Aladin with a difference in magnitude corresponding to a higher heat flux computed by AP1 at this point.

SNB shows very good results of the heat flux profile in all three cases, i.e. compared to Aladin and Calder in FIG. 2 and to Aladin in FIG. 3. However, one can observe that the EDF kinetic solution of SNB provides only a qualitative image with respect to the reference green line solution. This is illustrated for example in FIG. 3, where the kinetics at preheat point plot reveals an insufficient electric field treatment (no return current). The kinetics at maximum point plot shows that the solution SNB solution approaches closely the local Lorentz\* solution and that significantly recedes from the reference fully kinetic solution (green line). These discrepancies can be attributed to the use of an inconsistent electric field in the case of SNB which uses  $\mathbf{E}_L$ . An electric field comparison is shown in FIG. 3, where it is shown that the local electric field treatment  $\mathbf{E}_L$  used in SNB fails in the preheat region and consequently leads to a significant violation of the plasma quasi-neutrality, i.e. a non-zero current, where one can observe an uncontrolled stream of electrons in the preheat and also an overestimation of negative return current around the heat flux maximum.

The AP1 model equations (26), (27), and (41) in general show a very good performance in all three cases when compared to the fully kinetic results (green line) by Aladin and Calder, which can be assigned to the AWBS collision operator and the consistent treatment of  $\mathbf{E}$  via nonlocal Ohm's law (29) in (41) (no  $\mathbf{B}$  field in 1D).

In addition, the Knudsen number  $\text{Kn}^e$  has been varied among the simulation runs in order to address a broad range of nonlocality of the electron transport corresponding to the laser-heated plasma conditions, i.e.  $\text{Kn}^e \in (0.0001, 1)$ . The variation of  $\text{Kn}^e$  arises from the variation of the uniform electron density  $n_e \in (10^{19}, 10^{23}) \text{ cm}^{-3}$  or the length scale given by the slope of the temperature profile  $s \in (1/2500, 1/25) \mu\text{m}$ . Results showing the heat flux maximum of an extensive set of simulations of vary-

$\text{Kn}^e$	$10^{-4}$	$10^{-3}$	$10^{-2}$	$10^{-1}$	1
$v_{lim}/v_{th}$	70.8	22.4	7.3	3.1	1.8

TABLE II. Scan over varying nonlocality ( $\text{Kn}^e$ ) showing the limit of the collision friction dominance over the deceleration of electrons due to the electric field force. The electric field effect is dominant for electrons with higher velocity than  $v_{lim}$  defined in (56).  $\text{Kn}^e$  and  $v_{th}$  are evaluated from the same plasma profiles.

ing  $\text{Kn}^e$  is shown in FIG. 4. When analyzing the simulation results shown in FIG. 4, we observed that the maximum of  $q_1$  at the maximum point tends to decrease with increasing  $\text{Kn}^e$  and that the interval of electron velocities important for the heat transport always belongs to  $3v_{th} < v < 4v_{th}$  for an example refer to the kinetics at maximum point in FIG. 2 and FIG. 3. We have also found an important observation related to the electron-electron collisions. The stopping force is dominated by the electric field for electrons with velocity above the velocity limit

$$v_{lim} = \sqrt{\frac{\sqrt{3}\Gamma m_e n_e}{2q_e |\mathbf{E}|}}, \quad (56)$$

as shown in Appendix B and this limit drops down significantly with increasing Knudsen number as can be seen in TABLE II. As a consequence, the electrons responsible for the heat flux ( $3v_{th} < v < 4v_{th}$ ) are preferably affected by the electric field rather than by collisions when  $\text{Kn}^e > 10^{-1}$ . According to TABLE II collisions dominate stopping for  $v < 3.1v_{th}$  when  $\text{Kn}^e = 10^{-1}$  and even a much lower value  $v < 1.8v_{th}$  when  $\text{Kn}^e = 1.0$ . This explains the unsatisfactory results of the *decelerating* AP1 model for high  $\text{Kn}^e$  shown in FIG. 4. Notably, the AP1 limited electric field effect (described in Appendix B) leads to a steep increase of error with respect to VFP code Aladin for  $\text{Kn}^e > 10^{-1}$ . For example  $v_{lim} \sim 4.3v_{th}$  for the maximum point EDF in FIG. 2.

## B. Hohlraum problem

Additionally to the steep temperature gradients, the laser-heated plasma experiments also involve steep density gradients and variation in ionization, which is a dominant effect in multi-material hohlraums at the interface between the helium gas-fill and the ablated high  $Z$  plasma.

In [49], a kinetic simulation of laser pulse interaction with a gas filled hohlraum was presented. Plasma profiles provided by a HYDRA simulation in 1D geometry of a laser-heated gadolinium hohlraum containing a helium

gas at time of 20 ns were used as input for the IMPACT [21] VFP code. For simplicity, the Coulomb logarithm was treated as a constant  $\ln\Lambda_{ei} = \ln\Lambda_{ee} = 2.1484$ . In reality, in the low-density corona  $\ln\Lambda$  reaches 8, which, however, does not affect the heat flux profile significantly. FIG. 5 shows the electron temperature  $T_e$  evolved during 10 ps by Impact and the electron density  $n_e$  profile. Along with plasma profiles the heat flux profiles of AP1, Impact, and SNB are also shown.

One can observe a very good match between AP1 and Impact computations in the preheat region. It is worth mentioning that in the surroundings of the heat flux maximum ( $\sim 1662\mu\text{m}$ ) the profiles of all plasma variables exhibit steep gradients with a change from  $T_e = 2.5\text{ keV}$ ,  $n_e = 5 \times 10^{20}\text{ cm}^3$ ,  $Z = 2$  to  $T_e = 0.3\text{ keV}$ ,  $n_e = 6 \times 10^{21}\text{ cm}^3$ ,  $Z = 44$  across approximately  $100\mu\text{m}$  (between  $1600\mu\text{m}$  and  $1700\mu\text{m}$ ), starting at the helium-gadolinium interface. In this region, we can see a qualitative match between AP1 and Impact providing a same sign of the heat flux divergence contributing to hydro, however, we observe that the electric field limitation explained in Appendix B leads to a stronger drop of the *decelerating* AP1 heat flux on the material interface, which then closely aligns to the Impact heat flux in the corona. On the other hand, SNB overestimates significantly the heat flux in the lower density part of the plasma up to the point of the heat flux maximum given by Impact (green line in FIG. 5), and more importantly, the opposite sign of the heat flux divergence compared to Impact (and AP1) in the steep gradients region close to the material interface. In the preheat region SNB performs very well. Nevertheless, it is important to stress that SNB required only 25 velocity groups compared to 250 velocity groups used by Impact and AP1 for this ICF relevant plasma conditions, thus making it a very efficient modeling approach though its description of kinetics is rather qualitative.

## VI. CONCLUSIONS

In conclusion, we have performed a thorough analysis of the AWBS transport equation for electrons originally introduced in [2] and extended it by adding the nonlocal version of Ohm's law. After properly redefining the e-e collision term, we have shown that the AWBS simplified linear form of the Fokker-Planck collision operator keeps some important kinetic properties analyzed in local diffusive regime, e.g. provides a correct dependence on  $Z$  (BGK requires an additional fix) and inherently includes the anisotropic part of the distribution function  $\mathbf{f}_1$ , which compares very well to the full Fokker-Planck operator. Under nonlocal transport plasma conditions, we benchmarked AP1 against the reference VFP codes Aladin and Impact, collisional PIC code Calder, and we also included the standard nonlocal approach SNB. It was a first time comparison of collisional PIC and VFP codes. AP1 performed very well over all simulation cases while capturing the important kinetic features compared

to the reference kinetic codes. Furthermore, our detailed analysis of the anisotropic part of the EDF provided by AP1 showed an excellent match with Calder and outperformed Aladin and Impact in the case of low- $Z$  plasma, which is attributed to the effect of anisotropy in the collision model. This suggests a promising AP1's capability in predicting general transport coefficients and the seeding of parametric laser plasma instabilities sensitive to the Landau damping of longitudinal plasma waves [2, 54], which is of great importance in ICF related plasmas [55]. Other kinetic effects as perpendicular transport, e.g. heat flow or magnetic field advection, occurring in magnetised plasma [56] are introduced in AP1 via the nonlocal Ohm's law, which correctly recovers the generalized Ohm's law in its local diffusive asymptotic. The importance of the nonlocal Ohm's law becomes obvious for  $\text{Kn}^e > 10^{-1}$ , where the stopping of nonlocal electrons is rather due to the electric field effect than the collisional friction. We have also shown a slightly new formulation of SNB based on the scaled BGK collision operator (21), which performed really well in the heat-bath problem and the corresponding heat flux profile, however, its EDF output is rather qualitative which also lead to non-precise results of the hohlraum problem. On the other hand we also observed an inaccurate kinetic results of the *decelerating* AP1 computation for highly nonlocal plasma conditions, which was explained by the velocity limit applied to the action of the electric field.

## ACKNOWLEDGMENTS

C. P. Ridgers would like to acknowledge funding from EPSRC (grant EP/M011372/1). This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 20142018 under grant agreement No 633053 (project reference CfP-AWP17-IFE-CCFE-01). The views and opinions expressed herein do not necessarily reflect those of the European Commission.

## Appendix A: Analysis of local diffusive regime

In order to analyze the local diffusive regime of BGK collision operator (8), we adopt the P2 form of EDF

$$\tilde{f}(z, v, \mu) = f^0(z, v) + \mu f^1(z, v) + \frac{3\mu^2 - 1}{2} f^2(z, v), \quad (\text{A1})$$

where  $f^2$  helps us to quantify the validity of the local transport regime. The left hand side of (7) acts on (A1)

as

$$\begin{aligned} & \mu \left( \frac{\partial \tilde{f}}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial \tilde{f}}{\partial v} \right) + \frac{q_e E_z}{m_e} \frac{1 - \mu^2}{v^2} \frac{\partial \tilde{f}}{\partial \mu} = \frac{q_e E_z}{m_e v^2} f^1 \\ & + \mu \left[ \frac{\partial}{\partial z} \left( f^0 - \frac{f^2}{2} \right) + \frac{q_e E_z}{m_e v} \frac{\partial}{\partial v} \left( f^0 - \frac{f^2}{2} \right) + \frac{3q_e E_z}{m_e v^2} f^2 \right] \\ & + \mu^2 \left[ \frac{\partial f^1}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial f^1}{\partial v} - \frac{q_e E_z}{m_e v^2} f^1 \right] + O(\mu^3). \quad (\text{A2}) \end{aligned}$$

The action on (A1) of the BGK operator (8) reads

$$\begin{aligned} \frac{1}{v} C_{BGK}(\tilde{f}) &= \frac{f_M - \tilde{f}}{\lambda_e} + \frac{1}{2} \left( \frac{Z}{\lambda_e} + \frac{1}{\lambda_e} \right) \frac{\partial}{\partial \mu} (1 - \mu^2) \frac{\partial \tilde{f}}{\partial \mu}, \\ &= \frac{f_M - f^0}{\lambda_e} + \frac{(3Z + 4)f^2}{2\lambda_e} \\ &\quad - \mu \frac{Z + 2}{\lambda_e} f^1 - \mu^2 \frac{9(Z + 1)}{2\lambda_e} f^2. \quad (\text{A3}) \end{aligned}$$

Consequently, if the isotropic and anisotropic parts of (A2) and (A3) are compared, one finds the following equations

$$f^0 = f_M + \frac{\lambda_e q_e E_z}{m_e v^2} f^1 + \frac{3Z + 4}{2} f^2, \quad (\text{A4})$$

$$\begin{aligned} f^1 &= -\frac{\lambda_e}{Z + 2} \left[ \frac{\partial f^0}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial f^0}{\partial v} + \frac{3q_e E_z}{m_e v^2} f^2 \right. \\ &\quad \left. - \frac{1}{2} \frac{\partial f^2}{\partial z} - \frac{q_e E_z}{2m_e v} \frac{\partial f^2}{\partial v} \right], \quad (\text{A5}) \end{aligned}$$

$$f^2 = \frac{2\lambda_e}{9(Z + 1)} \left[ \frac{\lambda_e q_e E_z}{m_e v^2} f^1 - \frac{\partial f^1}{\partial z} - \frac{q_e E_z}{m_e v} \frac{\partial f^1}{\partial v} \right]. \quad (\text{A6})$$

It is valid to assume that  $f^0 \approx f_M$ , i.e. that  $f_M \gg \frac{\lambda_e q_e E_z}{m_e v^2} f^1 + \frac{3Z + 4}{2} f^2$  in (A4) and  $f^0 \gg f^2$  in (A5), for local transport conditions  $Kn \ll 1$  and the *quasi-neutrality* constraint, corresponding to a zero current

$$j \equiv \frac{4\pi q_e}{3} \int v f^1 v^2 dv = 0, \quad (\text{A7})$$

leads to the electric field (same as the classical Lorentz electric field  $\mathbf{E}_L$  [30])

$$E_z = \frac{m_e v_{th}^2}{q_e} \left( \frac{1}{L_{n_e}} + \frac{5}{2} \frac{1}{L_{T_e}} \right), \quad (\text{A8})$$

and the anisotropic part of EDF takes the form (11).

It should be noticed that  $f^0$  equilibrates to  $f_M$  as  $O(Kn^2 v/v_{th})$ , since one can show that  $\frac{\lambda_e q_e E_z}{m_e v^2} f^1 \sim Kn^2$  and  $f^2 \sim Kn^2 v/v_{th}$ , when  $f^0 = f_M$  and  $E_z = \mathbf{E}_L$  in (A5) and (A6).

In the case of the AWBS operator (5) used in (7), its

action on (6) reads

$$\begin{aligned} \frac{1}{v} C_{AWBS}(\tilde{f}) &= \frac{vr_A}{\lambda_e} \frac{\partial}{\partial v} (\tilde{f} - f_M) \\ &\quad + \frac{1}{2} \left( \frac{Z}{\lambda_e} + \frac{r_A}{\lambda_e} \right) \frac{\partial}{\partial \mu} (1 - \mu^2) \frac{\partial \tilde{f}}{\partial \mu} \\ &= \frac{vr_A}{\lambda_e} \frac{\partial}{\partial v} (f^0 - f_M) \\ &\quad + \mu \left( \frac{vr_A}{\lambda_e} \frac{\partial f^1}{\partial v} - \frac{Z + r_A}{\lambda_e} f^1 \right), \quad (\text{A9}) \end{aligned}$$

where  $\nu_e^* = r_A \nu_e = \frac{vr_A}{\lambda_e}$  with  $r_A$  being a scaling parameter of the standard e-e collision frequency. Its purpose is to match AWBS heat flux to results obtained by Spitzer and Harm [7] obtained for any  $Z$ . Section III C shows that this match can be found with a constant  $r_A = 0.5$ .

One finds the following equations if the isotropic and anisotropic parts of (A2) and (A9) are compared

$$\frac{\partial}{\partial v} (f^0 - f_M) = \frac{\lambda_e q_e E_z}{r_A m_e v^2} \frac{f^1}{v}, \quad (\text{A10})$$

$$\frac{vr_A}{\lambda_e} \frac{\partial f^1}{\partial v} - \frac{Z + r_A}{\lambda_e} f^1 = \frac{\partial f^0}{\partial z} + \frac{q_e E_z}{m_e v} \frac{\partial f^0}{\partial v}. \quad (\text{A11})$$

If we assume that  $\frac{\partial f^0}{\partial v} = \frac{\partial f_M}{\partial v}$ , i.e.  $f^0 = f_M$ , the anisotropic part of the AWBS operator is governed by the equation (13).

## Appendix B: AP1 electric field limit

Interestingly, we have encountered a very specific property of the AP1 model with respect to the electric field magnitude. The easiest way how to demonstrate this is to write the model equations (26) and (27) in 1D (z-axis). Then, due to its linear nature, it is easy to eliminate one of the partial derivatives with respect to  $v$ , i.e.  $\frac{\partial f_0}{\partial v}$  or  $\frac{\partial f_{1z}}{\partial v}$ . In the case of elimination of  $\frac{\partial f_0}{\partial v}$  one obtains the following equation

$$\begin{aligned} \left( v \frac{\nu_e}{2} - \frac{2q_e^2 E_z^2}{3m_e^2 v \nu_e} \right) \frac{\partial f_{1z}}{\partial v} &= \frac{2q_e E_z}{3m_e \nu_e} \frac{\partial f_{1z}}{\partial z} + \frac{4\pi q_e E_z}{3m_e} \frac{\partial f_M}{\partial v} \\ &\quad + \frac{v}{3} \frac{\partial f_0}{\partial z} + \left( \frac{4q_e^2 E_z^2}{3m_e^2 v^2 \nu_e} + \left( \nu_{ei} + \frac{\nu_e}{2} \right) \right) f_{1z}. \quad (\text{B1}) \end{aligned}$$

It is convenient to write the bracket on the left hand side of (B1) as  $\frac{2}{3v\nu_e} \left( (\sqrt{3}v\frac{\nu_e}{2})^2 - \frac{q_e^2}{m_e^2} E_z^2 \right)$  from where it is clear that the bracket is negative if  $\sqrt{3}v\frac{\nu_e}{2} < \frac{q_e}{m_e} |\mathbf{E}|$ , i.e. there is a velocity limit for a given magnitude  $|\mathbf{E}|$ , when the collisions are no more fully dominant and the electric field introduces a comparable effect to the collision friction in the electron transport.

It can be shown, that the last term on the right hand side of (B1) is dominant and the solution behaves as

$$\Delta \mathbf{f}_1 \sim \exp \left( \frac{\frac{4q_e^2 E_z^2}{3m_e^2 v^2 \nu_e} + \left( \nu_{ei} + \frac{\nu_e}{2} \right)}{v \frac{\nu_e}{2} - \frac{2q_e^2 E_z^2}{3m_e^2 v \nu_e}} \Delta v \right), \quad (\text{B2})$$

where  $\Delta v < 0$  represents a velocity step of the implicit Euler numerical integration of decelerating electrons. However, (B2) exhibits an exponential growth for velocities above the friction limit (bracket on the left hand side of (B1))

$$v_{lim} = \sqrt{\frac{\sqrt{3}\Gamma m_e}{2q_e} \frac{n_e}{|\mathbf{E}|}}, \quad (\text{B3})$$

which makes the problem to be ill-posed.

In order to provide a stable model, we introduce a reduced electric field to be acting as the accelerating force of electrons

$$|\mathbf{E}_{red}| = \sqrt{3}v \frac{m_e}{q_e} \frac{\nu_e}{2}, \quad (\text{B4})$$

ensuring that the bracket on the left hand side of (B1) remains positive. We define a quantity  $\eta_{red} = \frac{|\mathbf{E}_{red}|}{|\mathbf{E}|}$ . Then, the AP1 model (26), (27) can be formulated as well posed

$$v \frac{\nu_e}{2} \frac{\partial}{\partial v} (f_0 - f_M) = \frac{v}{3} \nabla \cdot \mathbf{f}_1 + \frac{q_e}{m_e} \frac{\mathbf{E}}{3} \cdot \left( \eta_{red} \frac{\partial \mathbf{f}_1}{\partial v} + \frac{2(2 - \eta_{red})}{v} \mathbf{f}_1 \right), \quad (\text{B5})$$

$$v \frac{\nu_e}{2} \frac{\partial \mathbf{f}_1}{\partial v} - \nu_{scat} \mathbf{f}_1 = v \nabla f_0 + \frac{q_e \eta_{red}}{m_e} \mathbf{E} \frac{\partial f_0}{\partial v} + \frac{q_e \mathbf{B}}{m_e c} \times \mathbf{f}_1, \quad (\text{B6})$$

while introducing the reduction factor of the accelerating electric field and the compensation of the electric field effect via its angular term.

- 
- [1] J. R. Albritton, E. A. Williams, I. B. Bernstein, K. P. Swartz, Nonlocal Electron Heat Transport by Not Quite Maxwell-Boltzmann Distributions, *Phys. Rev. Lett.* 57 (1986) 1887–1890.
  - [2] D. D. Sorbo, J.-L. Feugeas, P. Nicolai, M. Olazabal-Loume, B. Dubroca, S. Guisset, M. Touati, V. Tikhonchuk, Reduced entropic model for studies of multidimensional nonlocal transport in high-energy-density plasmas, *Phys. Plasmas* 22 (2015) 082706.
  - [3] R. S. Cohen, L. Spitzer, Jr., P. M. Routly, The electrical conductivity of an ionized gas, *Phys. Rev.* 80 (1950) 230–238.
  - [4] J. H. Jeans, *Astronomy and Cosmogony*, Cambridge University Press, London, 1929.
  - [5] S. Chandrasekhar, Stochastic problems in physics and astronomy, *Rev. Mod. Phys.* 15 (1943) 1.
  - [6] M. Planck, Über einen Satz der statistischen Dynamik und seine Erweiterung in der Quantentheorie, *Sitzungsber. Preuss. Akad. Wiss.* 24 (1917) 324–341.
  - [7] L. Spitzer, Jr. and R. Härm, Transport phenomena in a completely ionized gas, *Phys. Rev.* 89 (1953) 977.
  - [8] J. H. Jeans, The equations of radiative transfer of energy, *Month. Not. Royal Astr. Soc.* 78 (1917) 28–36.
  - [9] M. N. Rosenbluth, W. M. MacDonald, D. L. Judd, Fokker-planck equation for an inverse-square force, *Phys. Rev.* 107 (1957) 1.
  - [10] A. R. Bell, R. G. Evans, D. J. Nicholas, *Phys. Rev. Lett.* 46 (1981) 243.
  - [11] J. P. Matte, J. Virmont, Electron heat transport down steep temperature gradients, *Phys. Rev. Lett.* 49 (1982) 1936–1939.
  - [12] S. I. Braginskii, Transport processes in a plasma, *Reviews of Plasma Physics* 1 (1965) 205.
  - [13] J. Hawreliak, D. M. Chambers, S. H. Glenzer, A. Gouveia, R. J. Kingham, R. S. Marjoribanks, P. A. Pinto, O. Renner, P. Soundhauss, S. Topping, E. Wolfrum, P. E. Young, J. S. Wark, Thomson scattering measurements of heat flow in a laser-produced plasma, *Journal of Physics B: Atomic, Molecular and Optical Physics* 37 (7) (2004) 1541.
  - [14] C. P. Ridgers, R. J. Kingham, A. G. R. Thomas, Magnetic cavitation and the reemergence of nonlocal transport in laser plasmas, *Phys. Rev. Lett.* 100 (2008) 075003.
  - [15] L. Willingale, A. G. R. Thomas, P. M. Nilson, M. C. Kaluza, S. Bandyopadhyay, A. E. Dangor, R. G. Evans, P. Fernandes, M. G. Haines, C. Kamperidis, R. J. Kingham, S. Minardi, M. Notley, C. P. Ridgers, W. Rozmus, M. Sherlock, M. Tatarakis, M. S. Wei, Z. Najmudin, K. Krushelnick, Fast advection of magnetic fields by hot electrons, *Phys. Rev. Lett.* 105 (2010) 095001.
  - [16] J. J. Bissell, C. P. Ridgers, R. J. Kingham, Field compressing magnetothermal instability in laser plasmas, *Phys. Rev. Lett.* 105 (2010) 175001.
  - [17] A. S. Joglekar, A. G. R. Thomas, W. Fox, A. Bhattacharjee, Magnetic reconnection in plasma under inertial confinement fusion conditions driven by heat flux effects in ohm's law, *Phys. Rev. Lett.* 112 (2014) 105004.
  - [18] A. S. Joglekar, C. P. Ridgers, R. J. Kingham, A. G. R. Thomas, Kinetic modeling of nernst effect in magnetized hohlraums, *Phys. Rev. E* 93 (2016) 043206.
  - [19] R. J. Henchen, M. Sherlock, W. Rozmus, J. Katz, D. Cao, J. P. Palastro, D. H. Froula, Observation of nonlocal heat flux using thomson scattering, *Phys. Rev. Lett.* 121 (2018) 125001. doi:10.1103/PhysRevLett.121.125001.
  - [20] A. Thomas, M. Tzoufras, A. Robinson, R. Kingham, C. Ridgers, M. Sherlock, A. Bell, A review of Vlasov-FokkerPlanck numerical modeling of inertial confinement fusion plasma, *Journal of Computational Physics* 231 (3) (2012) 1051 – 1079.
  - [21] R. J. Kingham, A. R. Bell, An implicit Vlasov-Fokker-Planck code to model non-local electron transport in 2-D with magnetic fields, *J. Comput. Phys.* 194 (194) (2004) 1–34.
  - [22] F. Perez, L. Gremillet, A. Decoster, M. Drouin, E. Lefebvre, Improved modeling of relativistic collisions and collisional ionization in particle-in-cell codes, *Phys. Plasmas* 19 (2012) 083104.
  - [23] L. Landau, Kinetic equation for the coulomb effect, *Phys. Z. Sowjetunion* 10 (1936) 154.
  - [24] N. J. Fisch, Theory of current drive in plasmas, *Rev.*

- Mod. Phys. 59 (1987) 175.
- [25] D. D. Sorbo, J.-L. Feugeas, P. Nicolai, M. Olazabal-Loume, B. Dubroca, V. Tikhonchuk, Extension of a reduced entropic model of electron transport to magnetized nonlocal regimes of high-energy-density plasmas, *Laser Part. Beams* 34 (2016) 412–425.
  - [26] J. F. Luciani, P. Mora, J. Virmont, Nonlocal heat transport due to steep temperature gradients, *Phys. Rev. Lett.* 51 (1983) 1664–1667.
  - [27] P. Bhatnagar, E. Gross, M. Krook, A Model for Collision Processes in Gases. I. Small Amplitude Processes in Charged and Neutral One-Component Systems, *Phys. Rev.* 94 (1954) 511–525.
  - [28] I. P. Shkarofsky, T. W. Johnston, M. P. Bachynskii, *The particle Kinetics of Plasmas*, Addison-Wesley, Reading, 1966.
  - [29] E. M. Epperlein, R. W. Short, A practical nonlocal model for electron heat transport in laser plasmas, *Phys. Fluids B* 3 (1991) 3092–3098.
  - [30] H. A. Lorentz, The motion of electrons in metallic bodies, in: *Proceedings of the Royal Netherlands Academy of Arts and Sciences*, Amsterdam, Vol. 7, 1905, pp. 438–453.
  - [31] J. F. Luciani, P. Mora, A. Bendib, Magnetic field and nonlocal transport in laser-created plasmas, *Phys. Rev. Lett.* 55 (1985) 2421–2424.
  - [32] J. P. Brodrick, M. Sherlock, W. A. Farmer, A. S. Joglekar, R. Barrois, J. Wengraf, J. J. Bissell, R. J. Kingham, D. D. Sorbo, M. P. Read, C. P. Ridgers, Incorporating kinetic effects on nernst advection in inertial fusion simulations, *Plasma Physics and Controlled Fusion* 60 (8) (2018) 084009.
  - [33] R. J. Kingham, A. R. Bell, Nonlocal magnetic-field generation in plasmas without density gradients, *Phys. Rev. Lett.* 88 (2002) 045004. doi:10.1103/PhysRevLett.88.045004.
  - [34] M. Holec, J. Limpouch, R. Liska, S. Weber, High-order discontinuous Galerkin nonlocal transport and energy equations scheme for radiation hydrodynamics, *Int. J. Numer. Meth. Fl.* 83 (2017) 779.
  - [35] M. Holec, J. Nikl, S. Weber, Nonlocal transport hydrodynamic model for laser heated plasmas, *Phys. Plasmas* 25 (2018) 032704.
  - [36] A. B. Langdon, Nonlinear inverse bremsstrahlung and heated-electron distributions, *Phys. Rev. Lett.* 44 (1980) 575–579.
  - [37] C. P. Ridgers, A. G. R. Thomas, R. J. Kingham, A. P. L. Robinson, Transport in the presence of inverse bremsstrahlung heating and magnetic fields, *Physics of Plasmas* 15 (9) (2008) 092311.
  - [38] T. Group, SESAME report on the Los Alamos equation-of-state library, Tech. Rep. Tech. Rep. LALP-83-4, Los Alamos National Laboratory, Los Alamos (1983).
  - [39] S. P. Lyon, J. D. Johnson, SESAME: The Los Alamos national laboratory equation of state database, Tech. Rep. LA-UR-92-3407, Los Alamos National Laboratory, Los Alamos (1992).
  - [40] V. Dobrev, T. Kolev, R. Rieben, High-order curvilinear finite element methods for Lagrangian hydrodynamics, *SIAM J. Sci. Comput.* 34 (2012) B606–B641.
  - [41] MFEM: Modular finite element methods, [mfem.org](http://mfem.org).
  - [42] M. Touati, J.-L. Feugeas, P. Nicolai, J. Santos, L. Gremillet, V. Tikhonchuk, A reduced model for relativistic electron beam transport in solids and dense plasmas, *New J. Phys.* 16 (2014) 073014.
  - [43] R. C. Malone, R. L. McCroy, R. L. Morse, *Phys. Rev. Lett.* 34 (1975) 721.
  - [44] D. G. Colombant, W. M. Manheimer, M. Busquet, Test of models for electron transport in laser produced plasmas, *Phys. Plasmas* 12 (2005) 072702.
  - [45] A. V. Brantov, V. Y. Bychenkov, V. T. Tikhonchuk, Nonlocal electron transport in laser heated plasmas, *Phys. Plasmas* 5 (1998) 2742–2753.
  - [46] G. Schurtz, P. Nicolai, M. Busquet, A nonlocal electron conduction model for multidimensional radiation hydrodynamics codes, *Phys. Plasmas* 4238 (2000) 7.
  - [47] A. Marocchino and M. Tzoufras and S. Atzeni and A. Schiavi and Ph. Nicolai and J. Mallet and V. Tikhonchuk and J.-L. Feugeas, Comparison for non-local hydrodynamic thermal conduction models, *Phys. Plasmas* 20 (2013) 022702.
  - [48] M. Sherlock, J. P. Brodrick, C. P. Ridgers, A comparison of non-local electron transport models for laser-plasmas relevant to inertial confinement fusion, *Phys. Plasmas* 24 (2017) 082706.
  - [49] J. P. Brodrick, R. J. Kingham, M. M. Marinak, M. V. Patel, A. V. Chankin, J. T. Omotani, M. V. Umansky, D. D. Sorbo, B. Dudson, J. T. Parker, G. D. Kerbel, M. Sherlock, C. P. Ridgers, Testing nonlocal models of electron thermal conduction for magnetic and inertial confinement fusion applications, *Phys. Plasmas* 24 (2017) 092309.
  - [50] E. Lefebvre, N. Cochet, S. Fritzler, V. Malka, M.-M. Alonard, J.-F. Chemin, S. Darbon, L. Disdier, J. Faure, A. Fedotoff, O. Landoas, G. Malka, V. Mot, P. Morel, M. R. L. Gloahec, A. Rouyer, C. Rubbelynck, V. Tikhonchuk, R. Wrobel, P. Audebert, C. Rousseaux, Electron and photon production from relativistic laser-plasma interactions, *Nuclear Fusion* 43 (7) (2003) 629.
  - [51] G. Schurtz, P. Nicolai, M. Busquet, A nonlocal electron conduction model for multidimensional radiation hydrodynamics codes, *Phys. Plasmas* 7 (2000) 4238–4249.
  - [52] S. T. Beliaev, G. I. Budker, *The Relativistic Kinetic Equation*, Soviet Physics Doklady 1 (1956) 218.
  - [53] T. W. Johnston, Cartesian Tensor Scalar Product and Spherical Harmonic Expansions in Boltzmann’s Equation, *Phys. Rev.* 120 (1960) 1103–1111.
  - [54] R. Goldston, P. Rutherford, *Introduction to Plasma Physics*, CRC Press, 1995.
  - [55] R. K. Kirkwood, J. Moody, J. Kline, E. Dewald, S. Glenzer, L. Divol, P. Michel, D. Hinkel, R. Berger, E. Williams, J. Milovich, L. Yin, H. Rose, B. MacGowan, O. Landen, M. Rosen, J. Lindl, A review of laserplasma interaction physics of indirect-drive fusion, *Plasma Phys. Contr. F.* 55 (2013) 103001.
  - [56] C. A. Walsh, J. P. Chittenden, K. McGlinchey, N. P. L. Niasse, B. D. Appelbe, Self-generated magnetic fields in the stagnation phase of indirect-drive implosions on the national ignition facility, *Phys. Rev. Lett.* 118 (2017) 155001. doi:10.1103/PhysRevLett.118.155001.




FIG. 3. A moderate- $Z$  heat-bath problem. The temperature profile evolved up to 12 ps by Aladin. Top plots show heat flux profiles and electric fields by AP1, Aladin, and SNB. The resulting current of SNB using explicitly the local electric field  $E_L$  is also shown. Bottom plots show a kinetic detail of the anisotropic part of EDF (its flux velocity moment) at two different spatial points by AP1 and SNB compared to Aladin.




FIG. 4. The heat-flux inhibition compared to the local SH theory along varying nonlocality ( $\text{Kn}^e$ ) in the heat bath problem. AP1 compares well to the full kinetic simulations by Aladin, Impact, and Calder, up to  $\text{Kn}^e \sim 10^{-1}$ . For higher nonlocality *decelerating* AP1 departs significantly from the reference solutions because of the electric field limiting in accordance with the velocity limit in TABLE II.




FIG. 5. Heat flux profiles by AP1, Impact and SNB along the electron temperature  $T_e$  and electron density  $n_e$  profiles in a laser-heated gadolinium hohlraum with a helium gas-fill.