

Contents

1	Introduction	3
1.1	Wheat Breeding	5
1.2	Wheat Genetics	5
1.3	Wheat Genomics	5
1.4	Sequencing	5
1.5	Sequence analysis	9
1.5.1	Ambiguity Codes	10
1.5.2	RNA-Seq	10
1.6	Wheat specific resources resources	11
2	PolyMarker: A fast polyploid primer design pipeline	12
2.1	Pipeline	12
2.2	Applications of PolyMarker	16
2.2.1	KASP assays for public sets of SNPs	17
2.2.2	SNPs in a mutant population	17
2.2.3	Deletions on a mutant population	17
2.2.4	PolyMarker public web service	18
2.2.5	Genotyping of <i>Puccinia striiformis</i> f. sp. <i>tritici</i> isolates.	19
2.3	Discussion	19
3	Genetic map of <i>Yr15</i> with RNA-Seq	20
3.1	Mapping population	24
3.2	Sequencing and mapping	26
3.3	SNP Calling	28
3.4	Bulk Frequency Ratios	31
3.5	<i>In silico</i> mapping	34
3.6	Assay selection	37
3.7	SNP Validation	42

<i>CONTENTS</i>	2
3.8 Genetic map	44
3.9 Bioinformatic methods	47
3.9.1 Alignment reads to gene models	47
3.9.2 Bulk Frequency Ratios	47
3.9.3 Alignment between gene models	48
3.10 Discussion	48
4 Gene expression (expVIP)	52
4.1 Expression experiments (Introduction)	52
4.2 Database design	52
4.3 Analysis pipeline	52
4.4 Graphical interface	52
4.5 Conclusions	52
5 Conclusions and final remarks	53
A Supplemental tables	54
A.1 PolyMarker supplemental tables.	55
B Quality control	65
B.1 Sequence read quality	66
B.2 Sequence GC content	68

Chapter 1

Introduction

It defines the objectives and the importance of the research. It focus on the the application of Next Generation Sequencing to molecular biology, wheat genetics and ultimately to breeding programs. It also mentions the current status of the wheat reference genome and other resources (genetic maps, markers) the need of tools to query them effectively.




Figure 1.1: Timeline of the projects carried on during this PhD and the wheat resources that were released in the same period of time.

1.1 Wheat Breeding

An overview of how breeding is carried on currently, the different sources of genetic diversity and the relevance of fixing agriculturally important traits.

1.2 Wheat Genetics

The section describes alleles and the concept of gene, both as a locus in the genome (Quantitative Trait Locus, QTL) and an specific transcript (central dogma of molecular biology). Finally, it discusses traditional Mendelian inheritance and the effect of polyploidy. Some of this is described in the Yr15 chapter, maybe it is not needed anymore here.

1.3 Wheat Genomics

A description of the current status of the wheat genome (Mayer et al. (2014), Chapman et al. (2015)), the different available assemblies and approaches to sort the scaffolds (Genome Zipper, the various genetic maps).

1.4 Sequencing

The Human Genome Project used Sanger sequencing Lander et al. (2001). This technology is the current gold standard in terms of quality of the sequence. It evolved from electrophoresis gels where the bands represented bases to a fully automated technique. However, the throughput is limited and doing genome wide analysis has prohibitive costs. In the second half of the 2000s high-throughput sequencing technologies emerged which had reduced the cost of sequencing. The main principle of the second-generation sequencing is to produce clusters of clones (i.e. ePCR), fix them in a plate and then add bases with a fluorescent marker. The reaction happens in parallel in millions of clusters at the same time. With each cycle, a picture is taken, showing the fluorescence of each base. Then, image processing algorithms find where in the image the clusters are and the bases are called. At this scale, the volume and complexity of the information is not trivial to manipulate, hence computing is required.

According to the objectives of the experiment and the quality and volume of the available DNA, the library can be prepared on fragments of different sizes, the classification of the available sequencing for the fragments is the following Myllykangas et al. (2012); Metzker (2010); Shendure and Ji (2008); Hutchison (2007):

Single end When the fragments are short, it is possible to just sequence from the 5'-end the read.

Read Pairs When the sample consists fragments of up to 500bp, it is possible to read the 5' end up to the read length where the quality starts to drop, the molecule can be turned upside down, reverse complemented and sequence backwards. It is not required, but ideally, the fragments sequenced with read pairs should be selected to have a homogeneous size. The reads are in opposite orientation relative to each other.

Overlapping Read Pairs are a variation to read pairs, where the size of the fragment is shorter than two times the read length. This allows an alignment between the two fragments to get a longer read with the limitations of the instrument.

Mate pairs are used to get reads separated at distances between 1kbp and 5kbp. To achieve this, the molecule is circularised and the point where the two ends of the fragment were joined a biotin marker is inserted. Then, the molecule is fragmented again and the fragments containing the biotin are sequenced in the same fashion as read pairs. The resulting reads have the same orientation.

There are several types of experiments that can be analysed with high throughput sequencing, accordingly, different protocols for the sample preparation exist. The following is a short list of some of them

Whole genome shotgun When a sample is prepared for WGS, the DNA is extracted and chopped in fragments and sequenced. The reads obtained are, in principle, randomly distributed across the whole genome

RNA-Seq . Instead of sequencing DNA, mRNA is captured and sequenced. The fragments are not amplified in any way, to enable a portrait of the gene expression levels.

ChIP-SEQ . Chromatin Immunoprecipitation is used to find relationships between proteins and DNA sequence. It is useful to find transcription factors and replication-related proteins.

Amplicon sequencing . Used primarily to do barcoding of species. A known gene is amplified (i.e. 16S) with the intention of characterising the species present in the sample.

Metagenomic capture From a mixed sample (soil, root, animal fluids) all the DNA is extracted and sequenced, this gives a snapshot of the microbial community in the sample

RAD-seq Restriction site associated DNA markers are useful to do population analysis. The technique focuses on sequencing regions around restriction sites and the variations around them can be used to genotype individuals.

Exon capture The DNA is extracted and baits are used to attract the regions with motifs common around exons. This allows to sequence only the genes and regions near them.

The different sequencing technologies available as of 2013 have different yields, advantages and disadvantages, as described below:

Illumina Each fragment is amplified using bridge amplification over and over in the same place in the plate to form clusters. After the clusters are formed, a last cycle of amplification is carried on with the bases being added to the template, with the intervention of a polymerase, have a fluorescent marker which makes the cluster glow depending on the added base. It adds one base per cycle. With a read length between 75bp and 250bp is currently the most widely adopted platform. As a de facto standard, many tools exist to cope bioinformatically with the biases of the machine. The run takes 4 or 9 days, depending on days, depending if one or two reads are generated for each fragment. It produces up to 35 gigabases per run.

SOLiD The preparation of the fragments is similar to Illumina, however, when adding the bases they are added in pairs. This technique is called sequencing by ligation as it uses a DNA Ligase, as opposed

to a polymerase, to determine the transition between bases. The resulting sequence is not in base space, but in colour space, which represents the transition state between bases. This technique is robust for finding SNPs when you have a good reference where to align the reads. However, the number of tools available and the research done to analyse sequences in colour space is low compared to the tools using base space. The runs take between one and two weeks to complete, with a yield of up to 50 gig abases per run. The read length can be up to 50 bases

Roche/454 The fragments are cloned in beads, which then fall in wells in the slide. The sequencing is done by adding nucleotides in a determined order. The next nucleotides to be added in the reaction contain a fluorescent marker. The bases are not added one by one, but all the bases that are the same are added together. The amount of glow on each well can tell how many times a base is added. As the glow is not a discrete number, when a long homopolymer appear (above 5 bases) the likelihood of having a wrong count of the homopolymer is increased. The average read length varies between 300 and 700bp. A run usually takes half a day, but it only yelds 0.45 gigabases. The cost of the reagents is relatively expensive, but if the experiment requires longer reads it is a good option.

PacBio Opposed to all the previous technologies, Pacific Biosciences has developed a sequencing technology where the molecules doesn't need to be PCR amplified before the sequencing. The glass slide used contains wells with a depth of 100nm where a polymerase lays at the bottom. The nucleotides to be added have a fluorescent marked that is freed when the polymerase adds the nucleotide, releasing a light signal, which then can be captured from the bottom of the glass. The error rate for this technology is still high (about 10% of the bases are miscalled), however reading several times the same molecule reduce the error rate. The main advantage is that the reads can be over 1kbp.

OpGen Additionally, high-througput optical mapping technologies, like OpGen, are becoming accessible. The maps are done by fixing single molecules of DNA are held on a slide. Then, restriction enzymes targeted to specific digestion sites cut the fragment and fluorescent

markers are added to the ends of the fragments. Finally, the fragments are visualised and the size of the molecules is measured by the distance between fluorescent points in the slide. This is done with several fragments at the same time. Then, the distances between restriction sizes can be compared across all the fragments to generate a consensus. Finally, if you have contigs from other technologies, it is possible to complement the information and get better assemblies. Even without the contigs, the data can be used to compare translocations within strains of different bacteria or homologous species at a chromosome level.

ION Torrent (Do some research on newer sequencing things)

1.5 Sequence analysis

This section discusses the criteria to decide analysis done after sequencing, when to do re-alignments or *de novo* assemblies, how to do SNP calling in diploid and polyploid organisms and the bulk frequency ratios.

DNA sequence alone is not enough to understand the biology behind, a context is required. There are databases like Ensembl and NCBI that act as repositories of the known public sequences.

From the computational point of view, the problem can be viewed as a string matching. The Smith-Waterman Smith and Waterman (1981) and Needleman-Wunsch Needleman and Wunsch (1970) algorithms are the gold standard interns of accuracy looking for similarity between sequences. However, the execution time for both of them is prohibitive to run in massive databases. The algorithm execution time is $O(mn)$, as it requires calculating a matrix of size mn where m is the target sequence and n is the query sequence. To scale this to a manageable problem algorithms like BLAST index the references and use heuristics to make the search more manageable, with some penalty in the accuracy. This alignments tools are useful for long stretches of DNA (like cDNA or contigs) Altschul et al. (1990).

TODO: List of global aligners -BLAST -BLAT -Exonerate -nucmer

When looking at a protein level, where the sequences may be only loosely similar, Hidden Markov Models (HMM) are used to search for

protein families. This can be useful to annotate putative proteins and their functions. HMMs require a training dataset, where proteins are previously annotated and the reference is a model encoding the characteristics of a family, with associated probabilities. Hence, this technique is something between a sequences aligner and a classifier Eddy (2004).

When analysing high-throughput sequencing, having millions of short sequences make unfeasible to try to align the data to every possible reference. However, one can take in advantage the fact that you know which organism you are looking for and, if available, use a genomic reference. For this, tools like MAQ, BWA, Bowtie, among others, provide indexed search. Once you have your reads aligned to a reference you can do more analysis, depending on the biological question being asked and the type of sequencing carried on. Fortunately, most of the Short-Read sequence alignment produce similar outputs and the SAM format is becoming a de facto standard. This is allowing to make more modularised downstream analysis where you can test different aligners with different settings and pick the algorithm that better fits your experiment Liu and Schmidt (2012); Li and Durbin (2009); Li et al. (2009).

1.5.1 Ambiguity Codes

Make a table with the ambiguity codes and why they are useful.

1.5.2 RNA-Seq

One way to narrow down which genes are involved in certain trait or response to the environment is to focus on studying only the expressed genes. One of the techniques involving high-throughput sequencing is RNA-Seq. This technique captures the messenger RNA in the tissue being studied and sequenced. The premise is that you will find a gene more expressed if it is being used by the organism. Some proteins with a vital role for the cell are always expressed (i.e. RuBisCO for carbon fixation in plants GM (2000)). On the simplest of the experiments you would need two datasets to compare, one with the gene being looked expressed and one where it is not. The expression can come from different environmental conditions, development stage or different genotypes. Mortazavi et al. (2008)

Depending on how much *a priori* information of the analysed organism is available different bioinformatic approaches can be used.

Transcriptome alignment The reads are aligned to a database of known cDNA. Ideally, alternative splicing sequences are available, so a simple alignment should work (i.e. BWA, bowtie).

Genomic alignment The reads are aligned to the genome. The splice junctions, introns and axons need to be accounted, so simple alignment doesn't work. Regular alignments are used, but the reads may be trimmed at fixed sizes to allow discontinuous alignments using regular tools (i.e. Stampy, tophat/cufflinkns)

De Novo transcriptome assembly If a reference of the organism is not available, it is possible to generate a draft transcriptome with the RNA-Seq reads with traditional assemblers (velvet, abyss) or with specialised assembler tools like Trinity.

Once you have the alignments it is possible to evaluate the relative expression of the genes in the sample calculating the Reads per Kilobase per Million mapped reads (RPKM) or the Transcripts per Million (TPM). This normalises the expression by the amount of sequenced data and can be used to find which genes change in expression volume across different samples.

1.6 Wheat specific resources resources

Gene models -UniGene -UCW Gene models -Gene annotation IWGSC
-Gene annotation TGACv1

Genetic maps -Wang -Chapman/PopSeq (is the same populaiton, improved)

Markers -90k -820k -MASwheat/SRR

Portal -CeralsDB -MASWheat -Ensembl -Wheat-expression

Assemblies -Chapman -IWGSC -TGACv1 -NRGene (unpublished?)
-454 Liverpool

A compilation of the currently available resource for whet genetics and genomics. MAS wheat, CeralsDB, Ensembl, etc.

Chapter 2

PolyMarker: A fast polyploid primer design pipeline

One of the main challenges of working with polyploid species is the design of genome specific molecular markers. This is particularly true when targeting conserved homoeologue regions, where a primer could bind to a pair, or triplet, of identical sequences. For that reason, designing primers for polyploids require to include bases that are specific to the target, in addition to the physicochemical properties of the primer. The traditional methodology to find primer candidates include a blast search and a local alignment, select the primer candidates manually, and finally, validate the primers with a tool, like **Primer3** (Rozen and Skaletsky, 2000). To reduce the time invested in designed primers I have developed PolyMarker (Ramirez-Gonzalez et al., 2015a), a pipeline to automate the primer design for polyploid organisms.

2.1 Pipeline

PolyMarker is an automated pipeline that takes as input a list of SNPs and a reference file and produces a list of primer triplets for SNP genotyping. The list of SNPs is first converted to a FASTA file with ambiguity codes (Cornish-Bowden, 1985). The template sequences are aligned with **exonerate** (Slater and Birney, 2005) to find the homoeologous regions to the target sequence. Then, the alignment between homoeologues is refined using **MAFFT** (Katoh and Standley, 2013). A list of candidate variations is produced and used as input for **Primer3** (Rozen and Skaletsky, 2000). Finally, the output of **Primer3** is parsed to find the best




Figure 2.1: Steps and tools called by PolyMarker. The colour of the boxes represent: the step is an action inside the script (red); actions of the script (orange); temporary files (yellow); inputs (blue) and; output (green)

primer pair that contains a the targeted SNP and a base that is specific to the target genome (Figure 2.1). The pipeline is written as a Ruby script, using parsers and wrappers from BioRuby (Goto et al., 2010) and bio-samtools (Etherington et al., 2015; Ramirez-Gonzalez et al., 2012). The software is open source and released as a biogem (Bonnal et al., 2012), `bio-polyploid-tools`, the source code is available in github: <https://github.com/TGAC/bioruby-polyploid-tools>.

The PolyMarker input consist on SNP list with: unique name for the marker, the target chromosome and the sequence for the marker. The alternative alleles are surrounded by square brackets within the sequence. PolyMarker can take a list of several markers and design them in batch (Figure 2.2a). A FASTA file is produced with all the template sequences, with the alternative alleles substituted by the IUAPC ambiguity codes (Cornish-Bowden, 1985). The flanking sequence surrounding the SNP is limited by default to 100bp to reduce the search time and avoid missing regions that diverge near the SNP, as when the variation is near an intron-exon junction.

The template sequences are aligned to the reference using `exonerate` (Slater and Birney 2005; Figure 2.2b). The alignment is refined with the `--model est2genome` option, to allow the search of sequences coming from transcripts, a common source of SNPs (Allen et al., 2011). The `exonerate` output is formatted with the `--ryo` (roll your own format) to get an output easy to parse. All the hits that contain the SNP are extracted from the reference with a flanking sequence that extend out of the hit, by default, to 100bp on each side of the SNP, Figure 2.2c. The size of the flanking sequence can be set to different sizes to allow the design of different types of primers. Different homoeologues may contain small indels, Figure 2.2d. To enable a comparasion base-per-base, a local alignment with `MAFFT` (Kato and Standley, 2013) is produced, Figure 2.2e.

PolyMarker searches across each base in the local alignment to identify the variations across homoeologues and the target marker. A mask is produced to highlight the bases with a variations, Figure 2.2f, on the following categories:

Specific	Homoeologous polymorphism which is only present in the target genome (upper case).
-----------------	--




Figure 2.2: Alignments done by PolyMarker. (a) input. The alternative alleles are surrounded by brackets. (b) Global search of templates in the reference contigs. (c) Selected regions around the SNP on every chromosome. (d) Sequence of found regions around the SNP. (e) Local alignment on regions around the SNP detects indels. (f) Alignment with mask and primer candidates.

Semi-specific	Homoeologous polymorphism which is found in 2 of the 3 genomes, hence it discriminates against one of the off-target genomes or when not all the homoeologous sequences were found (lower case).
Non-specific	No variation is found across homoeologues (-).
Homoeologous	The target SNP is present across different chromosomes, so candidate SNP markers on this category are not expected to be reliably identify the allele (:).
Non-homoeologous	The target SNP is not present across chromosomes, so it can be used to identify an allele (&).

PolyMarker was designed to produce SNP assays for KASP genotyping (LGC Genomics, 2013), which requires a common primer and two allele-specific primers. The common primer is selected to start on a position from a: Specific; Semi-specific or; Non-specific, on that priority. This means that the common primer will be as specific as possible in the region. For the allele-specific primers, the starting position of the primer is on the base with the SNP. To ensure that the stability of the candidate primers will be met, the putative starting positions are tested with *Primer3* (Rozen and Skaletsky, 2000).

PolyMarker was designed and validated with the markers described in section 3.8. For wheat, PolyMarker uses the contigs from Mayer et al. (2014), as deposited in Ensembl. As new releases of the wheat genome are made available, different parsers to assign the chromosome to each sequence can be added with little effort to PolyMarker.

2.2 Applications of PolyMarker

PolyMarker is not restricted to wheat or to KASP assays, the source code is flexible and can be extended for other types of analysis. On each of the following projects, PolyMarker has been adapted to design primers in species where KASP hasn't been used before, the primers are used for regular PCR amplification, or the use of KASP is not the conventional SNP calling.

2.2.1 KASP assays for public sets of SNPs

PolyMarker was used to design KASP assays for the 81,587 markers from (Wang et al., 2014), available on the PolyMarker website and in CerealsDB (Wilkinson et al., 2012). Of those markers, 40,267 were designed using the target chromosome using the genetic map published by the genetic map. Genes without a genetic position were aligned to scaffolds sorted by chromosome from the International Wheat Genome Sequencing Consortium (Mayer et al., 2014) with BLAT (Kent, 2002) and the best hit was selected as putative location. 97.5% of the assays were designed and 76% of them are semi-specific or specific, thereby improving their expected performance with respect to randomly designed primers (Table A.1). A subset of the designed assay was used to genotype a mapping population to find resistance to Fusarium head blight (Burt et al., 2015).

2.2.2 SNPs in a mutant population

PolyMarker was used to design primers to validate SNPs in a Targeted Induced Local Lesions in Genomes (TILLING) population, an approach to identify the function of genes by mutating them. To calibrate the SNP calling, KASP assays were designed to get the mutations from M_2 , M_3 and, M_5 mutants (King et al., 2015). Then primers were designed for the whole mutant population, consisting of 1,200 Cadenza (Hexaploid) and 1,535 Kronos (Tetraploid) wheat lines (Krasileva et al., submitted 2016). Genome-specific primers 172 and 80 SNP assays on 19 and 8 M_4 Cadenza and Kronos lines respectively. Of those, 71(85.5%) Kronos and 147(88.8%) of the Cadenza primers were valid assays (Tables A.4 and A.5).

2.2.3 Deletions on a mutant population

On some of the TILLING mutant lines long deletions were detected (Krasileva et al., submitted 2016). To validate the deletions it is possible to use KASP assays to produce primers that amplify homoeologues. PolyMarker was modified to search for variations across homoeologues to select a common primer that will amplify two genomes (Figure 2.3a, b). On lines without the targeted deletion, the amplification corresponds to an homozygous assay (Figure 2.3c). When a deletion is present the results of the assay look like an homozygous sample, with the intensity




Figure 2.3: KASP assays to validate homozygous deletions. (a) Primer positions for wildtype. (b) Primer positions on homozygous deletion on M_4 (c) Heterozygous amplification on wildtype, including both homoeologues. (d) Homozygous amplification on deletion line, only the non-deleted homoeologue is amplified.

of the assay towards the the conserved homoeologue (Figure 2.3d). A set of KASP assays for the the deletions and mutations located on the same chromosome where designed to validate 11 homozygous deletions on M_4 plants. In all cases the segregation of the mutations was as expected, except for a predicted heterozygous mutation that was called as homozygous. Also, all the KASP assays that contained a deletion were called homozygous, as expected. To ensure that the calls didn't come from a single cluster, 4 wildtype plants were genotyped and the markers for deletions where called as heterozygous. An example of a validated deletion, with the calls for each individual is shown on Table A.3.

2.2.4 PolyMarker public web service

To make PolyMarker accessible to the community, a web server that allow the submission of SNPs was developed. The web interface consists on two virtual machines, one with a web facing interface that stores the queries, and a dedicated node to submit jobs to an HPC cluster. The on-line interface further simplifies the design of KASP assays, a process

that used to take a couple of weeks now is done in a couple of hours. Since the release of the public service in July 2014 until August 2016, 1,739 requests to PolyMarker have been done.

2.2.5 Genotyping of *Puccinia striiformis* f. sp. *tritici* isolates.

In Hubbard et al. (2015), *Puccinia striiformis* f. sp. *tritici* (PST) isolates were sequenced and assigned to clusters, according to their genotype. The clusters are useful to monitor the changes in the pathogen population, which can be used to predict if certain wheat lines will be resistant to the isolates in the field. PolyMarker was used to design primers for PST, using the assembly PST-130 (Cantu et al., 2011). Out of 15 assays 11 can be used to identify to which cluster of isolates a sample is likely to belong, Supplemental Table A.2.

2.3 Discussion

PolyMarker is a tool that was born as part of the validation of the SNPs found in Chapter 3. Originally, the primer design was ought to be done manually, a slow, error-prone and, repetitive process. The steps require the use of several bioinformatics tools, but once I figured out the steps I decided to automate the process. Since designing genome-specific primers is a common task in wheat research and breeding, the community showed interest on the tool and I decided to refine it and make it open source. PolyMarker has been used successfully in several projects and it even allowed the novel use of KASP assays to validate long deletions in polyploids.

The current web interface of PolyMarker is limited to KASP assays, however the command line version is more flexible and has been used to design primers for PCR amplicons, capillary sequencing and on other organisms. The ideas behind PolyMarker had been taken by other projects like the scripts described in Ma et al. (2015) and the corresponding web interface, GSP (Wang et al., 2016). As new references of wheat come available, PolyMarker should be updated to work with pseudomolecules and the web interface updated accordingly.

Chapter 3

Genetic map of *Yr15* with RNA-Seq

Wheat breeding programs aim to improve the wheat lines available for production. One of the traits desired in an elite line is the resistance to pathogens, such as *Puccinia striiformis* f. sp. *tritici*, the fungi responsible of yellow rust. A source of resistance genes is are introgressions from other species, such as *Triticum diccoides*. In the University of Sydney a collection of Near Isogenic Lines (NILs) with introgressions to several Yellow Rust resistance genes on a susceptible background were developed (Wellings and McIntosh, 1998). On this chapter the NIL for the *Yr15* locus is used to produce a mapping population to improve diagnostic markers.

Line selection can be done with molecular markers that can be used to test if certain allele is present in a line, without the need to do a phenotype. To find which regions are linked to a trait the use of F_2 mapping populations is a common practice. The population is produced by crossing two homozygous parents (P_1 and P_2) with different alleles, A/A (dominant) and a/a (recessive). When the trait is dominant and has a mendelian segregation, the F_1 population show the dominant trait, as it has a copy of each allele (A/a). The F_1 is then self-crossed to and the population segregates with a ration 1:2:1, dominant:heterozygous:recessive respectively. This generates a population with a phenotype ratio of 3:1 (dominant:recessive), since the effect of the recessive allele is masked by the dominant gene (Van Ooijen and Jansen 2013; Figure 3.1).

Bulk Segregant Analysis (BSA) consists on pooling the DNA of individuals with contrasting phenotypes (Michelmore et al., 1991) on a




Figure 3.1: The cross of two homozygous parents, P_1 and P_2 , with a dominant and a recessive allele of a gene produce an heterozygous F_1 . The F_1 crossed with itself produce a segregating F_2 population with a 1:2:1 ratio ($A/A:A/a:a/a$). The upper and lower cases represent dominant and recessive alleles

segregating population. The bulks show as heterozygous except for the region that is linked to the trait of interest. This approach can be used to identify SNPs using High Throughput Sequencing, such as: exome capture (Hodges et al., 2007), RNA-Seq (Pickrell et al., 2010), whole genome resquencing (Schneeberger et al., 2009), among others.

To Call for SNPs from RNA-Seq a reference transcriptome is used as target when aligning the reads. The Bulk Frequency Ratio (BFR) methodology can work on organisms has more than one pseudo genome and that the genes are not necessarily fully characterised independently among homoeologues or paralogues, you can have in a single reference collapsing similar regions. The UniGenes database, from NCBI, contains the genes of each species, with all the variations of each gene automatically collapsed and represented as with the longest cDNA (Pontius et al., 2002). The UCW genes described in Krasileva et al. (2013) contains 94,177 models from tetraploid and hexaploid wheat, assembled and phased to separate different homoeologues. Both gene sets are complement each other, however, the UCW gene models should provide an improved alignment, since the different homoeologues aren't merged in a single model, a possible side effect of the UniGene pipeline.

Homoeologous variants, as exemplified by the G>T variant at position 181; K in consensus (Figure 3.2), will produce the same ambiguity code for both parental consensus sequences and can therefore be excluded. Real allelic SNPs between the parental genotypes, exemplified by the G>A variant at position 184; R in consensus, are distinguished by the presence in one, but not the other parental consensus sequence. The




Figure 3.2: Illustration of a non-informative homoeologous SNP (G181T) present in both parental lines, and an informative allelic SNP (G184A), only present in the resistant progenitor Avocet S + Yr15. The consensus sequences from the parental genotypes include this information in the form of ambiguity codes (K and R, respectively). In the bulks, the individual reads align across the reference sequence, with matches indicated by dots, and polymorphisms at positions 181 and 184 indicated by the corresponding nucleotide variants at those positions. The SNP index is calculated as the frequency of the informative allelic SNP in each bulk. The Bulk Frequency Ratio is the quotient of the resistant and susceptible bulk SNP Indexes. Figure previously published in Ramirez-Gonzalez et al. (2015b).




Figure 3.3: Layers of information to do *In Silico* mapping. SNPs are called from gene models. The genes and markers from genetic maps are aligned to scaffolds. The order of the markers in a genetic map can be used to sort the scaffolds.

allelic SNPs are then examined further with the alignments of the bulks to identified the SNPs that are enriched on the resistant plants. The SNP index is the proportion of times an alternative allele is observed over the coverage at certain, in the example the the susceptible bulk has an SNP index of $1/8 = 0.125$ and $6/8 = 0.75$ for the resistant bulk (Takagi et al., 2013b). traditional The BFR are then calculated by dividing the SNP Index of sample containing the target phenotype (resistance) over the sample without the trait (susceptible), on the example is $0.75/0.125 = 6$. A high BFR suggests that the SNP is linked to the target trait (Trick et al., 2012). The results of the BFR analysis on the F_2 population are discussed in Section 3.4.

There are several layers of information that can be used to add a context to the SNPs. When the SNPs are called from genes like the Uni-Genes (Pontius et al., 2002) or the UCW gene models (Krasileva et al., 2013), the location of the genes can be assigned by aligning them to a genomic reference, even if it is fragmented. A source to get the order of the scaffolds are genetic maps previously published, such as the genetic map described in Wang et al. (2014), which has the sequence of the markers available. The markers and the genes can be aligned to the scaffolds with a high percentage of identity (over 98%), to avoid them being assigned

to an homoeologue or paralogue region in a different chromosome. The use of genetic maps to sort genomic sequence is frequently used to produce pseudo-chromosomes on genome wide projects, usually with ad-hoc tools (Tang et al., 2015). Since the CSS assembly is quite fragmented the genetic maps don't have enough resolution to produce a pseudomolecule, however it is enough to sort the scaffolds in bins when several markers map to the same location. In this way, it is possible to use the scaffolds as a proxy to map the genes to their genetic position (Figure 3.3). The results of mapping the genes with SNPs to the CSS assembly and the genetic map are described in Section 3.5. For a longer description of resources available for wheat see Section 1.6.

Finally, the best candidate SNPs were selected to produce a genetic map which lead to a triplet of markers diagnostic to the target locus.

The steps described in this chapter were first published in Ramirez-Gonzalez et al. (2015b) and the results of this chapter are published in Ramirez-Gonzalez et al. (2015c).

To do:
section
talking
about
genetic
map.

To do:
Mi-
crosatel-
lites vs
SNP
markers.

3.1 Mapping population

The population was developed by crossing the resistant line Avocet + *Yr15* (*Yr15*) (Wellings and McIntosh, 1998), Figure 3.4a, to the susceptible line Avocet S (AVS), Figure 3.4b. *Yr15* is a NIL of a 6th generation Back-cross (BC) and the AVS background is highly susceptible to yellow rust, hence the resistance is conferred by the *Yr15* locus. F_2 seeds from three independent F_1 plants were sown and tissue was collected, before the fungal inoculation to avoid the effect of the response on the gene expression. The plants were challenged at the three leaf stage as it is known that *Yr15* confers resistance in seedlings (Gerechter-Amitai et al., 1989). The expected segregation on an F_2 population is 3:1 (resistant:susceptible), since *Yr15* is a dominant gene. From the 232 plants in the F_2 population that germinated, 187 were resistant and 45 were susceptible, which deviates slightly from the expected ratio ($\chi^2 = 0.049$). Segregation distortion has been shown for the same *Yr15* donor (Randhawa et al., 2009), however the decreased number of susceptible plants can be explained by escapes in the virulence essays (i.e. plants scored as resistant without the *Yr15* locus). For this study we extracted DNA from individual plants in the F_2 population and we bulked RNA on 6




Figure 3.4: Response of (a) Avocet + Yr15 and (b) Avocet when inoculated with *Puccinia striiformis* f. sp. *tritici* at the three leaf stage. (c) The phenotype of the F_2 population was used to produce 6 bulks, 3 resistant and 2 susceptible. The RNA was pooled in bulks accordingly. Adapted from (Ramirez-Gonzalez et al., 2015c)

Table 3.1: Arrangement and number of sequenced base pairs per sample.

Library	name	Bar code	Lane	Reads ($\times 10^8$ bp)
LIB1715	Bulk R1	ATCACG	1	0.77
LIB1716	Bulk R2	TAGCTT	1	1.20
LIB1717	Bulk R3	ACTTGA	2	0.96
LIB1718	Bulk S1	GGCTAC	2	1.64
LIB1719	Bulk S2	CGTACG	2	1.49
LIB1720	Bulk S3	GTGGCC	1	1.88
LIB1721	AvocetS	N/A	3	4.13
LIB1722	AvocetS + <i>Yr15</i>	N/A	4	3.99

different bulks: 3 resistant and, 3 susceptible (Figure 3.4c).

3.2 Sequencing and mapping

RNA-Seq was used to avoid sequencing the non-coding regions and reduce the search space. The sequencing of the bulks and the parents were done on a single Illumina Hi-Seq2000 each. The bulks were multiplexed and sequenced on a third of a lane each, as shown on Table 3.1. To ensure that the quality of the sequencing was good, **fastqc-0.10** (Babraham Bioinformatics, 2012) was run with its default parameters in each one of the fastq files. The GC content was around 52% in all the samples (Appendx B.2), which is expected as the sample should be of coding regions, and for wheat the reported GC content in genes is around 55%. The quality of the reads is fairly consistent, in general dropping after the base 80 across the samples (Appendix B.1).

When the analysis was started, the draft genome and the corresponding annotation were not yet released, hence gene models were used. All the samples were aligned to the Unigenes v60 (56,954 genes) and the gene models from UCW (Krasileva et al., 2013) using **BWA 0.5.9** (Li and Durbin, 2009). The alignment provided showed that a few genes were overexpressed, however we still have 22,107 and 36,808 genes, on the Unigenes and the UCW gene set respectively, with a coverage greater than 20x in the progenitor with *Yr15*. Both gene sets performed similarly in terms of the percentage of genes with reads and percentage of aligned reads. For AVS and *Yr15*, the percentage of genes with a coverage of at least 20x is 45% and 39% respectively across both references (Figure 3.5a). Since each individual bulk has a lower coverage, the susceptible

Table 3.2: Number of genes with a coverage over 20x, 10x and at least one read (>0x).

Coverage	Reference	Bulks					Bulk mixes				Progenitors		
		R1	R2	R3	S1	S2	S3	R1+R2	S1+S2	R1+R2+R3	S1+S2+S3	Yr15	AVS
20x	UCW	16,434 17%	27,871 30%	27,223 29%	32,287 34%	28,669 30%	34,898 37%	33,968 36%	41,019 44%	40,985 44%	47,507 50%	36,808 39%	42,248 45%
	UniGene v60	9,643 17%	16,182 28%	15,222 27%	19,549 34%	17,397 31%	20,567 36%	20,219 36%	25,270 44%	24,598 43%	29,052 51%	22,107 39%	25,842 45%
10x	UCW	27,371 29%	38,282 41%	37,777 40%	42,658 45%	38,999 41%	44,610 47%	43,266 46%	49,473 53%	49,182 52%	54,781 58%	46,356 49%	50,760 54%
	UniGene v60	16,201 28%	22,948 40%	22,130 39%	26,200 46%	24,130 42%	26,914 47%	26,318 46%	30,579 54%	29,857 52%	33,557 59%	28,044 49%	31,095 55%
>0x	UCW	68,302 73%	72,484 77%	72,957 77%	74,694 79%	73,290 78%	75,201 80%	74,397 79%	77,093 82%	76,715 81%	78,796 84%	76,275 81%	77,080 82%
	UniGene v60	40,717 71%	42,489 75%	42,595 75%	43,625 77%	43,059 76%	43,748 77%	43,393 76%	44,655 78%	44,364 78%	45,392 80%	43,732 77%	44,596 78%




Figure 3.5: (a) Box plot distribution of the gene coverage of the parent reads (AVS and *Yr15*) across the UCW (blue) and the UniGene (red) gene models. The dashed line represents the 209 minimum coverage required for SNP calling. The full line represents the average coverage across all gene models. (b) Percentage of genes exhibiting SNPs across references. The number of SNPs between the parent reads and the corresponding references was calculated (per 100 bp, rounded). The between-parents category corresponds to putative SNPs when comparing the consensus sequence between AVS and *Yr15*. Adapted from Ramirez-Gonzalez et al. (2015c)

and resistant reads were merged *in silico* as: (i) susceptible bulks 1 with 2 (S1 + S2) and resistant bulks 1 with 2 (R1 + R2) and (ii) all the susceptible (S1 + S2 + S3) and resistant bulks (R1 + R2 + R3). The merged samples increased the percentage of genes with coverage over 20x to 44% and 50% in the resistant and susceptible bulks (Table 3.2), which is close to the coverage from the progenitors.

3.3 SNP Calling

The SNP calling was done on positions with a coverage of at least 20x on the progenitor lines against the gene reference. The AVS progenitor had roughly 3% more genes with polymorphisms than *Yr15*, consistent with the difference in coverage, suggesting that with a higher coverage we could recover more SNPs from *Yr15*. The UniGenes have a higher number of SNPs because the UCW gene models have a higher number of monomorphic genes when compared to the UniGenes. (Figure 3.5b; Table 3.3). The difference in the number of relative monomorphic SNPs between references can be explained by the fact that the UniGenes have homoeologues can be represented as a single sequence, as opposed to the UCW set which are homoeologue-specific, improving the mapping to the

Table 3.3: Count of SNPs per 100 bp on genes with at least 20x coverage.

SNPs per 100bp	UCW			UniGene v60		
	AVS	AVS+ <i>Yr15</i>	Between progenitors	AVS	AVS+ <i>Yr15</i>	Between progenitors
0	67,389 71.6%	70,338 74.7%	81,921 87.0%	36,210 63.6%	38,339 67.3%	47,097 82.7%
1	16,111 17.1%	14,770 15.7%	10,107 10.7%	10,058 17.7%	9,175 16.1%	8,061 14.2%
2	8,904 9.5%	7,676 8.2%	1,893 2.0%	8,529 15.0%	7,648 13.4%	1,621 2.9%
3	1,517 1.6%	1,192 1.3%	215 0.2%	1,870 3.3%	1,568 2.8%	59 0.3%
4+	253 0.3%	198 0.2%	38 0.0%	287 0.5%	224 0.4%	16 0.0%




Figure 3.6: Gene models with putative SNPs in common between the UCW and UniGenes reference. The intersection represents the genes that are common in both sets. Adapted from Ramirez-Gonzalez et al. (2015c)

correct homoeologue in the genes from the UCW set over the UniGenes.

Both gene sets were done from varieties different to AVS and are likely to be incomplete, hence we set a low threshold of at least 20% of the observed nucleotides on any position to call an SNP. To represent cases where more than one consensus base is called we use International Union of Pure and Applied Chemistry (IUPAC) codes (Cornish-Bowden (1985); Section 1.5.1; Figure 3.2). To focus the analysis on informative SNPs, the common varietal SNPs and variations between homoeologues were removed by finding the cases when the consensus call on both progenitors is the same. The SNPs that are unique to a single parental were examined in detail. There are 66,426 putative SNPs across 16,022 (17%) UCW genes and 52,262 SNPs on 11,056 UniGenes (19.4%; Figure 3.6).

The high number of genes with SNPs was unexpected as a BC6 NIL used for an F_2 mapping population expects to have $< 1\%$ of the genetic background segregating. The both sets of gene models were aligned

Table 3.4: Number of genes with SNPs assigned to the wheat chromosome arm CSS scaffolds (Mayer et al., 2014) using the best hit from BLAT (Kent, 2002)

Wheat Chromosome Arm	UCW (94,177)	UniGene v60 (56,954)	Total (151,131)
1AL	3,251 (3.45%)	1,404 (2.47%)	4,655 (3.08%)
1AS	1,366 (1.45%)	560 (0.98%)	1,926 (1.27%)
1BL	2,610 (2.77%)	1,280 (2.25%)	3,890 (2.57%)
1BS	1,487 (1.58%)	693 (1.22%)	2,180 (1.44%)
1DL	997 (1.06%)	1,057 (1.86%)	2,054 (1.36%)
1DS	753 (0.80%)	687 (1.21%)	1,440 (0.95%)
2AL	3,491 (3.71%)	1,460 (2.56%)	4,951 (3.28%)
2AS	2,305 (2.45%)	974 (1.71%)	3,279 (2.17%)
2BL	3,658 (3.88%)	1,546 (2.71%)	5,204 (3.44%)
2BS	2,790 (2.96%)	1,139 (2.00%)	3,929 (2.60%)
2DL	1,098 (1.17%)	1,069 (1.88%)	2,167 (1.43%)
2DS	796 (0.85%)	833 (1.46%)	1,629 (1.08%)
3AL	2,135 (2.27%)	978 (1.72%)	3,113 (2.06%)
3AS	1,543 (1.64%)	718 (1.26%)	2,261 (1.50%)
3B	6,559 (6.96%)	2,839 (4.98%)	9,398 (6.22%)
3DL	915 (0.97%)	938 (1.65%)	1,853 (1.23%)
3DS	412 (0.44%)	450 (0.79%)	862 (0.57%)
4AL	3,393 (3.60%)	1,335 (2.34%)	4,728 (3.13%)
4AS	2,011 (2.14%)	817 (1.43%)	2,828 (1.87%)
4BL	2,119 (2.25%)	898 (1.58%)	3,017 (2.00%)
4BS	1,946 (2.07%)	892 (1.57%)	2,838 (1.88%)
4DL	1,069 (1.14%)	945 (1.66%)	2,014 (1.33%)
4DS	800 (0.85%)	699 (1.23%)	1,499 (0.99%)
5AL	2,640 (2.80%)	1,132 (1.99%)	3,772 (2.50%)
5AS	963 (1.02%)	407 (0.71%)	1,370 (0.91%)
5BL	5,324 (5.65%)	1,943 (3.41%)	7,267 (4.81%)
5BS	1,360 (1.44%)	591 (1.04%)	1,951 (1.29%)
5DL	2,067 (2.19%)	1,688 (2.96%)	3,755 (2.48%)
5DS	620 (0.66%)	614 (1.08%)	1,234 (0.82%)
6AL	2,397 (2.55%)	896 (1.57%)	3,293 (2.18%)
6AS	2,285 (2.43%)	936 (1.64%)	3,221 (2.13%)
6BL	1,564 (1.66%)	820 (1.44%)	2,384 (1.58%)
6BS	1,308 (1.39%)	731 (1.28%)	2,039 (1.35%)
6DL	1,399 (1.49%)	1,050 (1.84%)	2,449 (1.62%)
6DS	870 (0.92%)	680 (1.19%)	1,550 (1.03%)
7AL	1,918 (2.04%)	849 (1.49%)	2,767 (1.83%)
7AS	1,717 (1.82%)	764 (1.34%)	2,481 (1.64%)
7BL	1,592 (1.69%)	776 (1.36%)	2,368 (1.57%)
7BS	1,239 (1.32%)	713 (1.25%)	1,952 (1.29%)
7DL	2,040 (2.17%)	1,301 (2.28%)	3,341 (2.21%)
7DS	1,224 (1.30%)	1,016 (1.78%)	2,240 (1.48%)
Assigned	80,031 (84.98%)	41,118 (72.20%)	121,149 (80.16%)

Table 3.5: Total number of SNPs scored in parents, individual bulks and in silico merged bulks.

Gene set	$\frac{R1}{S1}$	$\frac{R2}{S2}$	$\frac{R3}{S3}$	$\frac{R1+R2}{S1+S2}$	$\frac{R1+R2+R3}{S1+S2+S3}$	SNPs in parents
UCW	16,269 24.49%	29,703 44.72%	31,891 48.01%	44,224 66.58%	64,522 97.13%	66,426
UniGene v60	15,261 29.20%	25,143 48.11%	24,548 46.97%	35,698 68.31%	49,738 95.17%	52,262

with BLAT (Kent, 2002) to the Chinese Spring Chromosome arm survey sequence (CSS; Mayer et al. 2014); the alignment resulted on 80,031 (85.0%) UCW gene models and 41,118 (72.2%) UniGenes assigned to a chromosome arm (Table 3.4). The SNPs found in the mapped genes are evenly distributed across all the chromosomes (Figure 3.10a), suggesting that the Avocet S (JIC, UK) used as parent in the F_2 is different to the Avocet S used for the *Yr15* NIL development (University of Sydney, Australia).

To confirm that the Avocet S seed stocks from JIC are distinct to the stocks in Sydney DNA from both stocks was procured and compared with the iSelect 90k wheat SNP chip. Between two independent Avocet S seeds from JIC only 58 out of 71,972 (0.08%) valid assays were polymorphic. Nonetheless, there are over 5,000 ($> 7.5\%$) assays with polymorphisms between JIC-Avocet S and Avocet S from Sydney. The difference was not expected originally, but considering that the Avocet S seeds are coming from different stocks and the fact that in both countries commercial varieties with the same name had been released, it is not surprising.

3.4 Bulk Frequency Ratios

The objective was to find the SNPs enriched on each bulk and hence linked to the phenotype, variations from *Yr15* to resistance and from AVS to susceptibility in the segregating population. Across individual bulks, it was possible to score between 15,261 (24.5%) to 31,891 (48.0%) SNPs across both reference sets. On the *in silico* mixes over 95% of SNPs were scored (Table 3.5), suggesting that the coverage of individual bulks is not enough to score all the SNPs. The scoring was done with the Bulk Frequency Ratio (Trick et al. 2012; Figure 3.2; Section 3.9.2), which has a value that increases as the *Yr15* allele is observed more times relatively




Figure 3.7: Effect of BFR threshold on the number of SNPs across the short arm of chromosome group 1. Figure previously published in Ramirez-Gonzalez et al. (2015c).

to the AVS allele.

When increasing the minimum BFR threshold, enrichment of SNPs was observed in the short arm of the group 1 chromosomes (1S). Without taking in account the BFR, 3.6% of the SNPs are located in the 1S group, similar to the number of SNPs located in other groups 3.4. However, when increasing the threshold (between $BFR > 5$ and $BFR > 7$) the relative number of SNPs in group 1S increases. After $BFR > 7$ the gains in relative enrichment only improves marginally, but the number of called SNPs is reduced (Table 3.6; Figure 3.7). For that reason, SNPs with a $BFR > 6$ were selected for further validation. The method described by Trick et al. (2012) was extended by including cases where there is a complete lack of coverage in one of the samples ($BFR = \infty$), which is an ideal case where the linkage between the SNP and the phenotype is perfect. A total of 1,582 SNPs across 1,173 genes had a $BFR > 6$.

Table 3.6: SNPs in chromosome group 1S vs total number of SNPs with a minimum BFR from 0 to 10. AVS: SNPs coming from Avocet S. Yr15: SNPs coming from Avocet + Yr15.

Min BFR	Gene Set	R1/S1 Yr15	R1/S1 AVS	R2/S2 Yr15	R2/S2 AVS	R3/S3 Yr15	R3/S3 AVS	S1+2/ R1+2 Yr15	S1+2/ R1+2 AVS	S1+S2+S3/ R1+R2+R3 Yr15	S1+S2+S3/ R1+R2+R3 AVS
0	UCW	308/8,049 (3.83%)	305/8,220 (3.71%)	505/14,121 (3.58%)	556/15,582 (3.57%)	532/14,875 (3.58%)	623/17,016 (3.66%)	670/18,760 (3.57%)	885/25,464 (3.48%)	860/24,026 (3.58%)	1,505/40,496 (3.72%)
	UniGene v60	307/7,823 (3.92%)	299/7,438 (4.02%)	428/12,409 (3.45%)	421/12,734 (3.31%)	427/12,050 (3.54%)	415/12,498 (3.32%)	536/15,672 (3.42%)	595/20,026 (2.97%)	712/19,358 (3.68%)	901/30,380 (2.97%)
1	UCW	214/4,415 (4.85%)	194/4,108 (4.72%)	325/7,603 (4.26%)	314/7,374 (4.26%)	365/7,920 (4.61%)	415/8,850 (4.69%)	426/10,122 (4.21%)	494/12,185 (4.05%)	539/13,037 (4.13%)	842/19,466 (4.33%)
	UniGene v60	207/4,474 (4.63%)	194/3,630 (5.34%)	269/6,649 (4.05%)	269/6,193 (4.34%)	279/6,511 (4.29%)	272/6,436 (4.23%)	329/8,704 (3.78%)	369/9,343 (3.95%)	446/10,860 (4.11%)	541/14,226 (3.80%)
2	UCW	92/651 (14.13%)	75/671 (11.18%)	142/1,372 (10.35%)	111/1,101 (10.08%)	147/1,162 (12.65%)	149/1,411 (10.56%)	167/1,324 (12.61%)	163/1,478 (11.03%)	194/1,370 (14.16%)	207/1,765 (11.73%)
	UniGene v60	77/568 (13.56%)	58/527 (11.01%)	101/1,017 (9.93%)	81/720 (11.25%)	105/775 (13.55%)	84/867 (9.69%)	122/991 (12.31%)	116/973 (11.92%)	145/1,030 (14.08%)	132/1,210 (10.91%)
3	UCW	78/299 (26.09%)	45/295 (15.25%)	118/646 (18.27%)	70/409 (17.11%)	123/577 (21.32%)	85/494 (17.21%)	145/673 (21.55%)	98/563 (17.41%)	168/768 (21.88%)	122/665 (18.35%)
	UniGene v60	65/254 (25.59%)	26/186 (13.98%)	87/499 (17.43%)	54/294 (18.37%)	93/379 (24.54%)	48/315 (15.24%)	107/525 (20.38%)	66/379 (17.41%)	133/617 (21.56%)	78/489 (15.95%)
4	UCW	75/232 (32.33%)	28/160 (17.50%)	109/484 (22.52%)	44/217 (20.28%)	105/416 (25.24%)	44/246 (17.89%)	134/539 (24.86%)	53/277 (19.13%)	149/640 (23.28%)	64/323 (19.81%)
	UniGene v60	63/192 (32.81%)	17/104 (16.35%)	83/390 (21.28%)	29/155 (18.71%)	82/288 (28.47%)	29/173 (16.76%)	104/431 (24.13%)	40/214 (18.69%)	127/519 (24.47%)	29/266 (10.90%)
5	UCW	69/202 (34.16%)	19/108 (17.59%)	95/416 (22.84%)	33/138 (23.91%)	96/354 (27.12%)	23/143 (16.08%)	127/477 (26.62%)	28/175 (16.00%)	140/580 (24.14%)	42/222 (18.92%)
	UniGene v60	58/163 (35.58%)	11/70 (15.71%)	76/337 (22.55%)	14/102 (13.73%)	70/228 (30.70%)	20/112 (17.86%)	100/389 (25.71%)	23/146 (15.75%)	118/469 (25.16%)	21/178 (11.80%)
6	UCW	65/179 (36.31%)	12/85 (14.12%)	86/380 (22.63%)	22/98 (22.45%)	87/299 (29.10%)	11/94 (11.70%)	122/429 (28.44%)	21/130 (16.15%)	126/514 (24.51%)	29/165 (17.58%)
	UniGene v60	57/151 (37.75%)	7/48 (14.58%)	73/300 (24.33%)	6/71 (8.45%)	65/191 (34.03%)	13/84 (15.48%)	98/358 (27.37%)	20/122 (16.39%)	115/439 (26.20%)	16/143 (11.19%)
7	UCW	58/161 (36.02%)	11/73 (15.07%)	77/340 (22.65%)	13/74 (17.57%)	73/248 (29.44%)	7/69 (10.14%)	116/393 (29.52%)	20/111 (18.02%)	114/468 (24.36%)	22/143 (15.38%)
	UniGene v60	56/132 (42.42%)	4/37 (10.81%)	68/273 (24.91%)	5/58 (8.62%)	60/171 (35.09%)	9/64 (14.06%)	94/334 (28.14%)	18/103 (17.48%)	113/412 (27.43%)	16/124 (12.90%)
8	UCW	58/149 (38.93%)	10/62 (16.13%)	68/310 (21.94%)	12/59 (20.34%)	66/214 (30.84%)	6/56 (10.71%)	104/359 (28.97%)	17/102 (16.67%)	108/429 (25.17%)	16/119 (13.45%)
	UniGene v60	55/126 (43.65%)	3/33 (9.09%)	64/255 (25.10%)	5/50 (10.00%)	55/150 (36.67%)	9/55 (16.36%)	91/313 (29.07%)	14/89 (15.73%)	105/376 (27.93%)	15/108 (13.89%)
9	UCW	54/135 (40.00%)	8/53 (15.09%)	63/289 (21.80%)	8/51 (15.69%)	61/182 (33.52%)	5/49 (10.20%)	100/331 (30.21%)	15/91 (16.48%)	100/387 (25.84%)	13/106 (12.26%)
	UniGene v60	53/117 (45.30%)	1/30 (3.33%)	62/244 (25.41%)	5/46 (10.87%)	50/136 (36.76%)	9/48 (18.75%)	88/291 (36.76%)	13/83 (15.66%)	97/345 (28.12%)	12/99 (12.12%)
10	UCW	52/126 (41.27%)	8/50 (16.00%)	62/279 (22.22%)	8/50 (16.00%)	56/165 (33.94%)	4/45 (8.89%)	96/309 (33.94%)	14/82 (17.07%)	91/355 (25.63%)	13/100 (13.00%)
	UniGene v60	50/105 (47.62%)	1/28 (3.57%)	60/226 (26.55%)	5/39 (12.82%)	43/119 (36.13%)	7/45 (15.56%)	85/272 (31.25%)	13/82 (15.85%)	92/318 (28.93%)	12/97 (12.37%)




Figure 3.8: Location of SNPs with $BFR > 6$ according to the best alignment of the UniGene (red) and UCW (blue) gene models to the flow-sorted group 1 chromosomes from the Chinese Spring Survey sequence (CSS) (Mayer et al., 2014). Figure adapted from Ramirez-Gonzalez et al. (2015c).

3.5 *In silico* mapping

From the mapped SNPs with a $BFR > 6$, 872 of 1470 ($\sim 60\%$) were assigned to the chromosomes in group 1 of hexaploid wheat, being the only group with more than 4% of the SNPs assigned to it (Table 3.7). From the group 1, the B genome contained the higher proportion of SNPs mapped (54%), having 255 (54%) and 214 (46%) assigned to the long and short arms respectively (Figure 3.8). This results are expected since previous studies have located *Yr15* near the centromere in the short arm of chromosome 1B and, the *Yr15* introgression contains regions from the long and short arm from *T. diccoides* (Murphy et al., 2009; Peng et al., 2000; Sun et al., 1997).

The CSS assembly was used as a common reference between the reference genes and the SNPs 40,266 SNP markers published at the time when this analysis was done (Wang et al., 2014) to locate the SNPs with a $BFR > 6$ (including $BFR = \infty$) in a genomic position (Figures 3.9, 3.10). From the 1,582 SNPs across 1,173 genes, only 678 SNPs (43%, 474 genes) were successfully located in the genetic map. Since the CSS assembly is quite fragmented, the low percentage of located SNPs can be because not all candidate SNPs had a corresponding scaffold that has at least one of the 40,266 markers in the genetic map. Even if the number of located SNPs was not enough to give a position for over 50% of the SNPs from the parental line, the resolution of the genetic position SNPs that were assigned improved over just having the chromosome arm information from the CSS assembly. The mapping position further confirmed an enrichment of SNPs near the centromere of chromosome 1B with 325 out of 678 SNPs. Furthermore, 311 of those were located within an interval of 30cM (Figures 3.10b, 3.9a).

Studies in diploid organisms using QTL-Seq (Takagi et al., 2013a) or other NGS-enable genetic approaches (James et al., 2013) have shown

Table 3.7: SNP and genes with BFR i 6 mapping to each of the chromosomes from the CSS assemblies. The chromosome assignment on the "Genetically mapped" column correspond to the map published in Wang et al. (2014)).

Reference Chromosome	CSS assemblies				Genetically mapped			
	UCW gene models		UniGene v60		UCW gene models		UniGene v60	
	SNPs	Genes	SNPs	Genes	SNPs	Genes	SNPs	Genes
1AL	113	13.15%	79	12.29%	78	10.79%	50	9.43%
1AS	26	3.03%	21	3.27%	20	2.77%	17	3.21%
1BL	157	18.28%	110	17.11%	98	13.55%	64	12.08%
1BS	120	13.97%	74	11.51%	94	13.00%	44	8.30%
1DL	30	3.49%	21	3.27%	58	8.02%	47	8.87%
1DS	40	4.66%	25	3.89%	38	5.26%	24	4.53%
2AL	22	2.56%	20	3.11%	14	1.94%	12	2.26%
2AS	11	1.28%	11	1.71%	10	1.38%	7	1.32%
2BL	17	1.98%	15	2.33%	18	2.49%	17	3.21%
2BS	11	1.28%	10	1.56%	12	1.66%	7	1.32%
2DL	2	0.23%	2	0.31%	15	2.07%	10	1.89%
2DS	0	0.00%	0	0.00%	5	0.69%	3	0.57%
3AL	7	0.81%	7	1.09%	2	0.28%	2	0.38%
3AS	1	0.12%	1	0.16%	4	0.55%	4	0.75%
3B	31	3.61%	26	4.04%	28	3.87%	24	4.53%
3BL	0	0.00%	0	0.00%	0	0.00%	0	0.00%
3BS	0	0.00%	0	0.00%	0	0.00%	0	0.00%
3DL	7	0.81%	6	0.93%	2	0.28%	2	0.38%
3DS	1	0.12%	1	0.16%	2	0.28%	2	0.38%
4AL	18	2.10%	15	2.33%	6	0.83%	6	1.13%
4AS	5	0.58%	5	0.78%	6	0.83%	5	0.94%
4BL	11	1.28%	10	1.56%	6	0.83%	6	1.13%
4BS	6	0.70%	5	0.78%	13	1.80%	10	1.89%
4DL	4	0.47%	4	0.62%	5	0.69%	5	0.94%
4DS	2	0.23%	2	0.31%	5	0.69%	4	0.75%
5AL	7	0.81%	5	0.78%	3	0.41%	3	0.57%
5AS	1	0.12%	1	0.16%	2	0.28%	2	0.38%
5BL	31	3.61%	28	4.35%	14	1.94%	14	2.64%
5BS	7	0.81%	5	0.78%	6	0.83%	5	0.94%
5DL	8	0.93%	7	1.09%	15	2.07%	14	2.64%
5DS	4	0.47%	3	0.47%	6	0.83%	5	0.94%
6AL	22	2.56%	17	2.64%	9	1.24%	7	1.32%
6AS	8	0.93%	8	1.24%	11	1.52%	10	1.89%
6BL	7	0.81%	6	0.93%	3	0.41%	2	0.38%
6BS	7	0.81%	5	0.78%	2	0.28%	2	0.38%
6DL	11	1.28%	10	1.56%	7	0.97%	7	1.32%
6DS	5	0.58%	3	0.47%	2	0.28%	2	0.38%
7AL	9	1.05%	8	1.24%	7	0.97%	6	1.13%
7AS	5	0.58%	5	0.78%	8	1.11%	7	1.32%
7BL	10	1.16%	10	1.56%	4	0.55%	4	0.75%
7BS	3	0.35%	3	0.47%	4	0.55%	4	0.75%
7DL	15	1.75%	10	1.56%	12	1.66%	12	2.26%
7DS	8	0.93%	4	0.62%	6	0.83%	6	1.13%
Unmapped	49	5.70%	35	5.44%	63	8.71%	46	8.68%
Mapped	810	94.30%	608	94.56%	660	91.29%	484	91.32%
					460	53.55%	358	55.68%
					399	46.45%	285	44.32%
					444	61.41%	341	64.34%
					279	38.59%	189	35.66%




Figure 3.9: (a) Number of SNPs with $BFR > 6$ per cM in chromosome 1B. (b) BFRs of mapped genes with SNPs on chromosome 1B. The area of the circle represents the number of SNPs clustered by location (windows size: 10 cM) and BFR (window size: 5cM). R11 is the only marker near the *Yr15* locus that had a corresponding position in the genetic map. The percentage of genes with SNPs per cM is also illustrated based on UCW (blue) and UniGene (red) gene models. The centromere is imputed by the centre of a window of 10 cM where the short arm switches to the long in the genetic map. BFRs correspond to those from the mixed in silico bulk S1 + S2 + S3/R1 + R2 + R3. Adapted from (Ramirez-Gonzalez et al., 2015c).

smooth curves with a defined peak in the region linked to the studied trait. In practice, we only observe clusters of SNPs with enriched BFRs near the centromere of chromosome 1B (Figures 3.9a, 3.10b).

The location of the clusters with an enrichment of SNPs near the centromere is not expected on a random selection of genes, as the gene density increases with the distance to the centromere (Akhunov et al., 2003). This suggests that the experiment was successful on finding SNPs linked to *Yr15*. There are several factor that prevent a clear peak; like the biases induced by the differential expression, the fragmented reference sequence with scaffolds that are not long enough to go across genetic positions. Since there are several SNPs with a high BFR and the genetic map is not enough to locate a single region linked to *Yr15*, multiple criteria was needed to prioritise SNPs that were more likely to yield on successful genetic markers.

3.6 Assay selection

Three independent criteria were use to prioritize the SNPs for marker development and validation:

High BFR. SNPs with a $BFR > 6$ in at least two independent bulk replicates or in either of the *in silico* mixes were selected to ensure consistency and recover SNPs with a low coverage on a particular bulk.

Group 1S. SNPs that are in CSS scaffolds in the short arm of chromosome group 1 were selected. This is to be consistent with the *in silico* genetic map and with previous studies (Murphy et al., 2009; Peng et al., 2000; Sun et al., 1997).

***Yr15* parent.** The SNPs should originate from the *Yr15* parent to ensure that the SNP is coming from the *T. diccocooides* introgression and not from a SNP in the AVS genetic background, who would be less useful in breeding programs with a different background.

Only SNPs meeting the three criteria were selected for further analysis.

With the multiple criteria the number of genes with a putative SNP went down from $> 27,000$ to just 175; 77 and 98 from the UniGene and




Figure 3.10: Genetic location of genes with SNPs between AVS and Yr15. The colour scale indicates the percentage of genes with SNPs per centi-Morgan (cM) across the 21 wheat chromosomes. The location of the genes was determined by the best alignment to the CSS scaffolds, and the location of these was determined by their position on a genetic map (Wang et al., 2014) (a). All the SNPs between progenitors. Note the lack of enrichment across any individual chromosome. (b) SNPs with BFR > 6. Note the enrichment in Chromosome 1B




Figure 3.11: Selection criteria for marker design. Venn diagrams based on the three selection criteria (SNP in the short arm of chromosome group 1; SNP has a $BFR > 6$; and SNP is from the *Yr15* parent) for the UCW (blue) and UniGene (red) gene models. The centre diagram shows the intersection between common genes matching all three criteria across both data sets. Note that the numbers are not directly additive as in cases, multiple models from one reference set will relate to a single gene model in the other values. Published in (Ramirez-Gonzalez et al., 2015c)

UCW gene sets respectively. The selected genes from both references were aligned between references, as they come from independent sources an overlap in the selection between them is expected and, as expected, around half of the genes between gene sets overlap (Figure 3.11). The 50 SNPs with the highest BFRs, out of the 175 genes, were selected for validation, 15 of them were redundant between references, resulting on 35 SNPs to validate.

The separate bulks and the *in silico* mixes were evaluated in detail to understand the behaviour and value of having multiple bulks. The initial expectation was that as the number of SNPs with $BFR = \infty$ should drop in the mixes, as the improved coverage should reduce the instances were the absence of an allele is because of the lack of coverage on a particular sample. However, the opposite happened, the additional coverage in the *in silico* mixes recovered SNPs in genes with a low expression at the time of the sampling (Figure 3.12). Some SNPs were present across all the samples, however the value of the BFR changed depending on the sample(marker R5). On some cases a SNP are missing in an individual bulk, but present in the rest of them and in the mixes (marker R8). The main reason affecting the scoring is the coverage in the sample for each particular gene, hence an strategy with a consistent coverage would be preferred for this kind of analysis. Previous studies have shown that a




Figure 3.12: Bulk frequency ratio (BFR) of selected SNPs across the individual bulks and in silico mixes (UCW, red; UniGene, blue). The dotted line represents the BFR threshold of 6 (logarithmic scale). Left: Distribution of the BFRs for each selection criteria and the selected SNPs for validation. The circles on the top of each plot represent the percentage of SNPs with $BFR = \infty$. The Selection may include SNPs with $BFR < 6$ when the same SNP has a higher score on the complementing reference (ie. $BFR > 6$ on UCW, but $BFR < 6$ on UniGenes). Right: The BFR values of selected SNPs were sorted in descending order across the different bulks and according to their origin. Validated SNPs are indicated by open triangles, and SNPs corresponding to markers R5, R8 and R11 are labelled across different bulks and mixes. Note that some SNPs are below the threshold in a specific bulk as they meet the BFR criteria across others.

Table 3.8: Number of genes (and SNPs) with a unique hit ($> 99\%$ sequence identity) to a single wheat survey sequence scaffold.

Chromosome 1		All SNPs		BFR >6		% BFR >6	
		SNP	Genes	SNP	Genes	SNPs	Genes
UCW	Unique	5,283	1,245	311	214	5.89%	17.19%
	Total	8,086	1,954	486	330	6.01%	16.89%
	Percentage	65.34%	63.72%	63.99%	64.85%		
UniGene	Unique	3,687	745	213	139	5.78%	18.66%
	Total	6,422	1,318	386	246	6.01%	18.66%
	Percentage	57.41%	56.53%	49.17%	56.07%		
UCW +	Unique	8,970	1,990	524	353	5.84%	17.74%
	Total	14,508	3,272	872	576	6.01%	17.60%
	Percentage	61.83%	60.82%	60.09%	61.28%		

All SNPs		All SNPs		BFR >6		% BFR >6	
		SNP	Genes	SNP	Genes	SNPs	Genes
UCW	Unique	39,247	9,585	481	368	1.23%	3.84%
	Total	66,426	16,022	859	643	1.29%	4.01%
	Percentage	59.08%	59.82%	56.00%	57.23%		
UniGene	Unique	27,292	5,698	344	252	1.26%	4.42%
	Total	52,262	11,056	723	530	1.38%	4.79%
	Percentage	52.22%	51.54%	47.58%	47.55%		
UCW +	Unique	66,539	15,283	825	620	1.24%	4.06%
	Total	118,688	27,078	1,582	1,173	1.33%	4.33%
	Percentage	56.06%	56.44%	52.15%	52.86%		

coverage of $< 5x$ is enough to call for SNPs in model organisms with a high-quality reference (Schneeberger and Weigel, 2011). However, the results on this study are in line with other studies using populations for SNP calling (Abe et al., 2012; Takagi et al., 2013a). The non-uniform distribution of the coverage in RNA-Seq experiments affects the number of reads that can be used to call for SNPs, specially on genes with a low expression level (Mortazavi et al., 2008).

Around 60% of the gene models, across both references, had a unique hit with $> 99\%$ sequence identity to a single CSS scaffold (Table 3.8). This is likely because there is no unique homoeologue in the gene models, leading to reads, from two different homoeologues, mapping to the same region. To reduce the number of spurious SNPs we used IUAPC ambiguity codes (Section 1.5.1, Cornish-Bowden (1985)) when two different alleles were observed. This had as side effect that in order to keep only high confidence SNPs we required a higher coverage ($> 20x$). On the original study introducing the BFR in tetraploid wheat, the authors show that increasing the coverage, from $8x$ to $16x$, reduces the putative




Figure 3.13: KASP output from the wheat variety panel with (Ochre, Boston, Cortez) and without (Robigus, Cadenza and Shamrock) *Yr15*. Marker R2 (a) is monomorphic while R8 (b) is polymorphic between varieties known to carry the gene. Marker R8 results for the F₂ population (c) showing three distinct clusters. The central cluster (light green) is comprised of heterozygous individuals, whereas clusters near the axes are homozygous for either AVS (VIC; orange) or *Yr15* (FAM; dark green).

SNPs by 60%, but the validated SNPs increase from 57% to 83% (Trick et al., 2012). Hence, a compromise between increasing the minimum coverage at the cost of reducing the SNP candidates has to be reached in line with the objectives and available resources for a particular study.

3.7 SNP Validation

KASP assays were designed to validate and generate a genetic map of the *Yr15* locus for the 35 selected SNPs. To automate the design of genome-specific primers for polyploid organisms PolyMarker was developed (Chapter 2). Out of the 35 assays to design, 17 were designed as specific, 9 as semi-specific to chromosome 1BS, and 9 were not specific because there was no information for the homoeologues on the CSS scaffolds. PolyMarker also identified putative homoeologous variants (between genomes, as opposed to between varieties) that were in the list of candidate SNPs, but were not identified previously (Figure 2.2f; Table 3.9).

To validate if the 35 SNPs were polymorphic across the parents and, diagnostic to *Yr15* we tested them in the progenitors plus six commercial varieties, three containing *Yr15* (Ochre, Boston and, Cortez) and three without it (Shamrock, Robigus and, Cadenza). Two of the lines without *Yr15* have *T. diccoides* in their pedigree (Shamrock and Robigus), as

Table 3.9: Primer details for the markers to validate.

Assay ID	Polymorphism_type	AV'S-specific primer	Yr15-specific primer	Common primer	Specificity	Orientation
R1	non-homeologous	aactgtaattggtgcagCgG	aactgtaattggtgcagCgG	ttcaggataaacAggagatGT	chromosome_semispecific	reverse
R2	non-homeologous	acatcaattcttcaggaaactcttaC	acatcaattcttcaggaaactcttaT	gcacagcttctcgtgttcTT	chromosome_specific	forward
R3	non-homeologous	acgtggagaacttagatgcG	acgtggagaacttagatgcC	cccttiaggcgcaccaatT	chromosome_semispecific	reverse
R4	non-homeologous	agactcttttgggcagtggtG	agactcttttgggcagtggtT	ccctgggcagctattctcT	chromosome_semispecific	forward
R5	non-homeologous	agtcaacttggattacacigaagtT	agtcaacttggattacacigaagtC	agataacacatgaacatactgatG	chromosome_specific	reverse
R6	non-homeologous	caagatgaagatgaaggagaatataT	caagatgaagatgaaggagaatataG	gCttgacacctgtaatcatactcG	chromosome_semispecific	forward
R7	non-homeologous	caccaccaTggaggccaC	caccaccaTggaggccaT	cgcctggtgtagtgcgG	chromosome_specific	forward
R8	non-homeologous	cagatcccccgggtctctcaA	cagatcccccgggtctctcaA	cccccaaatgatcagaata	chromosome_inspecific	reverse
R9	non-homeologous	caggtctgaatgatcC	caggtctgaatgatcT	cggctatcttcaggctctgt	chromosome_inspecific	reverse
R10	non-homeologous	cattcgtcgccctctacG	cattcgtcgccctctacA	tcttaactatatagtcagctcAC	chromosome_specific	reverse
R11	non-homeologous	ccattctgatcaaggtcactgtcG	ccattctgatcaaggtcactgtcA	ttctgtiaTggcaaCgggagC	chromosome_specific	reverse
R12	homeologous	cttagccagtgaaacAaggcC	cttagccagtgaaacAaggcT	ggcgtgttggttacCgtggagG	chromosome_specific	reverse
R14	non-homeologous	gacTacAaggtgcatcccC	gacTacAaggtgcatcccT	ctgcctgcctcagtcgTaT	chromosome_specific	forward
R15	homeologous	gactagggtctaccAtgttgA	gactagggtctaccAtgttgC	agccctgCtaacaatggcaA	chromosome_specific	reverse
R16	non-homeologous	gatgtaagcTAgtactggCgC	gatgtaagcTAgtactggCgT	tgcacctgatcttttagcagC	chromosome_semispecific	reverse
R17	non-homeologous	geaAcaacaaCaaCaaagtggG	geaAcaacaaCaaCaaagtggC	ctctcaactcgtgtgtgtgtG	chromosome_specific	forward
R19	non-homeologous	gccgatattttaattcgtccaG	gccgatattttaattcgtccaA	agagcactgatgatgaccC	chromosome_specific	reverse
R20	non-homeologous	gctgtatctctcttgaanaaggcT	gctgtatctctcttgaanaaggcC	ttaggcatgtcagaatagtagaaaa	chromosome_semispecific	forward
R21	non-homeologous	gcttcaaacatgccgctG	gcttcaaacatgccgctT	cggctttttcaaccaggG	chromosome_semispecific	forward
R22	homeologous	gctTgtCttaaagccAttccA	gctTgtCttaaagccAttccG	gcctatcgttCgttaactcttaacT	chromosome_specific	reverse
R23	non-homeologous	gctttaggcactatggattcAcC	gctttaggcactatggattcAcT	caggtttctgttcgacctcA	chromosome_specific	forward
R24	non-homeologous	ggaggtctcacacgctctT	ggaggtctcacacgctctG	ctccaaaagaggggcattcattT	chromosome_semispecific	forward
R25	non-homeologous	gggttctcactgcgcC	gggttctcactgcgcT	ctctTtgcacatcgccagc	chromosome_inspecific	reverse
R26	non-homeologous	gtCttgcCggcacCacC	gtCttgcCggcacCacT	agtgatcttcgcgactcgc	chromosome_inspecific	forward
R28	non-homeologous	tagatgagaccttggacCggA	tagatgagaccttggacCggG	cagtcactaatgcggaacattcA	chromosome_semispecific	reverse
R29	non-homeologous	TatggtGtggccTtccccG	TatggtGtggccTtccccA	cagctcgtcgtgatgaactG	chromosome_specific	forward
R30	non-homeologous	tcagcagcccttttaacccaA	tcagcagcccttttaacccaT	agtaaatggcgcaggtgt	chromosome_inspecific	reverse
R31	homeologous	tcattcatgtatatGaaTccaagcC	tcattcatgtatatGaaTccaagcA	tcacgccctgcaacAttcaaaT	chromosome_specific	reverse
R32	homeologous	tccaatcttatggcttctgtctG	tccaatcttatggcttctgtctT	caggtgatgtagatgctgagaC	chromosome_semispecific	reverse
R33	non-homeologous	tcttctcgtctatagctgaaagG	tcttctcgtctatagctgaaagT	ccctttgctgcatgtgaga	chromosome_inspecific	forward
R34	non-homeologous	tctgagatgatgatactTtgtggG	tctgagatgatgatactTtgtggA	actgggatgacctctgtat	chromosome_inspecific	forward
R35	non-homeologous	tgaagaagtggaattctctgtgtG	tgaagaagtggaattctctgtgtC	ctttTagtctcttaattctattgcttC	chromosome_inspecific	forward
R36	non-homeologous	tgaaatgctctgtcaatgccA	tgaaatgctctgtcaatgccG	ATGCGAATTGGGGAATTAAA	chromosome_inspecific	reverse
R37	non-homeologous	tgcataatgctgaagagactcG	tgcataatgctgaagagactcA	tgtccacctactcaagtctgc	chromosome_inspecific	reverse
R38	non-homeologous	tgcCcaagTtTtctgcaagaT	tgcCcaagTtTtctgcaagaG	tgtaggagaagactCgaagtA	chromosome_specific	forward
R40	non-homeologous	tgcataatgctgaagagactcA	tgcataatgctgaagagactcG	agtcctgaagcatgtgct	chromosome_nonspecific	reverse
R43	non-homeologous	tcgctgatttcatcatgtcccA	tcgctgatttcatcatgtcccG	tcaggtgctgcaaatgttagG	chromosome_semispecific	forward

it is the donor species of *Yr15* (McIntosh et al., 1995). This test panel allows to test if the SNPs are only diagnostic to *T. diccocooides* instead of *Yr15*. On the test panel, 28 (80%) SNPs were polymorphic across the parents and three of them were diagnostic to *yr15* (R5, R8, R33). From the five homoeologous SNPs, three of them were monomorphic and two polymorphic, suggesting that PolyMarker is effective on detecting which assays are less likely to work (Table 3.10; Figure 3.13a,b). The segregation of the SNPs in the full F_2 population (Section 3.1, Figure 3.13c) and a genetic map was produced (Section 3.8).

3.8 Genetic map

Initially, the 28 polymorphic markers were used to genotype a subset of 66 plants from the F_2 population. From those, 23 (82%) were linked to *Yr15* and several markers fall in a small interval around *Yr15* (Figure 3.14a; Table 3.10), confirming that the multiple-criteria strategy (Section 3.6) for selecting candidate SNPs was effective. Then, the complete F_2 population was assessed with:

- the seven markers that were most linked to *Yr15*, including two of the diagnostic markers from the variety panel (R5 and R8),
- The flanking SSR microsatellite markers used by UK breeders for germoplasm selection (Xbarc8 and Xgwm413).
- A marker based on barley-wheat synteny (R43) which met the selection criteria, but wasn't on the original set of 50 markers with high BFR.

The F_2 population consisted on 232 plants with phenotypic information, of those 196 were genotyped reliably (no more than one data point missing). Using the eight SNP markers and 2 SSRs, the *Yr15* locus was mapped to an interval of 0.77cM, with R8/xgwm413 0.26cM distal, and R5/R11 0.77cM proximal from *Yr15* (Figure 3.14b,c).

The sub-cM resolution is expected on an F_2 population of 196 individuals, as 392 gametes provide a resolution of 0.26cM. Despite the fact that none of the selected markers have perfect linkage to *Yr15*, the produce genetic map is an improvement in the resolution of the map for the locus and it enables the shift to SNP markers from microsatellites, which

Table 3.10: Results of validation of primers on the progenitors (AVS and *Yr15*, varieties known to contain *Yr15* (Cortez, Ochre and, Boston) and, varieties without *Yr15* (Robigus, Cadenza and, Shamrock). Shamrock and Robigus have *T. dicoccoides* introgressions. The bold markers are diagnostic in the panel (R5, R8, R88) or in the genetic map (R11).

Assay ID	Gene set	Gene model name	SNP	Yr15+					Yr15-			Polymorphic	Linked Yr15	comment
				Yr15	Ochre	Boston	Cortez	Shamrock	Robigus	Cadenza	AVS			
R1	UCW	UCW_Tt-k55_contig.8830;tt-k21_contig.10204	C341G	A	H	A	A	A	-	A	B	Yes	*	segregation distortion
R2	UniGene v60	gnl UG Ta#S13126619	C491T	B	B	B	B	B	B	B	B	No	-	
R3	UCW	contig95240	C220G	H	B	B	B	B	B	B	B	Yes	Yes	
R4	UCW	contig105384	C1227T	A	B	B	B	B	B	B	B	Yes	Yes	
R5	UniGene v60	gnl UG Ta#S58861868	A214G	A	A	A	A	B	B	B	B	Yes	Yes	
R6	UCW	KukriC706.1	T2979C	A	H	A	A	B	B	H	H	Yes	No	
R7	UniGene v60	gnl UG Ta#S37932863	C281T	H	A	A	A	B	B	A	B	Yes	No	
R8	UniGene v60	gnl UG Ta#S58863387	T241C	B	B	B	B	A	A	A	A	Yes	Yes	
R9	UniGene v60	gnl UG Ta#S58892239	C303T	H	B	A	B	B	B	H	B	Yes	No	
R10	UCW	UCW_Tt-k63_contig.79829	C207T	H	A	B	A	B	B	B	B	Yes	Yes	
R11	UCW	UCW_Tt-k45_contig.39011	C726T	A	A	A	H	-	B	B	B	Yes	Yes	
R12	UCW	contig50308	G587A	-	H	H	H	B	B	A	B	Yes	Yes	
R14	UniGene v60	gnl UG Ta#S44692929	C549T	A	A	-	A	A	B	-	B	Yes	Yes	
R15	UCW	UCW_Tt-k51_contig.2344;tt-k55_contig.2091	T686G	A	A	A	A	A	A	A	A	No	-	
R16	UniGene v60	gnl UG Ta#S17898149	G227A	A	A	B	A	B	B	B	B	Yes	Yes	
R17	UCW	CL3339Contig1	T509C	H	H	H	H	H	H	H	H	No	-	
R19	UCW	UCW_Tt-k21_contig.8407;tt-k61_contig.5972	C1405T	A	B	B	B	B	B	B	B	Yes	Yes	
R20	UCW	UCW_Tt-k21_contig.8407;tt-k61_contig.5972	T1102C	A	B	B	B	-	B	B	B	Yes	Yes	
R21	UCW	UCW_Tt-k31_contig.53804;tt-k41_contig.31582	G1810T	H	B	B	B	B	B	B	B	Yes	Yes	
R22	UCW	UCW_Tt-k31_contig.14966	T408C	A	A	A	A	A	A	A	A	Yes	Yes	
R23	UCW	UCW_Tt-k51_contig.12731;tt-k55_contig.13077;tt-k61_contig.18734	C50T	A	H	H	H	H	-	H	B	Yes	Yes	
R24	UCW	UCW_Tt-k55_contig.8830;tt-k21_contig.10204	T3005G	H	H	B	H	B	B	B	B	Yes	Yes	
R25	UCW	UCW_Tt-k63_contig.79829	G184A	A	A	A	A	H	H	A	A	No	-	
R26	UCW	UCW_Tt-k21_contig.3794	C702T	H	A	A	A	B	B	B	B	Yes	Yes	
R28	UCW	KukriC3701.1	T1053C	A	A	B	A	B	B	B	B	Yes	Yes	
R29	UCW	UCW_Tt-k55_contig.8640;tt-k41_contig.8875	G783A	H	A	B	A	B	B	B	B	Yes	Yes	
R30	UCW	UCW_Tt-k55_contig.8830;tt-k21_contig.10204	T2184A	A	A	B	A	B	B	A	B	Yes	Yes	
R31	UCW	UCW_Tt-k45_contig.22098	G683T	A	B	A	B	B	B	A	B	Yes	Yes	
R32	UCW	UCW_Tt-k21_contig.33188;tt-k25_contig.30647	C596A	H	A	A	A	B	B	A	B	Yes	Yes	
R33	UniGene v60	gnl UG Ta#S58861868	G486T	A	A	A	A	B	B	B	B	No	-	
R34	UCW	UCW_Tt-k31_contig.34099	G1713A	H	A	B	-	B	B	B	B	Yes	No	
R35	UniGene v60	gnl UG Ta#S58900202	T889C	A	B	B	B	B	B	B	B	Yes	Yes	
R36	UCW	UCW_Tt-k55_contig.8830;tt-k21_contig.10204	T2349C	H	H	H	H	H	-	H	B	Yes	Yes	
R37	UCW	UCW_Tt-k31_contig.34099	C846T	B	B	B	B	B	B	B	B	No	-	
R38	UniGene v60	gnl UG Ta#S58840501	T179G	B	B	B	B	B	B	B	B	No	-	
R40	UCW	UCW_Tt-k31_contig.34099	C846T	A	H	B	A	-	B	B	B	Yes	No	based on barley synteny
R43	UniGene v60	gnl UG Ta#S58843705	G268A	A	B	B	-	-	B	-	B	Yes	Yes	based on barley synteny




Figure 3.14: Genetic maps for *Yr15*. (a) Genetic map of the test panel from 50 individuals. (b) Genetic map from 196 individuals from the full population only with the 8 markers previously identified as closer to the *Yr15* locus. (c) Graphical genotype of the 196 F_2 individuals used to develop the genetic map. The alleles are abbreviated according to their origin: A: AVS; B: *Yr15* and H: Heterozygous. Missing calls are indicated by a hyphen.

has become the preferred marker system in MAS pipelines in breeding programmes.

3.9 Bioinformatic methods

3.9.1 Alignment reads to gene models

The raw output from the Illumina HiSeq 2000 was processed with Casava v1.8 (Illumina, 2011). Lanes 1 and 2, containing multiplexed bulks (Table 3.1) was demultiplexed with a tolerance of 1 mismatch in the barcode. Lanes 3 and 4 contained the parental sequences without a barcode. The FastQ files were left compressed and in chunks of 40,000, as the default for the BCL conversion pipeline from Casava to allow parallel processing in a cluster environment. The quality of the sequencing lanes was assessed with FastQC v0.10.1 (Babraham Bioinformatics, 2012). The RNA-Seq reads were aligned with BWA 0.5.9 (Li and Durbin, 2009) to the wheat UniGene database v60 (Pontius et al., 2002) and to the UCW gene models (Krasileva et al., 2013), including the *T. turgidum* and complementary ORFs (MAS Wheat, 2013). The alignments were sorted and stored as single BAM files to have random access (Li et al., 2009).

3.9.2 Bulk Frequency Ratios

Listing 3.1: Method to find best BLAT alignment

```
def self.each_best_hit(text = '')
  emptyHit = Bio::Blat::Report::Hit.new
  emptyHit.score = 0
  best_aln = Hash.new(emptyHit)
  self.each_hit(text) do |hit|
    current_score = hit.score
    if current_score > best_aln[current_name].score
      best_aln[current_name] = hit
    end
  end
  best_aln.each_value { |val| yield val }
end
```

Listing 3.2: Extensions to Bio::Blat::Report::Hit to get the percentage of coverage

```
class Bio::Blat::Report::Hit
  def covered
    match + mismatch
  end
  def query_percentage_covered
    covered * 100.0 / query_len.to_f
  end
  def target_percentage_covered
    covered * 100.0 / target_len.to_f
  end
end
```

3.9.3 Alignment between gene models

3.10 Discussion

Resequencing the ~ 17 Gbp genome of hexaploid wheat is costly and approaches to reduce the required sequenced volume to effectively call for SNPs had been evolving since the conception of this project. The RNA and DNA extraction and the sequencing for this project was carried on before the beginning of my PhD (before October 2012). At that point exome capture was already established for genotyping humans (Ng et al., 2009), however the first exome capture on wheat was just recently published, with probes coming from unassembled 454 reads (Winfield et al., 2012), and a probe designed from transcripts (Henry et al., 2014) was not published after the analysis of this section was completed and validated. An even more targeted capture for resistance genes (RenSeq) was published while this study was executed (Jupe et al., 2013). On the other hand RNA-Seq was already tested for Bulk Segregant Analysis on tetraploid wheat (Trick et al., 2012). Hence, the decision of reducing the sequenced space with RNA-Seq was appropriate at the time (Figure 1.1). Unfortunately, one of the shortcomings of RNA-Seq used to call for SNPs is that the coverage is not uniform and the genes that have low expression don't have enough coverage to call for SNPs (Section 3.2). If a similar study is to be started today, a better alternative would be to use exome capture in general from a segregating population for any trait,

or RenSeq if the target gene is a resistance gene.

The quality and completeness of the reference genome or gene models directly affects the mapping NGS reads. This is particularly true on polyploid organisms: if one of the homoeologues is absent, the reads are likely to map to the wrong genome if the parameters of the aligner are relaxed or; not map at all if the the required identity is high. When the bioinformatic analysis of this project started, the only available wheat genomic reference was a whole genome shotgun 454 sequencing, unassembled (Brenchley et al., 2012); the Chinese Spring Chromosome arm survey sequence (CSS) assembly was being finished (Mayer et al., 2014); the longer scaffolds from Chapman et al. (2015) were not public yet and; the efforts to make a whole genome shotgun assembly were being planned independently by the International Wheat Genome Sequencing consortium (Pozniak, 2016) and TGAC (Clark, 2016). Because a contiguous assembly with the corresponding annotation wasn't available at the time of the analysis and the fact that the data available was from a transcriptome, the use of gene models as a reference for the alignment was a suitable approach.

In terms of available gene sets when the analysis started, the canonical reference was the UniGenes from the NCBI (Pontius et al., 2002). The UniGenes are produced with an automated pipeline that clusters all the ESTs deposited in the NCBI by identity and selects the longest transcript, which can merge homoeologous transcripts as a single reference. Shortly after I started the bioinformatic analysis, two additional gene models were available, the draft annotation for the CSS assembly (MIPSV1) in January 2013 and the UCW gene models (Krasileva et al., 2013) in May 2013. I selected the UCW gene models, as they were more mature and were phased to distinguish between genomes and already published, over the the MIPSV1 genes, still being refined from an initial approach lifting proteins from related organisms and a few RNA-Seq experiments. The MIPS gene models were improved by removing duplications in the assembly in a later stage and the nomenclature before the release of the assembly (Mayer et al., 2014), but at that point the results of this project where already submitted for publication (Figure 1.1; Ramirez-Gonzalez et al. 2015c).

To locate the gene models in the chromosome arms and see if there was an enrichment on the called SNPs the use of a high resolution consensus

Should I talk briefly about barley?. I would need to add a section before

map is needed, as the genome assemblies available during the analysis are fragmented. Timely, a genetic map with $> 42,000$ markers was published (Wang et al., 2014). I was able to use it to locate several CSS scaffolds before the assembly was published, as I collaborated in the project. The located scaffolds were used as proxy to sort just under half of the reference genes in their chromosomal position (Section 3.5). Despite the resolution not being enough to find a single point of enrichment, it was enough to confirm that the SNPs were in the expected location, including one of the SNP candidates flanking the *Yr15* locus (SNP R11, Figure 3.9b). If the analysis was to be done today, the genetic map from Chapman et al. (2015) along with their longer scaffolds, or the scaffolds from TGACv1 or the NRGene should provide a better resolution. Even without having all the CSS scaffolds sorted, the fact that they come from individual chromosome arms they enabled the assignment of the genes to a chromosome.

The original expectation was to have a NIL for the BSA, however the number of SNPs called in the progenitors suggested that the background, Avocet S, was not the same. This happened because despite both susceptible lines being called the same and having the same response to the pathogen, they are different lines from different countries (Section 3.3). This highlights the importance of genotyping the material used when developing mapping populations, specially if the source of the seeds come from different seed banks.

Despite this shortcomings, the use of the BFRs to score the putative SNPs was effective as most of the SNPs with a high score mapped in chromosome 1B, as expected from previous studies ($BFR > 6$, Section 3.6). Using the extra criteria of only selecting SNPs from the resistant progenitor and in the expected chromosome arm I was able to produce a high resolution genetic map (Section 3.8). The genetic map was of the expected resolution for the size of the population (0.26cM on 196 individuals). Since the mapping population contained only one critical recombinant between *Yr15* and the flanking markers, the population couldn't yield to a better map. To improve the map, a cross from the two critical recombinants could be use to repet a similar analysis, but sequencing with either exome capture or RenSeq.

As described in Ramirez-Gonzalez et al. (2015c), the markers R11, R5 and R8 were tested "across 122 doubled haploid (DH) lines. These

Talk about Why is R33 diagnostic on the varieties, but maps away?.

SNP haplotype			Reaction to <i>P. striiformis</i>		
R11	R5	R8	Resistant	Intermediate	Susceptible
C	A	T	-	6	16
T	A	T	-	11	-
T	G	C	79	1	-

Figure 3.15: Haplotype analysis and phenotypic evaluation of the 113 doubled haploid lines used in the study. The TGC haplotype corresponds to that originally identified in the *Yr15* parent and which was diagnostic across 112 of the 113 lines studied.

DH lines were derived from crosses between five different UK varieties/breeding lines to *Yr15* derivatives known to carry the resistance gene. The expected *Yr15* haplotype corresponded to T, G and C alleles at markers R11, R5 and R8, respectively (TGC haplotype). The DH lines were tested at seedling stage for reaction to *P. striiformis*, with 84 showing complete resistance and 34 presenting an intermediate or completely susceptible reaction. The resistant lines all carried the complete *Yr15* haplotype (TGC, Figure 3.15) across the three SNP markers with the exception of five lines which had a single missing data point, but were otherwise consistent. This compared favourably with the most diagnostic in-house SNP markers available within the breeding programmes. Using the three in-house markers, 79 resistant lines carried the expected haplotype, but five completely resistant DH lines were scored as false negative due to the presence of the non-*Yr15* haplotype. Within the intermediate and susceptible DH lines, all but one had a non-*Yr15* haplotype (CAT or TAT) across R11, R5 and R8 (Figure 3.15). This single DH line was scored as a false positive as it carried the TGC *Yr15* haplotype, but was found to have an intermediate (chlorotic) reaction to *P. striiformis*. This line was also the only one scored as a false positive using the three in-house markers". The fact that the developed markers perform better than the markers developed by breeders show the value of this particular experiment and further confirms that BSA combined with NGS is an effective way to develop novel markers.

This is a very long quote, but I'm finding hard to shorten it.

Mention other people using a similar strategy since this was published.

Chapter 4

Gene expression (expVIP)

4.1 Expression experiments (Introduction)

Describe the list of previously published expression experiments and how they can potentially be used as a framework for new experiments.

4.2 Database design

Description of how the database was designed and the flexibility given by having the factors and units as variables

4.3 Analysis pipeline

Implementation of the pipeline, from running kallisto to load the data in the database

4.4 Graphical interface

How the expression can be displayed filtered, and sorted

4.5 Conclusions

The use of previously published studies is a valuable resource. Also, mention that despite the fact that there are several expression/gene browsers, none of them allow comparisons between species and don't consider polyploids.

Chapter 5

Conclusions and final remarks

This section wraps up by showing the relationship and importance of a comprehensive approach to data analysis, from the field, genetics, molecular biology and genomics. I will also remark how the technology and the resources have changed in the last 4 years. As at the references used at beginning where superseded during the PhD.

Appendix A

Supplemental tables

Table A.1: Count of KASP assays designed for the 40,267 SNP markers located in the genetic map from Wang et al. (2014). 4,228 assays did not align to the target chromosome. Not designed: Primer3 could not find viable primers flanking the SNP.

	Homoeologous variant	Varietal SNP	Percentage
Non-specific	1,765	5,857	21.15%
Semi-specific	7,942	6,907	41.20%
Specific	6,813	5,957	35.43%
Not designed	242	556	2.21%
Total	16,762	19,277	36,039

A.1 PolyMarker supplemental tables.

Table A.2: PolyMarker used to genotype PST

Assay	Contig	Position	X	Y	Cluster I isolates		Cluster II isolates		Cluster III isolates		Cluster IV isolates	
					13/26	13/123	CL1	T-13/3	13/09	13/23	13/182	13/36
1	PST130.14470	268	C	T	X:Y	X:Y	X:X	X:X	X:X	X:X	X:X	X:X
2	PST130.8160	11876	C	T	Y:Y	Y:Y	X:Y	X:Y	X:Y	X:Y	X:Y	X:Y
3	PST130.14628	1712	A	C	X:Y	-	X:X	X:X	X:X	X:X	X:X	X:X
4	PST130.14898	503	G	A	X:X	X:X	X:Y	X:Y	X:Y	-	X:Y	X:Y
5	PST130.28344	2372	A	G	Y:Y	Y:Y	X:Y	X:Y	Y:Y	Y:Y	Y:Y	Y:Y
6	PST130.7634	3463	A	C	Y:Y	Y:Y	X:Y	X:Y	Y:Y	Y:Y	Y:Y	Y:Y
7	PST130.7629	11699	G	A	Y:Y	Y:Y	X:Y	X:Y	Y:Y	Y:Y	Y:Y	Y:Y
8	PST130.10943	2979	C	T	X:Y	X:Y	X:X	X:X	X:X	X:X	X:Y	X:Y
9	PST130.10126	6216	G	T	Y:Y	Y:Y	X:X	X:X	X:X	-	Y:Y	Y:Y
10	PST130.22010	172	C	T	Y:Y	Y:Y	Y:Y	Y:Y	X:Y	-	X:Y	X:Y
11	PST130.16961	1098	C	T	X:X	X:X	X:Y	X:Y	Y:Y	Y:Y	X:Y	X:Y
12	PST130.6915	2710	A	T	Y:Y	Y:Y	Y:Y	Y:Y	Y:Y	X:Y	Y:Y	Y:Y
13	PST130.12479	1428	C	T	X:X	X:X	Y:Y	Y:Y	X:X	X:X	Y:Y	X:X
14	PST130.7634	3883	C	G	X:X	X:X	X:Y	X:Y	X:X	X:X	X:Y	X:X
15	PST130.14470	456	T	C	Y:Y	Y:Y	X:Y	X:Y	Y:Y	Y:Y	Y:Y	Y:Y

Table A.3: Validation of homozygous deletions on line Cadenza0423.

Marker	Deletion	chr	cM	1	2	3	4	5	6	7	8	9	10	11	12	C	C	C	C	Result
5BS_2297308_Cadenza0423_12664_C12664T	-	5B	4.551	X	X	-	X	X	X	X	X	X	X	-	X	Y	Y	Y	Y	HOM Mutation
5BL_10812849_Cadenza0423_5664_G5664T	-	5B	38.769	X	X	-	X	X	X	X	X	X	X	-	X	Y	Y	Y	Y	HOM Mutation
5BL_10825062_Cadenza0423_7917_G7917A	-	5B	38.769	X	X	-	X	X	X	X	X	X	X	-	X	Y	Y	Y	Y	HOM Mutation
IWGSC_CSS_5BL_scaff_10847976:27068-27231	+	5B	38.769	X	X	-	X	X	X	X	X	X	X	-	X	H	H	H	H	Hom Deletion
IWGSC_CSS_5BL_scaff_10847976:28118-28674	+	5B	38.769	X	X	-	X	X	X	X	X	X	X	-	X	H	H	H	H	Hom Deletion
IWGSC_CSS_5BL_scaff_10865441:15863-15946	+	5B	38.769	X	X	-	X	X	X	X	X	X	X	-	X	H	H	H	H	Hom Deletion
5BL_10837222_Cadenza0423_4616_G4616A	-	5B	39.905	X	X	-	X	X	X	X	X	X	X	-	X	Y	Y	Y	Y	HOM Mutation
5BL_10891320_Cadenza0423_18847_C18847T	-	5B	45.594	Y	Y	-	Y	H	X	X	Y	H	Y	-	H	Y	Y	Y	Y	HET Mutation

Table A.4: Validation of mutations on M_4 on Cadenza

IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Cadenza)	Primer 2 (mutant)	Common Primer
IWGSC_CSS_3B_scaff_10445294	Cadenza1772	6019	C	T	het	het	caggatAgtGggactgtcaaaG	caggatAgtGggactgtcaaaA	ggagacGGctGtggacatT
IWGSC_CSS_3DL_scaff_6955403	Cadenza1772	2418	C	T	het*	hom	tcagCggattgtcgggatG	tcagCggattgtcgggatA	tgteCatgaaTcttgtccacG
IWGSC_CSS_4AL_scaff_7106846	Cadenza1772	11277	G	A	hom	hom	tgggatccatgcctacactG	tgggatccatgcctacactA	gatggTgatttgcgcctA
IWGSC_CSS_4AS_scaff_5991335	Cadenza1772	15710	G	A	hom	hom	ctggccctgcgctgctaC	ctggccctgcgctgctaT	gtggaaGttcagaaggaccaG
IWGSC_CSS_4BS_scaff_4956646	Cadenza1772	252	G	A	het*	hom	gcaggttgacttcccgaG	gcaggttgacttcccgaA	tGaggtacgaGcTaaagAaagC
IWGSC_CSS_4DS_scaff_1715962	Cadenza1772	1225	G	A	hom	hom	cagctgtggTatctcaactgG	cagctgtggTatctcaactgA	CcCtGaaACACcGtttggAT
IWGSC_CSS_5AL_scaff_2763407	Cadenza1772	2119	G	A	hom	hom	gcgacGaacctcgagatctG	gcgacGaacctcgagatctA	gaTggcaAtcgtCgtgcA
IWGSC_CSS_5AS_scaff_1548786	Cadenza1772	12625	C	T	het	het	AtaggcacattgctagactgaG	AtaggcacattgctagactgaA	ggattgggtgttcacgC
IWGSC_CSS_5BL_scaff_10849226	Cadenza1772	2289	C	T	het*	hom	cctgacatcattgttcacgatC	cctgacatcattgttcacgatT	cactccgaggtgtccatgaT
IWGSC_CSS_5BS_scaff_2270737	Cadenza1772	2262	G	A	hom	—	attcCTgtgttggtggCaaatgaG	attcCTgtgttggtggCaaatgaA	taaGcaciaAccctccagctgG
IWGSC_CSS_1AL_scaff_3022915	Cadenza1661	891	C	T	hom	hom	ccacagtgcgactcctattgaCG	ccacagtgcgactcctattgaCA	atgtctgattcGtcGtagtcC
IWGSC_CSS_1AS_scaff_3297240	Cadenza1661	1970	C	T	het	het	catcccgccGtttctcC	catcccgccGtttctcT	gtctcgccgatgaagagcT
IWGSC_CSS_1BL_scaff_3828996	Cadenza1661	1340	G	A	hom	hom	agccggatgttagtgttaacC	agccggatgttagtgttaacT	agcagcttgTcgcgttaaC
IWGSC_CSS_1DS_scaff_1884529	Cadenza1661	10575	G	A	hom	hom	aCagatacaAttgtcatgcaggC	aCagatacaAttgtcatgcaggT	acctgggTTgtccaatacttC
IWGSC_CSS_2AL_scaff_6318370	Cadenza1661	19142	C	T	het	—	cgtggcCgaatCtcGacG	cgtggcCgaatCtcGacA	ttcttggggagccgggC
IWGSC_CSS_2AS_scaff_5213460	Cadenza1661	1358	G	A	hom	hom	gtcacgaaCccgctcagA	gtcacgaaCccgctcagA	aggaaagagagaaaagaGcG
IWGSC_CSS_2BS_scaff_5179331	Cadenza1661	5604	G	A	het	het	actctcgtcaagaactgatacaG	actctcgtcaagaactgatacaA	gcaGagaatgttcttgaacT
IWGSC_CSS_2DS_scaff_5341235	Cadenza1661	4673	G	A	het	het	ggtgaggatctcggagctG	ggtgaggatctcggagctA	gcgcggtcgtacaggttG
IWGSC_CSS_3AL_scaff_4250995	Cadenza1661	7046	G	A	hom	hom	cCaagaaacgggtggccaG	cCaagaaacgggtggccaA	ctgcagctgtccatcatcgT
IWGSC_CSS_3B_scaff_10404421	Cadenza1661	4303	G	A	het	het	ccttcgtcgaCaggacctG	ccttcgtcgaCaggacctA	GCcagtaactCacAtgtctC
IWGSC_CSS_5DL_scaff_2390496	Cadenza1538	2125	C	T	hom	het	gcagttttatcctcagtagtcttgG	gcagttttatcctcagtagtcttgA	ttctgagaaTgtaagtgtcGatG
IWGSC_CSS_6AL_scaff_5753680	Cadenza1538	3920	C	T	hom	hom	tgctccaaatttgagcaciaTaaC	tgctccaaatttgagcaciaTaaT	aaatgcaaggggttaagtttttG
IWGSC_CSS_6AS_scaff_4425792	Cadenza1538	4307	G	A	hom	het	agatgcttgtCggGccaG	agatgcttgtCggGccaA	gctgaagcaacgcgatcaaT
IWGSC_CSS_6BS_scaff_3003630	Cadenza1538	6933	C	T	het	het	ggcagtaagtgtgtgctgagC	ggcagtaagtgtgtgctgagT	tTgaCttctggtttgtggcA
IWGSC_CSS_6DL_scaff_3246988	Cadenza1538	9186	G	A	het	het	gctaaagaagagcttgagagaattC	gctaaagaagagcttgagagaattT	aatttctgaagagaggtgtgtatG
IWGSC_CSS_7AL_scaff_4480114	Cadenza1538	3446	C	T	het	—	gatatctcccacacggcgG	gatatctcccacacggcgA	tgagccactcttcagtttT
IWGSC_CSS_7AS_scaff_4193541	Cadenza1538	8359	C	T	hom	het	agcaattctttggctatcaattagC	agcaattctttggctatcaattagT	tcactGtcttaactctactgctG
IWGSC_CSS_7BL_scaff_6721572	Cadenza1538	9223	C	T	het	het	gctCaggaggagagacaagaaG	gctCaggaggagagacaagaaA	tgctatgaagaattccgacctC
IWGSC_CSS_7BS_scaff_3152545	Cadenza1538	3960	G	A	hom	—	tcagcaaaatcacctgcCgC	tcagcaaaatcacctgcCgT	gCtgcccatcatcgtttaT
IWGSC_CSS_7DS_scaff_3963838	Cadenza1538	2913	G	A	het	het	tCgttgcaagcCttTtgtgT	tCgttgcaagcCttTtgtgT	agaGttaTcaagcTactgtcacA
IWGSC_CSS_1AL_scaff_3903380	Cadenza1469	6193	G	A	hom	hom	ctcttcAgagatgaacgcgG	ctcttcAgagatgaacgcgA	tcGtGagatgGtggtttGTTA
IWGSC_CSS_1AS_scaff_3287728	Cadenza1469	3817	C	T	het*	hom	ccgaccaAttcactaacggG	ccgaccaAttcactaacggA	acctctttccccAgacatgaT
IWGSC_CSS_1BL_scaff_3815304	Cadenza1469	513	G	A	hom	hom	aacatttgctTaCcaaaacGC	aacatttgctTaCcaaaacGT	acacagcaagttataatgCAAAGC
IWGSC_CSS_1DL_scaff_2266648	Cadenza1469	5926	C	T	het	het	caacatgagacacacaccttC	caacatgagacacacaccttT	gtcaacgcgtgaggattgtC
IWGSC_CSS_1DS_scaff_1906671	Cadenza1469	3697	C	T	hom	hom	tggTGTgtagacacttggcgaG	tggTGTgtagacacttggcgaA	catggcgaacaccAcctG
IWGSC_CSS_2AL_scaff_6337088	Cadenza1469	7334	G	A	het*	hom	acaatgccAagttgacaggttG	acaatgccAagttgacaggttA	gggagtggtgttCagaacaT

IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Cadenza)	Primer 2 (mutant)	Common Primer
IWGSC_CSS_2BL_scaff.7972799	Cadenza1469	8995	C	T	het	hom	gTgCtcctcGgcaccccttC	gTgCtcctcGgcaccccttT	gatccgGgcaaacacgTG
IWGSC_CSS_2DL_scaff.9832343	Cadenza1469	3262	G	A	het	het	TtgtctaAcagcacCGcagG	TtgtctaAcagcacCGcagA	agatctcggtcagcctttcT
IWGSC_CSS_2DS_scaff.5327939	Cadenza1469	3889	G	A	het	het	ttttTgccttatgtgactctagtaC	ttttTgccttatgtgactctagtaT	gaggccatcacagatagcG
IWGSC_CSS_3B_scaff.10395219	Cadenza1469	1292	G	A	hom	—	agggtccttgctgcttgctgG	agggtccttgctgcttgctgA	cctctctgggggctttataC
IWGSC_CSS_3B_scaff.10592217	Cadenza0580	2994	C	T	het	—	acagcagtatcaagccctcC	acagcagtatcaagccctcT	tgatactgttgTggCggagG
IWGSC_CSS_3DS_scaff.2596771	Cadenza0580	1037	G	A	het	het	tggttatgCAcaggataatCagG	tggttatgCAcaggataatCagA	tggcaaatgtgatgtcattaggT
IWGSC_CSS_4AL_scaff.7093953	Cadenza0580	9881	C	T	hom	hom	GacaggaagccggtaacaC	GacaggaagccggtaacaT	ctccAgcaggcatgggaT
IWGSC_CSS_4BL_scaff.7037448	Cadenza0580	1837	C	T	hom	hom	CgttgaaaaGctgcaagaacttaaC	CgttgaaaaGctgcaagaacttaaT	cagttcttccTtCaGagcagataT
IWGSC_CSS_4BS_scaff.4929479	Cadenza0580	10668	G	A	hom	—	tggattttcccgactgttC	tggattttcccgactgttT	gtaaacaggcatttcaagagtcA
IWGSC_CSS_4DL_scaff.14359838	Cadenza0580	1408	G	A	hom	—	gCtcAttcagggatTGTcCtaTatG	gCtcAttcagggatTGTcCtaTatA	tgaCagaacagttggtcatatcT
IWGSC_CSS_4DS_scaff.2276484	Cadenza0580	8034	G	A	hom	hom	gccgtggttgatggAgaG	gccgtggttgatggAgaA	cgctcagattactgatacttgcA
IWGSC_CSS_5AL_scaff.2756579	Cadenza0580	5278	G	A	het	het	tgaatggatttttctgccgttC	tgaatggatttttctgccgttT	ggAAatCCTATgCagaAgAaaCTG
IWGSC_CSS_5BL_scaff.10787208	Cadenza0580	10627	G	A	het	—	gcctctcacatcgcgagaC	gcctctcacatcgcgagaT	acgatgtcAggtggGcgT
IWGSC_CSS_5BS_scaff.2282179	Cadenza0580	5267	G	A	het	—	tgatgggctacgacgtgC	tgatgggctacgacgtgT	tcggcgcccttgaaAtcC
IWGSC_CSS_5DL_scaff.4498073	Cadenza0423	4937	C	T	hom	hom	gcaccctctggttggtcatC	gcaccctctggttggtcatT	tgacagcaAagcagccG
IWGSC_CSS_5DS_scaff.2738970	Cadenza0423	2319	C	T	het	—	cgtgaggtgggtgatttG	cgtgaggtgggtgatttT	tggaaactagtacactgcagtTC
IWGSC_CSS_6AL_scaff.5757109	Cadenza0423	2788	G	A	hom	hom	caggaGcctggcaataaaGG	caggaGcctggcaataaaGA	ctttcGagctctcttagtttcG
IWGSC_CSS_6AS_scaff.4387871	Cadenza0423	2543	G	A	hom	hom	gcatgctaacaggcgaaaagG	gcatgctaacaggcgaaaagA	ctcatgctctgatcttaaggtT
IWGSC_CSS_6BL_scaff.4271391	Cadenza0423	4660	C	T	hom	hom	tacgtgcatgatgtggtagtcgtaC	tacgtgcatgatgtggtagtcgtaT	gtttgaaagtgcacagatgTaccA
IWGSC_CSS_6DS_scaff.1880206	Cadenza0423	9159	G	A	het	het	ctgCgaaggctccacaaG	ctgCgaaggctccacaaA	ggatgagaagtttgcattgctC
IWGSC_CSS_7AS_scaff.4227506	Cadenza0423	952	G	A	het	—	ccatgtgtttccaatgttagagC	ccatgtgtttccaatgttagagT	tgccctagctggtatgcT
IWGSC_CSS_7BL_scaff.6681782	Cadenza0423	1486	C	T	hom	hom	agtaagCGtgacagcaatggG	agtaagCGtgacagcaatggA	AtgtctTtgGtggaagtacatcA
IWGSC_CSS_7BS_scaff.3160328	Cadenza0423	7801	C	T	het	het	tgttaaatGatacagCctgcagC	tgttaaatGatacagCctgcagT	tggaaatgggtCgttgtttT
IWGSC_CSS_7DS_scaff.407428	Cadenza0423	2051	G	A	het	het	gtcGCgccatcctgacaG	gtcGCgccatcctgacaA	actcatcAggtcagcccaA
IWGSC_CSS_3AL_scaff.442479	Cadenza0364	3198	C	T	het	het	gagtcATaagttggtaagattggC	gagtcATaagttggtaagattggT	GCaGaTaaCaacaggatcacG
IWGSC_CSS_3AL_scaff.4447942	Cadenza0364	11917	G	A	het	het	gtcataaaagattgctcctgtgaaG	gtcataaaagattgctcctgtgaaA	ctcGgatgtgggaggaagA
IWGSC_CSS_3AS_scaff.1557483	Cadenza0364	2547	C	T	het	het	aaagtcacatcatgcttaccataaG	aaagtcacatcatgcttaccataaA	cgaataccaacgcctcatcA
IWGSC_CSS_3AS_scaff.2648747	Cadenza0364	2688	G	A	het	het	tggAagcAcaaggggccC	tggAagcAcaaggggccT	GccgccgatggagactcG
IWGSC_CSS_3AS_scaff.3304956	Cadenza0364	1017	G	A	het	het	gtcccttgacacagctttG	gtcccttgacacagctttA	cctgctggactacaactcaaT
IWGSC_CSS_3AS_scaff.3321091	Cadenza0364	4585	C	T	het	het	caagaatgATgctgatgttggaG	caagaatgATgctgatgttggaA	acatgctgaatgccgaatC
IWGSC_CSS_3AS_scaff.3371333	Cadenza0364	538	G	A	het	het	gggaaaCgAgAcgagcgG	gggaaaCgAgAcgagcgA	ccgtgcttctctaccctT
IWGSC_CSS_3AS_scaff.3371815	Cadenza0364	1061	C	T	het	het	atccccacggcacagagG	atccccacggcacagagA	aAttggcccttggtgattcC
IWGSC_CSS_3AS_scaff.3440912	Cadenza0364	4498	G	A	het	het	ccgtaaaaactttctgtgcttgC	ccgtaaaaactttctgtgcttgT	atActgacaaactacatgatgtgC
IWGSC_CSS_3B_scaff.10343586	Cadenza0364	2242	G	A	het	—	gggttcTgTcctctcttccactG	gggttcTgTcctctcttccactA	tgtgttgaaccgcgaagcA
IWGSC_CSS_3AL_scaff.442479	Cadenza0364	3198	C	T	het	het	gagtcATaagttggtaagattggC	gagtcATaagttggtaagattggT	GCaGaTaaCaacaggatcacG
IWGSC_CSS_3AL_scaff.4447942	Cadenza0364	11917	G	A	het	het	gtcataaaagattgctcctgtgaaG	gtcataaaagattgctcctgtgaaA	ctcGgatgtgggaggaagA
IWGSC_CSS_3AS_scaff.1557483	Cadenza0364	2547	C	T	het	het	aaagtcacatcatgcttaccataaG	aaagtcacatcatgcttaccataaA	cgaataccaacgcctcatcA
IWGSC_CSS_3AS_scaff.2648747	Cadenza0364	2688	G	A	het	het	tggAagcAcaaggggccC	tggAagcAcaaggggccT	GccgccgatggagactcG
IWGSC_CSS_3AS_scaff.3304956	Cadenza0364	1017	G	A	het	het	gtcccttgacacagctttG	gtcccttgacacagctttA	cctgctggactacaactcaaT
IWGSC_CSS_3AS_scaff.3321091	Cadenza0364	4585	C	T	het	het	caagaatgATgctgatgttggaG	caagaatgATgctgatgttggaA	acatgctgaatgccgaatC

IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Cadenza)	Primer 2 (mutant)	Common Primer
IWGSC_CSS_3AS_scaff_3371333	Cadenza0364	538	G	A	het	het	gggaaaCgAgAcgagcgG	gggaaaCgAgAcgagcgA	cctgtccttctcaccctT
IWGSC_CSS_3AS_scaff_3371815	Cadenza0364	1061	C	T	het	het	atccccacggcacagagG	atccccacggcacagagA	aAttggcccttggtgattcC
IWGSC_CSS_3AS_scaff_3440912	Cadenza0364	4498	G	A	het	het	ccgtaaaaactttctgtgcttgC	ccgtaaaaactttctgtgcttgT	atActgacaaactacatgatgtgC
IWGSC_CSS_3B_scaff_10343586	Cadenza0364	2242	G	A	het	—	ggttcTgTcctctcttccactG	ggttcTgTcctctcttccactA	tgtgttgaaccgcaagcA
IWGSC_CSS_5DL_scaff_242342	Cadenza0281	2433	C	T	hom	hom	catggCgacggtGtcctG	catggCgacggtGtcctA	aAccctcatTTtggCTACTtCT
IWGSC_CSS_5DL_scaff_4538822	Cadenza0281	1208	G	A	hom	—	acgtcagaacaaccgtttgaC	acgtcagaacaaccgtttgaT	ttaaattggttggcgccacC
IWGSC_CSS_6AL_scaff_5813297	Cadenza0281	4532	C	T	hom	—	gggagaggggacgtctcgG	gggagaggggacgtctcgA	ttctctgcacaacgattccG
IWGSC_CSS_6AS_scaff_4378990	Cadenza0281	6748	C	T	hom	hom	cccaggttctgctcttttcC	cccaggttctgctcttttcT	caagtatcaaaaaatgaaggGgT
IWGSC_CSS_6BL_scaff_4360781	Cadenza0281	5426	C	T	het	het	aCtactcaaatggcttGgtgtaG	aCtactcaaatggcttGgtgtaA	tcagtccaacatgTcaagagatT
IWGSC_CSS_7AL_scaff_4488310	Cadenza0281	3808	G	A	hom	hom	gttctctttagtagcagccG	gttctctttagtagcagccA	ggcgctttcttcggcctA
IWGSC_CSS_7BL_scaff_6696509	Cadenza0281	9232	G	A	het	het	gctctaggGgtggcaaAagG	gctctaggGgtggcaaAagA	ggcttGaGgtcGcagtgT
IWGSC_CSS_7BS_scaff_3143575	Cadenza0281	1866	C	T	het	het	agatgttgagagggcgcttC	agatgttgagagggcgcttT	gcttggAtgtgggcaagtT
IWGSC_CSS_7DL_scaff_3346250	Cadenza0281	1663	G	A	het	het	acgtgcagcaacatcctaaC	acgtgcagcaacatcctaaT	TttcccaccaggccaagA
IWGSC_CSS_7DS_scaff_3933917	Cadenza0281	1243	C	T	het	het	tgCtgagcCttTcaccttgC	tgCtgagcCttTcaccttgT	agaggtttggttccatcGG
IWGSC_CSS_3B_scaff_10626860	Cadenza0148	7847	G	A	het	het	gcagctctgggaaggagA	gcagctctgggaaggagA	gttaatgtacCTcctagctcG
IWGSC_CSS_3DL_scaff_6915683	Cadenza0148	6904	C	T	het	het	cgtcaaCctgtgggcaattG	cgtcaaCctgtgggcaattA	tcatgctcataatgTcatagggT
IWGSC_CSS_4AS_scaff_5929057	Cadenza0148	4238	G	A	hom	hom	gcgcaacgtagCacctacC	gcgcaacgtagCacctacT	ttatctggtgaagtgcaggttCA
IWGSC_CSS_4AS_scaff_5950625	Cadenza0148	10590	C	T	het	het	agaTattCaaaTcggtggAttggC	agaTattCaaaTcggtggAttggT	cctgCtccctcacgtcC
IWGSC_CSS_4AS_scaff_5967119	Cadenza0148	11626	C	T	hom	hom	cgtGgacaccccgagctG	cgtGgacaccccgagctA	gacgacgcctgcacgaC
IWGSC_CSS_4DL_scaff_14455742	Cadenza0148	1946	C	T	hom	hom	gCctgagggagatcgcgC	gCctgagggagatcgcgT	aaccgGtAaCTGtGgGcA
IWGSC_CSS_4DS_scaff_2318993	Cadenza0148	4000	C	T	hom	hom	tccagtttgacacagattgaatggG	tccagtttgacacagattgaatggA	tgagaTtctgtttcctttcacAttG
IWGSC_CSS_5AL_scaff_2750707	Cadenza0148	4603	G	A	het	het	ccttgggtgtagccatttcaagTaG	ccttgggtgtagccatttcaagTaA	ccaggaTgcAgtgcaatatttcaagG
IWGSC_CSS_5BL_scaff_10794137	Cadenza0148	9235	C	T	hom	hom	gaagctgcttctcgcttG	gaagctgcttctcgcttA	agtatcccttccatataagcagtG
IWGSC_CSS_5BS_scaff_1646558	Cadenza0148	2916	C	T	het	het	gccGtacactcacctAtcctttG	gccGtacactcacctAtcctttA	gcaaTgtccacttAtcatcccT
IWGSC_CSS_1AL_scaff_3883106	Cadenza0110	27536	C	T	het	het	accttccatcactggctgG	accttccatcactggctgA	gtgaagaacaacaggttgaagC
IWGSC_CSS_1BL_scaff_3812829	Cadenza0110	10770	G	A	het*	hom	ccccactccattccagA	ccccactccattccagA	gGatgtgttctgtgctggaA
IWGSC_CSS_1DL_scaff_2266648	Cadenza0110	6156	G	A	het	het	actgcgtggttatgggacC	actgcgtggttatgggacT	ccccatcactgaacacaacA
IWGSC_CSS_1DS_scaff_1889435	Cadenza0110	8826	C	T	hom	hom	aaccatgaattactcggacagG	aaccatgaattactcggacagA	gcctgaagaattgtatcaaaacaG
IWGSC_CSS_2AS_scaff_5268634	Cadenza0110	4636	G	A	het	het	gatccatgtgattggcatgtttG	gatccatgtgattggcatgtttA	TgctgtTggatagtcagttacT
IWGSC_CSS_2BL_scaff_7965110	Cadenza0110	15801	C	T	hom	hom	cattgaagcAtacacAattgcAtaC	cattgaagcAtacacAattgcAtaT	gccagagatccagataaggTttA
IWGSC_CSS_2DL_scaff_9852812	Cadenza0110	13788	G	A	hom	hom	atttttgtatggtctcaatcttcgC	atttttgtatggtctcaatcttcgT	gaacgtTcattctgtactgtcT
IWGSC_CSS_2DS_scaff_5371379	Cadenza0110	2166	C	T	hom	hom	agacacaaaactagtGatgcgC	agacacaaaactagtGatgcgT	gctgctgagaatgttTtgtatttG
IWGSC_CSS_3AL_scaff_4384278	Cadenza0110	1276	C	T	het	het	agcTgaactgccccTgtaG	agcTgaactgccccTgtaA	agggaacctCgGtgatgaA
IWGSC_CSS_3AS_scaff_3340122	Cadenza0110	1467	C	T	hom	hom	attcctAgtgttgcggaacatG	attcctAgtgttgcggaacatA	gagaagactagaaaagttttcAgcaT
IWGSC_CSS_5DL_scaff_4554222	Cadenza2103	6528	C	T	het*	hom	gctgccctacaaagaaacaaattG	gctgccctacaaagaaacaaattA	aTcccaactatCGaTtttgctataC
IWGSC_CSS_6AL_scaff_5833640	Cadenza2103	7346	C	T	hom	hom	aagaaaaagccacaatggtttctC	aagaaaaagccacaatggtttctT	aCTctgTcagtgtttcccgC
IWGSC_CSS_6AS_scaff_4429974	Cadenza2103	3867	G	A	hom	hom	GagatgaAttatttgagcatgtggC	GagatgaAttatttgagcatgtggT	ggttccggctgcataagT
IWGSC_CSS_6DL_scaff_3307626	Cadenza2103	4970	C	T	hom	hom	tgcagatgttgcctgtgtaG	tgcagatgttgcctgtgtaA	tgtagaaggtgattttgtactGtC
IWGSC_CSS_6DS_scaff_2059604	Cadenza2103	5224	G	A	het	—	gctcaatgcatgcTgagtgG	gctcaatgcatgcTgagtgA	tgtcaagtattattttcctgctcG
IWGSC_CSS_7AL_scaff_4552322	Cadenza2103	1412	C	T	het	het	gcaaaggcTgatactccaacaG	gcaaaggcTgatactccaacaA	ggcAAGccAgtataaaagtaaGC

IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Cadenza)	Primer 2 (mutant)	Common Primer
IWGSC_CSS.7BS_scaff.3147455	Cadenza2103	4607	G	A	het	—	gcaccttaggatgtgagTtatgC	gcaccttaggatgtgagTtatgT	gcatgtagggtttatttgactgttA
IWGSC_CSS.7DL_scaff.3382467	Cadenza2103	3473	C	T	hom	—	GGTtctgCaGTTTCATAActcatC	GGTtctgCaGTTTCATAActcatT	attgaatcaactgatacGaaGactC
IWGSC_CSS.3B_scaff.10457010	Cadenza0277	10599	G	A	het	het	aaccttggccgcagaacaC	aaccttggccgcagaacaT	actggtcgcacgagaggG
IWGSC_CSS.3B_scaff.10593852	Cadenza0277	10124	C	T	het	het	tgacaggggacgtatacaG	tgacaggggacgtatacaA	gtctaaCTtACattAcccatcagC
IWGSC_CSS.3DS_scaff.2583390	Cadenza0277	663	G	A	hom	hom	actgcactcatacaatActtCtgC	actgcactcatacaatActtCtgT	tcCacctggacagacaagtG
IWGSC_CSS.4AL_scaff.7093953	Cadenza0277	10004	C	T	hom	hom	ccttgattcaatggaTtgTtttgG	ccttgattcaatggaTtgTtttgA	ttcccaaaaTaaaaggaagagC
IWGSC_CSS.4AL_scaff.7176064	Cadenza0277	6220	C	T	het	het	gtgccgtaTtcCgcctgG	gtgccgtaTtcCgcctgA	atgttcgaggggatgggG
IWGSC_CSS.4DL_scaff.14122349	Cadenza0277	1010	C	T	hom	hom	gtcgctgctgCttgtgaG	gtcgctgctgCttgtgaA	ggaacaggcccaaggagG
IWGSC_CSS.5AL_scaff.2736916	Cadenza0277	4296	G	A	het	het	aagaactATgAaaGtaacacacgaC	aagaactATgAaaGtaacacacgaT	ttcGcTttTaagGcAttCtcG
IWGSC_CSS.5BL_scaff.10883744	Cadenza0277	2080	C	T	hom	hom	gcctctttCtgttTagcctcaG	gcctctttCtgttTagcctcaA	cgacaaggttcgtgatTgcA
IWGSC_CSS.1AL_scaff.3932013	Cadenza0548	11765	C	T	hom	hom	accgccaaCccaagacaG	accgccaaCccaagacaA	cccatTAcccGgcGtgcAacG
IWGSC_CSS.1BS_scaff.3417505	Cadenza0548	373	C	T	het	het	gtggtgaggaGGgtgGaG	gtggtgaggaGGgtgGaA	tggtcgcGccagttgttgA
IWGSC_CSS.2AS_scaff.5305619	Cadenza0548	2786	C	T	hom	hom	atacagatgcctAAGtggTtC	atacagatgcctAAGtggTtT	ggaagacaAtGctccaggaC
IWGSC_CSS.2AS_scaff.5306489	Cadenza0548	46953	T	G	het	wt	aggttcattgtccatagaagGT	aggttcattgtccatagaagGG	aggctaTAGactcctgTACAgT
IWGSC_CSS.2BL_scaff.7984123	Cadenza0548	11660	G	A	het	het	cattgtggcatagtaatcagtacaG	cattgtggcatagtaatcagtacaA	aatacattgaggaatacaagccC
IWGSC_CSS.2DL_scaff.9907477	Cadenza0548	1363	C	T	hom	hom	tgctccctttgccagaaC	tgctccctttgccagaaT	ggcaaacctgatgtggcatC
IWGSC_CSS.2DS_scaff.5330886	Cadenza0548	5449	G	A	hom	hom	gcattgtccattataactgaacGtG	gcattgtccattataactgaacGtA	catgctgtctctctggacC
IWGSC_CSS.3AL_scaff.4449951	Cadenza0548	633	C	T	het	het	tccaaacctaacagtcataactaG	tccaaacctaacagtcataactaA	gtctgcagTGCaatgtgC
IWGSC_CSS.3B_scaff.10479889	Cadenza0097	3339	C	T	hom	—	ttgTttctGgagaagatgcCG	ttgTttctGgagaagatgcCA	ggtgtcattcaAcGgcA
IWGSC_CSS.3B_scaff.10562262	Cadenza0097	7819	C	T	het	het	agaggggtgctatccatAttgG	agaggggtgctatccatAttgA	agcgatccaaggcttcC
IWGSC_CSS.4AL_scaff.7040796	Cadenza0097	10772	G	A	hom	hom	acacaacattgccaccagaG	acacaacattgccaccagaA	CAatCgattgtctgtTctcC
IWGSC_CSS.4AL_scaff.7063488	Cadenza0097	6360	C	T	het	het	gcctctcacCttAattgaaagctgC	gcctctcacCttAattgaaagctgT	aggcagtgagatgtggaagttT
IWGSC_CSS.4AL_scaff.7091701	Cadenza0097	5050	G	A	het	het	catgagcatctgggaggaaaatG	catgagcatctgggaggaaaatA	agcaagggaAtaatgaacggaaA
IWGSC_CSS.4DS_scaff.1845841	Cadenza0097	7110	G	A	hom	hom	aatgTAGctccccatacCgG	aatgTAGctccccatacCgA	actgaacTgcaatcgtTtatggA
IWGSC_CSS.5AL_scaff.2767581	Cadenza0097	3737	G	A	het	het	gagaggtcctcactAtcggC	gagaggtcctcactAtcggT	cgTcatcacaatattgtcggG
IWGSC_CSS.5BL_scaff.10784643	Cadenza0097	1568	C	T	hom	hom	agaaaTAcatggatggatggaCG	agaaaTAcatggatggatggaCA	catctcCCttcaCgGaaaG
IWGSC_CSS.1AL_scaff.3952258	Cadenza2092	8107	C	T	het	—	tgagtagagaaattgacagtgtgG	tgagtagagaaattgacagtgtgA	tgccaccattgacatgagaG
IWGSC_CSS.1BL_scaff.3858008	Cadenza2092	10278	G	A	hom	hom	tttgagcaggcaggatcgC	tttgagcaggcaggatcgT	actcacggcctatacActattC
IWGSC_CSS.1DL_scaff.2265172	Cadenza2092	9094	C	T	hom	hom	tgcaTGTcatttgttcttatcagC	tgcaTGTcatttgttcttatcagT	agtgtccaacttccGttcatC
IWGSC_CSS.2AL_scaff.6435867	Cadenza2092	16201	G	A	hom	hom	tttctgTactttaacgtcaattgaC	tttctgTactttaacgtcaattgaT	gtgaggatgatgaggaagacC
IWGSC_CSS.2AL_scaff.6439430	Cadenza2092	25101	C	T	het	—	caagaaagggCagCtCagC	caagaaagggCagCtCagT	tcGttAcTctttcActggtgaA
IWGSC_CSS.2DL_scaff.9760848	Cadenza2092	4733	C	T	het	het	gcaccatgggtctcaggtaC	gcaccatgggtctcaggtaT	tcagtcagtttGCTCgtTCTG
IWGSC_CSS.3AL_scaff.4407012	Cadenza2092	2785	C	T	hom	hom	acatatAgtgttctcatccaccatC	acatatAgtgttctcatccaccatT	acctctcatgttaaataggtttgT
IWGSC_CSS.3AS_scaff.3441108	Cadenza2092	541	G	A	het	het	GtgatgaccttgagacGgaC	GtgatgaccttgagacGgaA	aggcaTgacaaCgcgcaA
IWGSC_CSS.3B_scaff.10449827	Cadenza1551	4779	G	A	hom	hom	ggcaaggtcaagaaacGgtC	ggcaaggtcaagaaacGgtT	aCagaGtgggttagaggcaG
IWGSC_CSS.3B_scaff.10550638	Cadenza1551	3250	C	T	het	het	ctccttcacttgttgcggC	ctccttcacttgttgcggT	gcaacATtTgatactgcaagG
IWGSC_CSS.3DL_scaff.6945816	Cadenza1551	589	C	T	hom	hom	agcatctcacctgcaacCaataC	agcatctcacctgcaacCaataT	TgtgcccTctgaAtattttcaTG
IWGSC_CSS.3DL_scaff.6954177	Cadenza1551	3508	C	T	het	het	tgtagcatcacattaaactttctG	tgtagcatcacattaaactttctA	gcttggtataaacCttacgacA
IWGSC_CSS.4AS_scaff.5938272	Cadenza1551	19080	G	A	hom	hom	agAcCccgAtcgccatgG	agAcCccgAtcgccatgA	GggAgatAcaggtaaaActcTtcG
IWGSC_CSS.4AS_scaff.5977594	Cadenza1551	11092	C	T	het	het	gccttgattcggaacaacaaaC	gccttgattcggaacaacaaaT	gcgtctctcagtcctgcA

IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Cadenza)	Primer 2 (mutant)	Common Primer
IWGSC_CSS_5AL_scaff_2671035	Cadenza1551	5859	C	T	het	het	cggatgatattTttagacttcgacG	cggatgatattTttagacttcgacG	ggcagttcagcGacccatT
IWGSC_CSS_5BL_scaff_10889480	Cadenza1551	2530	G	A	hom	hom	gagcttaactcgagatggaG	gagcttaactcgagatggaA	tccatgCAacGccttgG
IWGSC_CSS_3B_scaff_10528396	Cadenza2088	8059	G	A	hom	—	ctttccgctccgtaagcaataG	ctttccgctccgtaagcaataA	gtgcaactgttcaggcctgA
IWGSC_CSS_3B_scaff_10637573	Cadenza2088	16815	G	A	het	het	agcaagcttaccGgtctgC	agcaagcttaccGgtctgT	cgagcAactacgagcagctT
IWGSC_CSS_4AL_scaff_7086469	Cadenza2088	6697	G	A	het	het	gccgtctacttcaacgcG	gccgtctacttcaacgcA	ccaGaggcttgTGCattttT
IWGSC_CSS_4AL_scaff_7126302	Cadenza2088	3627	G	A	hom	hom	gttcaaaaacaagtggtAatttgC	gttcaaaaacaagtggtAatttgT	cacaaggatgatgaagcTctcttagA
IWGSC_CSS_4BL_scaff_7041808	Cadenza2088	10234	G	A	hom	hom	tcaatggatgagggtgcttC	tcaatggatgagggtgcttT	ccatagcagcatcagccacA
IWGSC_CSS_5AL_scaff_2794167	Cadenza2088	13162	G	A	het	—	agtattcaggacaagcatCttCaG	agtattcaggacaagcatCttCaA	caatgaacctctcgaagaaGaG
IWGSC_CSS_5BL_scaff_10889232	Cadenza2088	3885	G	A	het	het	cTcaaccacaatgggcaAatC	cTcaaccacaatgggcaAatT	tccttcatcaatcatcaattgttG
IWGSC_CSS_5BS_scaff_2267405	Cadenza2088	11113	C	T	hom	hom	ctttgatgatcctaggcctctTG	ctttgatgatcctaggcctctTA	tgatttggTctggtAgagtttGA
IWGSC_CSS_3B_scaff_10475354	Cadenza1409	2203	G	A	hom	hom	agCGaacaagagGtcaaacG	agCGaacaagagGtcaaacA	ctgaaacacaCtagaCAattAccG
IWGSC_CSS_3B_scaff_10674115	Cadenza1409	4555	C	T	het	het	gcttcagtgcatgccttcaG	gcttcagtgcatgccttcaA	cttcacaccGagataatGtattG
IWGSC_CSS_4AL_scaff_7153568	Cadenza1409	13073	C	T	hom	hom	tccgaccgAtcaaccttgG	tccgaccgAtcaaccttgA	gaccggaactcctcggcC
IWGSC_CSS_4DL_scaff_14314966	Cadenza1409	2010	G	A	het	hom	gtaggccccctctCAGgG	gtaggccccctctCAGgA	cggcgTcaCaAgttgCcT
IWGSC_CSS_4DS_scaff_2324074	Cadenza1409	7606	G	A	het	het	tGcatgaaaatgtgtGcaGaG	tGcatgaaaatgtgtGcaGaA	gggtaAgttcAaaactGaaagtgaG
IWGSC_CSS_5AS_scaff_1517889	Cadenza1409	3561	G	A	het	het	tctcgacatcttcccggtgaC	tctcgacatcttcccggtgaT	gtgccttgaacattgcttattA
IWGSC_CSS_5AS_scaff_1523866	Cadenza1409	8054	G	A	hom	—	ggatgatctaccgcaGgaC	ggatgatctaccgcaGgaT	tctcgagCcTctctcA
IWGSC_CSS_5BL_scaff_10917655	Cadenza1409	19073	G	A	hom	hom	caaatgacatgcaaaagaagttgC	caaatgacatgcaaaagaagttgT	cgcttcatcactacaAaatatgtcT
IWGSC_CSS_1AL_scaff_3886649	Cadenza1599	5204	C	T	het	het	tgatgcccaaccacaatGcC	tgatgcccaaccacaatGcT	ggactgactgtgaccatatttaG
IWGSC_CSS_1BL_scaff_3810267	Cadenza1599	6634	C	T	hom	hom	ccCaggaaatgagcacctC	ccCaggaaatgagcacctT	cgcaggcgaagatgtgaTtG
IWGSC_CSS_1DL_scaff_2291677	Cadenza1599	12856	C	T	hom	hom	GgtagacaagtcgccgaG	GgtagacaagtcgccgaA	cctcctccttcaacGCcG
IWGSC_CSS_2AL_scaff_6354492	Cadenza1599	7566	G	A	het	het	gGagaatgcaCAgtAacTtctgG	gGagaatgcaCAgtAacTtctgA	ttccgaagaaccacaTccTG
IWGSC_CSS_2AS_scaff_5282937	Cadenza1599	9736	G	A	het	het	gctgtagattttatagctgctatgC	gctgtagattttatagctgctatgT	cacCagaattgttCactgatttTC
IWGSC_CSS_2BL_scaff_7952427	Cadenza1599	19249	G	A	hom	hom	cgTccctCcttagcacgaC	cgTccctCcttagcacgaT	aTcactcattagcgcgAG
IWGSC_CSS_2DL_scaff_9897981	Cadenza1599	5627	C	T	het	het	cttgggtgctTgattgcttactC	cttgggtgctTgattgcttactT	gTttgctCtctgctatCtTgtG
IWGSC_CSS_3AL_scaff_4446105	Cadenza1599	1765	G	A	hom	—	aaatgctttcctaCcgctagtG	aaatgctttcctaCcgctagtA	ttctAgaggcaatagctTatatgcT

Table A.5: Validation of mutations on M_4 on Kronos

IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Kronos)	Primer 2 (mutant)	Common Primer
IWGSC_CSS_1AS_scaff_3284790	Kronos3085	7449	G	A	Het	Het	ccacaccttgagcctcgC	ccacaccttgagcctcgT	gtgattttgccaggggagA
IWGSC_CSS_1BL_scaff_3897513	Kronos3085	1515	C	T	Het	Het	gcttcactGggtcctgC	gcttcactGggtcctgT	acAaggactgcttcagaGaC
IWGSC_CSS_2AL_scaff_6434745	Kronos3085	3424	C	T	Het	Het	cctcGgttttgcaaatttctatgC	cctcGgttttgcaaatttctatgT	gGCaaTggcataacaacagatA
IWGSC_CSS_3AS_scaff_3408995	Kronos3085	732	C	T	Het	Het	aggccatttcgaattccgC	aggccatttcgaattccgT	ggTgttaTccagAacctgagTG
IWGSC_CSS_3B_scaff_10708748	Kronos3085	2675	G	A	Het	Het	gttgcatgcttcacccagG	gttgcatgcttcacccagA	gtaacaactctgagttcgtagcaC
IWGSC_CSS_4AL_scaff_7132733	Kronos3085	1799	C	T	Hom	Hom	caccctgtagtgaccctC	caccctgtagtgaccctT	aCcGcctaGaaagaaagcttC


IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Kronos)	Primer 2 (mutant)	Common Primer
IWGSC.CSS_5AS_scaff.1534693	Kronos3085	4605	C	T	Het	Het	cagcttctctggccctcAtC	cagcttctctggccctcAtT	gtaCctcagcAgtcaTgagAG
IWGSC.CSS_6AS_scaff.4361911	Kronos3085	8857	G	A	Het	Het	tcacgaaagacgacttcaacctcC	tcacgaaagacgacttcaacctcT	catgagggtctgcatctccatcA
IWGSC.CSS_6BS_scaff.3008326	Kronos3085	1528	G	A	Het	Het	ccatgttgacttggtggtgC	ccatgttgacttggtggtgT	ggaagcatggCaagtgcA
IWGSC.CSS_7AS_scaff.4214385	Kronos3085	27835	C	T	Hom	Hom	cgtaccttcggtgggaaagG	cgtaccttcggtgggaaagA	ctcttggtcagctgataaagacT
IWGSC.CSS_1AL_scaff.3929964	Kronos3191	1336	C	T	Het	Het	tttcggccataacctgacatC	tttcggccataacctgacatT	attgctccagcttcttgcaG
IWGSC.CSS_1BL_scaff.3899789	Kronos3191	7925	C	T	Het	Het	actctcacTggcagcagC	actctcacTggcagcagT	caactgtggtgcccatcGtA
IWGSC.CSS_2AL_scaff.6426728	Kronos3191	1481	G	A	Hom	Hom	gaaActgccgcagctCgC	gaaActgccgcagctCgT	ccaGcaGctcgtgagaaA
IWGSC.CSS_2BL_scaff.7960273	Kronos3191	690	C	T	Hom	Hom	gccattcatccttaggcgC	gccattcatccttaggcgT	acatgcaattgctgatgactG
IWGSC.CSS_3AS_scaff.3286603	Kronos3191	2975	G	A	Het*	Hom	ccgtgtggtttgttggtgG	ccgtgtggtttgttggtgA	gaaaggaacgtgTcaTgcaG
IWGSC.CSS_5AL_scaff.2694249	Kronos3191	2399	C	T	Het	Het	gccttcagatagagccGC	gccttcagatagagccGT	cgccacatcgacattctcG
IWGSC.CSS_5BL_scaff.10923577	Kronos3191	3713	C	T	Het	Het	gtggattgcctgagcttgC	gtggattgcctgagcttgT	tgggtggcctcttgggaC
IWGSC.CSS_6AL_scaff.5823017	Kronos3191	13225	C	T	Hom	Hom	ccctttcagcctctggaG	ccctttcagcctctggaA	ttcgagaaggcccatcgA
IWGSC.CSS_6BS_scaff.2955394	Kronos3191	1622	C	T	Het*	Hom	gtggagatgaaggtctagcaaG	gtggagatgaaggtctagcaaA	gatactcTgcaatgggtgT
IWGSC.CSS_7BL_scaff.6739382	Kronos3191	12261	G	A	Hom	Hom	gagacaagctttgaattgctcC	gagacaagctttgaattgctcT	CgagtgcctTcatttcccG
IWGSC.CSS_1AS_scaff.3276389	Kronos3288	9720	C	T	Hom	Hom	aCcaGcaggaccAatgtctC	aCcaGcaggaccAatgtctT	atgatgcaactcagccaT
IWGSC.CSS_2AL_scaff.6367515	Kronos3288	6976	G	A	Het	Het	caggtcgagTgtctccgG	caggtcgagTgtctccgA	ggggtgatCtggaaaggG
IWGSC.CSS_2AL_scaff.6422019	Kronos3288	4523	G	A	Het	Het	cgctaggtccctgcataG	cgctaggtccctgcataG	acgcAcgttaagccgtA
IWGSC.CSS_3AL_scaff.4284850	Kronos3288	7901	C	T	Hom	Hom	tgctttggacaacatcgG	tgctttggacaacatcgA	tgtcAgtcagcagcccaG
IWGSC.CSS_4AS_scaff.5962359	Kronos3288	13049	G	A	Het	Hom	ccatcaagaagtacgagttcgaC	ccatcaagaagtacgagttcgaT	accatgccagcttgtaA
IWGSC.CSS_6AL_scaff.5778773	Kronos3288	6853	G	A	Het	Het	gagtgaccttcccgtcttC	gagtgaccttcccgtcttT	ggagaacagctactcggcT
IWGSC.CSS_6AS_scaff.4392100	Kronos3288	3434	C	T	Het	Het	atggaagcacaggtgaccG	atggaagcacaggtgaccA	ggAagcgaaagtgaacaaA
IWGSC.CSS_7BL_scaff.6744240	Kronos3288	9772	G	A	Het	Het	agctgttcttctcctacttcaaG	agctgttcttctcctacttcaaA	caggtcgttcttgagctcC
IWGSC.CSS_1AL_scaff.3887185	Kronos3413	9708	C	T	Hom	Hom	gcacgcctttatcgaggtaaaG	gcacgcctttatcgaggtaaaA	AgaacagcagagcgcaA
IWGSC.CSS_2BS_scaff.3381362	Kronos3413	5160	C	T	Het*	Hom	caacttctgggctgtagtG	caacttctgggctgtagtA	tgAgaattctgacGcaaaagaC
IWGSC.CSS_3AS_scaff.3296605	Kronos3413	6154	G	A	Het	Het	ctggtcacgggctctagC	ctggtcacgggctctagT	cagcactgagacatggaC
IWGSC.CSS_3BL_scaff.10693516	Kronos3413	12632	C	T	Het	Het	ctaggcttgacaaaacaggC	ctaggcttgacaaaacaggT	agcttgcatctatgggcatT
IWGSC.CSS_5AS_scaff.1547699	Kronos3413	2686	G	A	Het	Het	gCtacaaccttcaccaatcgC	gCtacaaccttcaccaatcgT	gacggctttgaagtgtcatC
IWGSC.CSS_5BL_scaff.10856077	Kronos3413	5853	G	A	Het	Het	agagcttcaccccatgctC	agagcttcaccccatgctT	acgCacatttAatagctgaagC
IWGSC.CSS_6AL_scaff.5750718	Kronos3413	11046	G	A	Hom	Hom	cacgcTtcccgaacttctataG	cacgcTtcccgaacttctataA	AgacgatgtgatcaggattcaG
IWGSC.CSS_7AL_scaff.4433177	Kronos3413	3511	C	T	Het	Het	GaTgctccGtcaggctgG	GaTgctccGtcaggctgA	cactactggacaagctcttgG
IWGSC.CSS_7BL_scaff.6742567	Kronos3413	667	C	T	Het	Het	gttgcttgctggcgagaC	gttgcttgctggcgagaT	cattttgcacctgtgtcTG
IWGSC.CSS_1AL_scaff.3976389	Kronos3935	10941	C	T	Hom	Hom	gggtgaggagatcggCgatG	gggtgaggagatcggCgatA	cagtcactcatagagaggtcaG
IWGSC.CSS_1BL_scaff.3873362	Kronos3935	1392	G	A	Het	Het	cagatctgaagcctaGcacatG	cagatctgaagcctaGcacatA	actaccagaatcagcacaacaaAC
IWGSC.CSS_2BL_scaff.7882382	Kronos3935	2721	C	T	Het	Het	gcaagctaagatgtaccgtagC	gcaagctaagatgtaccgtagT	gccacagttaggagaagactT
IWGSC.CSS_3AL_scaff.4242376	Kronos3935	2410	C	T	Het	Het	agaacccaaaacccgTacttaG	agaacccaaaacccgTacttaA	gtagGgtCcatCTaaagcttG
IWGSC.CSS_3B_scaff.10485067	Kronos3935	3349	C	T	Hom	Hom	gcttgagcaactactccaatG	gcttgagcaactactccaatA	gcaatttcccttaTccgcagT
IWGSC.CSS_4AS_scaff.5984153	Kronos3935	6006	G	A	Het	Het	agCaggctctggccaagttG	agCaggctctggccaagttA	cgaatGtatgaGtaggcgT
IWGSC.CSS_4BL_scaff.7019402	Kronos3935	9081	C	T	Het	Het	tgcaatcatgtagtgcctgG	tgcaatcatgtagtgcctgA	agcatgatccctagaaCCataC
IWGSC.CSS_5BL_scaff.10842786	Kronos3935	3304	G	A	Het	Het	tggttcccGaagcctgaaC	tggttcccGaagcctgaaT	cgcatacttgaacaTGagcAC
IWGSC.CSS_6BS_scaff.3045205	Kronos3935	2293	G	A	Het	Het	aaggaccaagcccaactctcG	aaggaccaagcccaactctcA	agtgatcaagcccaatgtgcA


IWGSC contig	Line	Pos	WT	Mut	Predicted	M_4	Primer 1 (Kronos)	Primer 2 (mutant)	Common Primer
IWGSC.CSS_7AL_scaff.4555249	Kronos3935	4487	C	T	Het	Het	cAgtgctcgagatggcgC	cAgtgctcgagatggcgT	cCttgcaacctctctgatT
IWGSC.CSS_1BL_scaff.3918498	Kronos4240	6096	G	A	Het	Het	ttgcatgcccccaagaagaG	ttgcatgcccccaagaagaA	tgggcgaactgggtaatgtgG
IWGSC.CSS_2BS_scaff.5131713	Kronos4240	5900	G	A	Het	Het	cctttatcgaggaaaagagacacC	cctttatcgaggaaaagagacacT	caccattgtagggttcctTttC
IWGSC.CSS_5AL_scaff.2769540	Kronos4240	9626	C	T	Het	Het	tgCagtgtgggaaacggaG	tgCagtgtgggaaacggaA	catgagtGagatcttctctgcT
IWGSC.CSS_5BL_scaff.10871091	Kronos4240	7062	G	A	Het	Het	gccaaaggAaccataacctgC	gccaaaggAaccataacctgT	GgactcttggcAaccggA
IWGSC.CSS_6AL_scaff.5800333	Kronos4240	2360	G	A	Het	Het	cgacaggattgtgagCgC	cgacaggattgtgagCgT	tcagatgctgcaagattcatcT
IWGSC.CSS_7BL_scaff.6716931	Kronos4240	2613	G	A	Het	Het	gGtgGgtattTgcttgggtgaG	gGtgGgtattTgcttgggtgaA	tgGtggactcgacaGtGtA
IWGSC.CSS_2BL_scaff.8029221	Kronos4346	2860	G	A	Het	Het	tgcttccgctcttgctcC	tgcttccgctcttgctcT	atTtgcATCgAtcgggcC
IWGSC.CSS_3B_scaff.10460714	Kronos4346	14359	C	T	Hom	Hom	ctaccttgccatgcgacatG	ctaccttgccatgcgacatA	agcaccccgactctttgacG
IWGSC.CSS_4AS_scaff.5989735	Kronos4346	6404	G	A	Hom	Hom	acgcatgctaacatcagcG	acgcatgctaacatcagcT	actcaagataccaCcgcacG
IWGSC.CSS_5BL_scaff.7648030	Kronos4346	6893	C	T	Het	Het	taccttttctactggcagG	taccttttctactggcagA	ttttcagaggaaacacaggtatcA
IWGSC.CSS_6AL_scaff.5755840	Kronos4346	778	C	T	Het	Het	atcgagtaagctgtcacCgC	atcgagtaagctgtcacCgT	acctgcatgtcaCatccaC
IWGSC.CSS_6BS_scaff.2972151	Kronos4346	7876	G	A	Hom	Hom	gcagcaatgtcActgtttgG	gcagcaatgtcActgtttgA	gcttggactgggcattttatG
IWGSC.CSS_7AL_scaff.4542983	Kronos4346	18700	G	A	Het	Het	gcagggctAccggatacC	gcagggctAccggatacT	catctgccGgttaaacatgC
IWGSC.CSS_7BS_scaff.3098098	Kronos4346	5183	C	T	Het	Het	gCgatatgggtacttgcaatgaG	gCgatatgggtacttgcaatgaA	ttacattgcttataG'TttgCcgG
IWGSC.CSS_1AS_scaff.3259804	Kronos4485	219	C	T	Het	Het	gtcggcacaacccttgC	gtcggcacaacccttgT	gcttctttaaggaggggcga
IWGSC.CSS_2AL_scaff.6315418	Kronos4485	10490	G	A	Hom	Hom	gccccctctcaaCcttctcagC	gccccctctcaaCcttctcagT	ttcagacgtCGagggaatttcC
IWGSC.CSS_2BS_scaff.5181092	Kronos4485	3742	G	A	Het	Het	TggccagcacacctgcaG	TggccagcacacctgcaA	tggacgatgagTgatggAaaT
IWGSC.CSS_3B_scaff.10425015	Kronos4485	2372	C	T	Het	Het	gctactgaagttggCtcGG	gctactgaagttggCtcGA	cttcacatccttgggggTtC
IWGSC.CSS_3B_scaff.10775915	Kronos4485	4701	C	T	Het	Het	ccaagggctgcagagagG	ccaagggctgcagagagA	agacctcacgatGtcctcC
IWGSC.CSS_5AL_scaff.2754304	Kronos4485	2301	G	A	Het	Het	taaccTgccatcgcccG	taaccTgccatcgcccA	cattgGccagccaTgacT
IWGSC.CSS_5BL_scaff.10919959	Kronos4485	1867	C	T	Hom	Hom	gatgccctttgtggagaagG	gatgccctttgtggagaagA	tcttgttcccgaacatgtcA
IWGSC.CSS_7AS_scaff.4245431	Kronos4485	3402	G	A	Hom	Hom	aaggcgctgtgtttcC	aaggcgctgtgtttcT	agtaagtggAacagctaagatcaT
IWGSC.CSS_7BL_scaff.6667357	Kronos4485	641	C	T	Het	Het	gatcAgctgctcattcgagG	gatcAgctgctcattcgagA	ttccctgtcaattgatgccC

Appendix B


Quality Control









B.1 Sequence read quality





B.2 Sequence GC content



Sample	Read 1	Read 2
LIB1717 Bulk R3		
LIB1718 Bulk S1		
LIB1719 Bulk S2		
LIB1720 Bulk S3		

Bibliography

- Abe, A., Kosugi, S., Yoshida, K., et al. Genome sequencing reveals agronomically important loci in rice using MutMap. *Nature biotechnology*, 30(2):174–8, February 2012. ISSN 1546-1696. doi: 10.1038/nbt.2095.
- Akhunov, E. D., Goodyear, A. W., Geng, S., et al. The organization and rate of evolution of wheat genomes are correlated with recombination rates along chromosome arms. *Genome research*, 13(5):753–63, May 2003. ISSN 1088-9051. doi: 10.1101/gr.808603.
- Allen, A. M., Barker, G. L. a., Berry, S. T., et al. Transcript-specific, single-nucleotide polymorphism discovery and linkage analysis in hexaploid bread wheat (*Triticum aestivum* L.). *Plant biotechnology journal*, 9(9):1086–99, December 2011. ISSN 1467-7652. doi: 10.1111/j.1467-7652.2011.00628.x.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. Basic local alignment search tool. *Journal of molecular biology*, 215(3): 403–10, October 1990. ISSN 0022-2836. doi: 10.1016/S0022-2836(05)80360-2.
- Babraham Bioinformatics. FastQC A Quality Control tool for High Throughput Sequence Data, 2012. URL <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Bonnal, R. J. P., Aerts, J., Githinji, G., et al. Biogem: An effective tool-based approach for scaling up open source software development in bioinformatics. *Bioinformatics*, 28(7):1035–1037, April 2012. ISSN 13674803. doi: 10.1093/bioinformatics/bts080.
- Brenchley, R., Spannagl, M., Pfeifer, M., et al. Analysis of the bread wheat genome using whole-genome shotgun sequencing. *Nature*, 491 (7426):705–710, Nov 2012. ISSN 0028-0836. doi: 10.1038/nature11650.

- Burt, C., Steed, A., Gosman, N., et al. Mapping a type 1 fhb resistance on chromosome 4as of triticum macha and deployment in combination with two type 2 resistances. *Theoretical and Applied Genetics*, 128(9): 1725–1738, 2015. ISSN 1432-2242. doi: 10.1007/s00122-015-2542-9.
- Cantu, D., Govindarajulu, M., Kozik, A., et al. Next generation sequencing provides rapid access to the genome of *Puccinia striiformis* f. sp. *tritici*, the causal agent of wheat stripe rust. *PLoS ONE*, 6(8):1–8, 08 2011. doi: 10.1371/journal.pone.0024230.
- Chapman, J. a., Mascher, M., Buluç, A., et al. A whole-genome shotgun approach for assembling and anchoring the hexaploid bread wheat genome. *Genome Biology*, 16(1):1–17, 2015. ISSN 1465-6906. doi: 10.1186/s13059-015-0582-8.
- Clark, M. Unlocking bread wheat genomes diversity with new sequencing and assembly approaches, 1 2016. URL <https://pag.confex.com/pag/xxiv/webprogram/Paper20198.html>.
- Cornish-Bowden, A. Nomenclature for incompletely specified bases in nucleic acid sequences: recommendations 1984. *Nucleic acids research*, 13(9):3021–30, May 1985. ISSN 0305-1048.
- Eddy, S. R. What is a hidden Markov model? *Nature biotechnology*, 22(10):1315–6, October 2004. ISSN 1087-0156. doi: 10.1038/nbt1004-1315.
- Etherington, G. J., Ramirez-Gonzalez, R. H., and MacLean, D. bio-samtools 2: a package for analysis and visualization of sequence and alignment data with SAMtools in Ruby. *Bioinformatics*, pages 1–2, 2015. ISSN 1367-4803. doi: 10.1093/bioinformatics/btv178.
- Gerechter-Amitai, Z. K., van Silfhout, C. H., Grama, A., and Kleitman, F. Yr15 — a new gene for resistance to puccinia striiformis in triticum dicoccoides sel. g-25. *Euphytica*, 43(1):187–190, 1989. ISSN 1573-5060. doi: 10.1007/BF00037912.
- GM, C. Chloroplasts and Other Plastids. In *The Cell: A Molecular Approach*. Sinauer Associates, 2000. URL http://www.ncbi.nlm.nih.gov/books/NBK9905/?redirect-on-error=__HOME__.

- Goto, N., Prins, P., Nakao, M., et al. BioRuby: bioinformatics software for the Ruby programming language. *Bioinformatics (Oxford, England)*, 26(20):2617–9, October 2010. ISSN 1367-4811. doi: 10.1093/bioinformatics/btq475.
- Henry, I. M., Nagalakshmi, U., Lieberman, M. C., et al. Efficient Genome-Wide Detection and Cataloging of EMS-Induced Mutations Using Exome Capture and Next-Generation Sequencing. *The Plant cell*, 26(4):1382–1397, April 2014. ISSN 1532-298X. doi: 10.1105/tpc.113.121590.
- Hodges, E., Xuan, Z., Balija, V., et al. Genome-wide in situ exon capture for selective resequencing. *Nat Genet*, 39(12):1522–1527, Dec 2007. ISSN 1061-4036. doi: 10.1038/ng.2007.42.
- Hubbard, A., Lewis, C. M., Yoshida, K., et al. Field pathogenomics reveals the emergence of a diverse wheat yellow rust population. *Genome Biology*, 16(1):1–15, 2015. ISSN 1465-6906. doi: 10.1186/s13059-015-0590-8.
- Hutchison, C. a. DNA sequencing: bench to bedside and beyond. *Nucleic acids research*, 35(18):6227–37, January 2007. ISSN 1362-4962. doi: 10.1093/nar/gkm688.
- Illumina. *CASAVA v1.8.2 User guide*. Revc edition, 2011. URL http://support.illumina.com/sequencing/sequencing_software/casava/documentation.ilmn.
- James, G. V., Patel, V., Nordström, K. J., et al. User guide for mapping-by-sequencing in Arabidopsis. *Genome biology*, 14(6):R61, June 2013. ISSN 1465-6914. doi: 10.1186/gb-2013-14-6-r61.
- Jupe, F., Witek, K., Verweij, W., et al. Resistance gene enrichment sequencing (RenSeq) enables reannotation of the NB-LRR gene family from sequenced plant genomes and rapid mapping of resistance loci in segregating populations. *The Plant journal : for cell and molecular biology*, 76(3):530–544, November 2013. ISSN 1365-313X. doi: 10.1111/tpj.12307.
- Katoh, K. and Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molec-*

- ular biology and evolution, 30(4):772–80, April 2013. ISSN 1537-1719. doi: 10.1093/molbev/mst010.
- Kent, W. J. BLAT—The BLAST-Like Alignment Tool. *Genome Research*, 12(4):656–664, March 2002. ISSN 1088-9051. doi: 10.1101/gr.229202.
- King, R., Bird, N., Ramirez-Gonzalez, R., et al. Mutation scanning in wheat by exon capture and next-generation sequencing. *PLoS ONE*, 10(9):1–18, 09 2015. doi: 10.1371/journal.pone.0137549.
- Krasileva, K., Vasquez-Gross, H., Howell1, T., et al. Uncovering hidden variation in young polyploid wheat genomes. submitted 2016.
- Krasileva, K. V., Buffalo, V., Bailey, P., et al. Separating homeologs by phasing in the tetraploid wheat transcriptome. *Genome biology*, 14(6):R66, June 2013. ISSN 1465-6914. doi: 10.1186/gb-2013-14-6-r66.
- Lander, E. S., Linton, L. M., Birren, B., et al. Initial sequencing and analysis of the human genome. *Nature*, 409(6822):860–921, February 2001. ISSN 0028-0836. doi: 10.1038/35057062.
- LGC Genomics. <http://www.lgcgroup.com/services/genotyping/>, 2013. URL <http://www.lgcgroup.com/services/genotyping/>.
- Li, H. and Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)*, 25(14):1754–60, July 2009. ISSN 1367-4811. doi: 10.1093/bioinformatics/btp324.
- Li, H., Handsaker, B., Wysoker, A., et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*, 25(16):2078–9, August 2009. ISSN 1367-4811. doi: 10.1093/bioinformatics/btp352.
- Liu, Y. and Schmidt, B. Long read alignment based on maximal exact match seeds. *Bioinformatics (Oxford, England)*, 28(18):i318–i324, September 2012. ISSN 1367-4811. doi: 10.1093/bioinformatics/bts414.
- Ma, J., Stiller, J., Zheng, Z., et al. A high-throughput pipeline for detecting locus-specific polymorphism in hexaploid wheat (*triticum aestivum* l.). *Plant Methods*, 11(1), aug 2015. doi: 10.1186/s13007-015-0082-6.

- MAS Wheat. Mas wheat transcriptome. supplemental file 17., 2013. URL <http://maswheat.ucdavis.edu/Transcriptome/index.htm>.
- Mayer, K. F. X., Rogers, J., Dole el, J., et al. A chromosome-based draft sequence of the hexaploid bread wheat (*Triticum aestivum*) genome. *Science*, 345(6194):1251788–1251788, July 2014. ISSN 0036-8075. doi: 10.1126/science.1251788.
- McIntosh, R., Wellings, C.R., and Park, R.F. *Wheat Rusts: an Atlas of Resistance Genes*. CSIRO Publishing, East Melbourne, Victoria, Australia, 1995.
- Metzker, M. L. Sequencing technologies - the next generation. *Nature reviews. Genetics*, 11(1):31–46, January 2010. ISSN 1471-0064. doi: 10.1038/nrg2626.
- Michelmore, R. W., Paran, I., and Kesseli, R. V. Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proceedings of the National Academy of Sciences*, 88(21):9828–9832, November 1991. ISSN 0027-8424. doi: 10.1073/pnas.88.21.9828.
- Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L., and Wold, B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nature methods*, 5(7):621–8, July 2008. ISSN 1548-7105. doi: 10.1038/nmeth.1226.
- Murphy, L. R., Santra, D., Kidwell, K., et al. Linkage Maps of Wheat Stripe Rust Resistance Genes and for Use in Marker-Assisted Selection. *Crop Science*, 49(5):1786, 2009. ISSN 1435-0653. doi: 10.2135/cropsci2008.10.0621.
- Myllykangas, S., Buenrostro, J., and Ji, H. P. *Bioinformatics for High Throughput Sequencing*. Springer New York, New York, NY, 2012. ISBN 978-1-4614-0781-2. doi: 10.1007/978-1-4614-0782-9. URL <http://www.springerlink.com/index/10.1007/978-1-4614-0782-9>.
- Needleman, S. B. and Wunsch, C. D. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *Journal of Molecular Biology*, 48(3):443–453, March 1970. ISSN 00222836. doi: 10.1016/0022-2836(70)90057-4.

- Ng, S. B., Turner, E. H., Robertson, P. D., et al. Targeted capture and massively parallel sequencing of 12 human exomes. *Nature*, 461(7261): 272–276, Sep 2009. ISSN 0028-0836. doi: 10.1038/nature08250.
- Peng, J., Fahima, T., Röder, M., and Huang, Q. High-density molecular map of chromosome region harboring stripe-rust resistance genes YrH52 and Yr15 derived from wild emmer wheat, *Triticum dicoccoides*. *Genetica*, pages 199–210, 2000.
- Pickrell, J. K., Marioni, J. C., Pai, A. A., et al. Understanding mechanisms underlying human gene expression variation with rna sequencing. *Nature*, 464(7289):768–772, Apr 2010. ISSN 0028-0836. doi: 10.1038/nature08872.
- Pontius, J., Wagner, L., and Schuler, G. UniGene: A Unified View of the Transcriptome. In *The NCBI Handbook [Internet]*, chapter 21. National Center for Biotechnology Information (US), October 2002. URL <http://www.ncbi.nlm.nih.gov/books/NBK21083/>.
- Pozniak, C. J. IWGSC whole genome shotgun sequencing of chinese spring: Towards a reference sequence of wheat, 1 2016. URL <https://pag.confex.com/pag/xxiv/webprogram/Paper22378.html>.
- Ramirez-Gonzalez, R. H., Uauy, C., and Caccamo, M. PolyMarker: A fast polyploid primer design pipeline. *Bioinformatics*, pages 2–3, 2015a. ISSN 1367-4803. doi: 10.1093/bioinformatics/btv069.
- Ramirez-Gonzalez, R. H., Bonnal, R., Caccamo, M., and Maclean, D. Bio-samtools: Ruby bindings for SAMtools, a library for accessing BAM files containing high-throughput sequence alignments. *Source code for biology and medicine*, 7(1):6, January 2012. ISSN 1751-0473. doi: 10.1186/1751-0473-7-6.
- Ramirez-Gonzalez, R. H., Segovia, V., Bird, N., Caccamo, M., and Uauy, C. *Next Generation Sequencing Enabled Genetics in Hexaploid Wheat*, pages 201–209. Springer Japan, Tokyo, 2015b. ISBN 978-4-431-55675-6. doi: 10.1007/978-4-431-55675-6_22. URL http://dx.doi.org/10.1007/978-4-431-55675-6_22.
- Ramirez-Gonzalez, R. H., Segovia, V., Bird, N., et al. Rna-seq bulked segregant analysis enables the identification of high-resolution genetic

- markers for breeding in hexaploid wheat. *Plant Biotechnology Journal*, 13(5):613–624, 2015c. ISSN 1467-7652. doi: 10.1111/pbi.12281.
- Randhawa, H. S., Mutti, J. S., Kidwell, K., et al. Rapid and targeted introgression of genes into popular wheat cultivars using marker-assisted background selection. *PloS one*, 4(6):e5752, January 2009. ISSN 1932-6203. doi: 10.1371/journal.pone.0005752.
- Rozen, S. and Skaletsky, H. Primer3 on the WWW for general users and for biologist programmers. *Methods in molecular biology (Clifton, N.J.)*, 132:365–86, January 2000. ISSN 1064-3745.
- Schneeberger, K. and Weigel, D. Fast-forward genetics enabled by new sequencing technologies. *Trends in Plant Science*, 16(5):282–288, 2011.
- Schneeberger, K., Ossowski, S., Lanz, C., et al. Shoremap: simultaneous mapping and mutation identification by deep sequencing. *Nat Meth*, 6(8):550–551, Aug 2009. ISSN 1548-7091. doi: 10.1038/nmeth0809-550.
- Shendure, J. and Ji, H. Next-generation DNA sequencing. *Nature biotechnology*, 26(10):1135–45, October 2008. ISSN 1546-1696. doi: 10.1038/nbt1486.
- Slater, G. S. C. and Birney, E. Automated generation of heuristics for biological sequence comparison. *BMC bioinformatics*, 6:31, January 2005. ISSN 1471-2105. doi: 10.1186/1471-2105-6-31.
- Smith, T. and Waterman, M. Identification of common molecular subsequences. *Journal of Molecular Biology*, 147(1):195–197, March 1981. ISSN 00222836. doi: 10.1016/0022-2836(81)90087-5.
- Sun, G. L., Fahima, T., Korol, A. B., et al. Identification of molecular markers linked to the Yr15 stripe rust resistance gene of wheat originated in wild emmer wheat, *Triticum dicoccoides*. *TAG Theoretical and Applied Genetics*, 95(4):622–628, September 1997. ISSN 0040-5752. doi: 10.1007/s001220050604.
- Takagi, H., Abe, A., Yoshida, K., et al. QTL-seq: rapid mapping of quantitative trait loci in rice by whole genome resequencing of DNA from two bulked populations. *The Plant journal : for cell and molecular biology*, 74(1):174–83, May 2013a. ISSN 1365-313X. doi: 10.1111/tpj.12105.

- Takagi, H., Uemura, A., Yaegashi, H., et al. MutMap-Gap: whole-genome resequencing of mutant F2 progeny bulk combined with de novo assembly of gap regions identifies the rice blast resistance gene Pii. *The New phytologist*, 200(1):276–83, October 2013b. ISSN 1469-8137. doi: 10.1111/nph.12369.
- Tang, H., Zhang, X., Miao, C., et al. Allmaps: robust scaffold ordering based on multiple maps. *Genome Biology*, 16(1):1–15, 2015. ISSN 1465-6906. doi: 10.1186/s13059-014-0573-1.
- Trick, M., Adamski, N., Mugford, S. G., et al. Combining SNP discovery from next-generation sequencing data with bulked segregant analysis (BSA) to fine-map genes in polyploid wheat. *BMC plant biology*, 12(1):14, January 2012. ISSN 1471-2229. doi: 10.1186/1471-2229-12-14.
- Van Ooijen, J. and Jansen, J. *Estimation of recombination frequencies; Genetic Mapping in Experimental Populations*, pages 73–133. Cambridge University Press, Cambridge, 2013. ISBN 978-1-107-0132-16.
- Wang, S., Wong, D., Forrest, K., et al. Characterization of polyploid wheat genomic diversity using a high-density 90 000 single nucleotide polymorphism array. *Plant biotechnology journal*, 12(6):787–796, March 2014. ISSN 1467-7652. doi: 10.1111/pbi.12183.
- Wang, Y., Tiwari, V. K., Rawat, N., et al. GSP: a web-based platform for designing genome-specific primers in polyploids. *Bioinformatics*, page btw134, mar 2016. doi: 10.1093/bioinformatics/btw134.
- Wellings, C. and McIntosh, R. A. Host-pathogen studies of wheat stripe rust in australia. In Slinkard, A., editor, *Proceedings 9th International Wheat Genetics Symposium*, pages 336–338. University of Saskatchewan, Saskatoon, SK, Canada, 1998.
- Wilkinson, P. a., Winfield, M. O., Barker, G. L. a., et al. CerealsDB 2.0: an integrated resource for plant breeders and scientists. *BMC bioinformatics*, 13(1):219, January 2012. ISSN 1471-2105. doi: 10.1186/1471-2105-13-219.
- Winfield, M. O., Wilkinson, P. A., Allen, A. M., et al. Targeted re-sequencing of the allohexaploid wheat exome. *Plant Biotechnology Journal*, 10(6):733–742, 2012. ISSN 1467-7652. doi: 10.1111/j.1467-7652.2012.00713.x.