

Lecture 16: Eigenspaces and Diagonalization

MATH203 Applied Linear Algebra

Updated May 6, 2026

What Does P_i Project Onto?

Standing Assumption

Throughout this lecture:

$$\det(tI - A) = (t - \lambda_1)(t - \lambda_2) \cdots (t - \lambda_m), \quad \lambda_i \text{ all distinct}$$

Same assumption as Lecture 15.

We have the spectral decomposition:

$$A = \lambda_1 P_1 + \lambda_2 P_2 + \cdots + \lambda_m P_m$$

where $P_1, \dots, P_m$ are compatible projections that partition I .

Today's Question

$$A = \lambda_1 P_1 + \lambda_2 P_2 + \cdots + \lambda_m P_m$$

$$\underbrace{\mathbf{v}}_{\text{any vector}} = \underbrace{P_1 \mathbf{v}}_{\in \text{Col}(P_1)} + \underbrace{P_2 \mathbf{v}}_{\in \text{Col}(P_2)} + \cdots + \underbrace{P_m \mathbf{v}}_{\in \text{Col}(P_m)}$$

What is $\text{Col}(P_i)$?

What does A do to vectors in $\text{Col}(P_i)$?

A New Example

$$\underbrace{A = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix}}_{\text{our running example}}, \quad \det(tI - A) = (t - 1)(t - 3)$$

Eigenvalues: $\lambda_1 = 1$, $\lambda_2 = 3$.

A New Example

$$\underbrace{A = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix}}_{\text{our running example}}, \quad \det(tI - A) = (t - 1)(t - 3)$$

Eigenvalues: $\lambda_1 = 1$, $\lambda_2 = 3$.

Spectral projections (Lagrange formula from L15):

$$P_1 = \frac{A - 3I}{1 - 3} = \begin{pmatrix} 2 & -1 \\ 2 & -1 \end{pmatrix}, \quad P_3 = \frac{A - I}{3 - 1} = \begin{pmatrix} -1 & 1 \\ -2 & 2 \end{pmatrix}$$

Value tables (from L15):

	$t = 1$	$t = 3$
A	1	3
P_1	1	0
P_3	0	1

$$A = \underbrace{1}_{\lambda_1} \cdot P_1 + \underbrace{3}_{\lambda_2} \cdot P_3$$

Goal: Discover what $\text{Col}(P_i)$ is.

1. Pick a test vector $\mathbf{v}$, split it as $\mathbf{v} = P_1\mathbf{v} + P_3\mathbf{v}$.
2. Apply A to each piece — see what happens.
3. Explain **why** using value tables.
4. Name the subspace $\text{Col}(P_i)$.

What Does P_i Do to a Vector?

Pick $\mathbf{v} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$. Apply each projection:

$$P_1 \mathbf{v} = \begin{pmatrix} 2 & -1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$P_3 \mathbf{v} = \begin{pmatrix} -1 & 1 \\ -2 & 2 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

What Does P_i Do to a Vector?

Pick $\mathbf{v} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$. Apply each projection:


$$P_1 \mathbf{v} = \begin{pmatrix} 2 & -1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$P_3 \mathbf{v} = \begin{pmatrix} -1 & 1 \\ -2 & 2 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$\text{Check: } \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \mathbf{v} \quad \checkmark$$

Every Vector Splits Into Eigenspace Pieces

$$\mathbf{v} = P_1 \mathbf{v} + P_3 \mathbf{v} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} \quad \checkmark$$



$P_1 + P_3 = I$: every vector decomposes into eigenspace pieces (parallelogram law).

What Does A Do to Each Piece?

Apply A to the first piece:

$$A \cdot P_1 \mathbf{v} = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \mathbf{1} \cdot P_1 \mathbf{v}$$

What Does A Do to Each Piece?

Apply A to the **first piece**:

$$A \cdot P_1 \mathbf{v} = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \mathbf{1} \cdot P_1 \mathbf{v}$$

Apply A to the **second piece**:

$$A \cdot P_3 \mathbf{v} = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \end{pmatrix} = \mathbf{3} \cdot P_3 \mathbf{v}$$

What Does A Do to Each Piece?

Apply A to the **first piece**:

$$A \cdot P_1 \mathbf{v} = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 1 \cdot P_1 \mathbf{v}$$

Apply A to the **second piece**:


$$A \cdot P_3 \mathbf{v} = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \end{pmatrix} = 3 \cdot P_3 \mathbf{v}$$

A **scales each piece by its eigenvalue**:

$$\lambda_1 = 1 \text{ and } \lambda_2 = 3.$$

A Scales Each Piece by Its Eigenvalue

$$A\mathbf{v} = A(P_1\mathbf{v} + P_3\mathbf{v}) = \underbrace{1}_{\lambda_1} \cdot P_1\mathbf{v} + \underbrace{3}_{\lambda_2} \cdot P_3\mathbf{v}$$



A doesn't rotate or mix the pieces — it **only scales** each one along its eigenspace.

Why? — Value Table Proof

Why does A scale the P_1 -piece by λ_1 ?

Multiply value tables entry-wise:

	$t = 1$	$t = 3$
A	1	3
$\times P_1$	1	0
AP_1	1	0

Why? — Value Table Proof

Why does A scale the P_1 -piece by λ_1 ?

Multiply value tables entry-wise:

	$t = 1$	$t = 3$
A	1	3
$\times P_1$	1	0
AP_1	1	0

Value table of $\lambda_1 P_1 = 1 \cdot P_1$:

	$t = 1$	$t = 3$
$1 \cdot P_1$	1	0

Same value table $\implies$ **same matrix**: $AP_1 = \lambda_1 P_1 = 1 \cdot P_1$.

Why? — Value Table Proof

Why does A scale the P_1 -piece by λ_1 ?

Multiply value tables entry-wise:

	$t = 1$	$t = 3$
A	1	3
$\times P_1$	1	0
AP_1	1	0

Value table of $\lambda_1 P_1 = 1 \cdot P_1$:

	$t = 1$	$t = 3$
$1 \cdot P_1$	1	0

Same value table $\implies$ **same matrix**: $AP_1 = \lambda_1 P_1 = 1 \cdot P_1$.

Similarly for P_3 :

	$t = 1$	$t = 3$
A	1	3
$\times P_3$	0	1
AP_3	0	3

= value table of $3P_3$ ✓

A Commutes With Each P_i

What about P_1A ? **Multiply** value tables in the other order:

	$t = 1$	$t = 3$
P_1	1	0
$\times A$	1	3
P_1A	1	0

Same value table as AP_1 ! So $P_1A = AP_1 = \lambda_1 P_1$.

A Commutes With Each P_i

What about P_1A ? **Multiply** value tables in the other order:

	$t = 1$	$t = 3$
P_1	1	0
$\times A$	1	3
P_1A	1	0

Same value table as AP_1 ! So $P_1A = AP_1 = \lambda_1 P_1$.

Value table multiplication is **entry-wise**: order doesn't matter.

$$AP_i = P_iA = \lambda_i P_i \quad \text{for each } i$$

Answer: $\text{Col}(P_i)$ is the **Eigenspace**

Definition (Eigenspace)

The **eigenspace** of A for eigenvalue λ_i is

$$E_{\lambda_i} = \{ \mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \lambda_i \mathbf{x} \}$$

Answer: $\text{Col}(P_i)$ is the Eigenspace

Definition (Eigenspace)

The eigenspace of A for eigenvalue λ_i is

$$E_{\lambda_i} = \{ \mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \lambda_i \mathbf{x} \}$$

If $\mathbf{x} \in \text{Col}(P_i)$, then P_i fixes it: $P_i \mathbf{x} = \mathbf{x}$.

$$A\mathbf{x} = A \underbrace{P_i \mathbf{x}}_{=\mathbf{x}} = \underbrace{AP_i}_{=\lambda_i P_i} \mathbf{x} = \lambda_i \underbrace{P_i \mathbf{x}}_{=\mathbf{x}} = \lambda_i \mathbf{x} \quad \implies \quad \text{Col}(P_i) \subseteq E_{\lambda_i}$$

Answer: $\text{Col}(P_i)$ is the Eigenspace

Definition (Eigenspace)

The **eigenspace** of A for eigenvalue λ_i is

$$E_{\lambda_i} = \{ \mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \lambda_i \mathbf{x} \}$$

If $\mathbf{x} \in \text{Col}(P_i)$, then P_i fixes it: $P_i \mathbf{x} = \mathbf{x}$.

$$A\mathbf{x} = A \underbrace{P_i \mathbf{x}}_{=\mathbf{x}} = \underbrace{AP_i}_{=\lambda_i P_i} \mathbf{x} = \lambda_i \underbrace{P_i \mathbf{x}}_{=\mathbf{x}} = \lambda_i \mathbf{x} \quad \implies \quad \text{Col}(P_i) \subseteq E_{\lambda_i}$$

Conversely: P_i is a polynomial of A (Lagrange formula). If $A\mathbf{x} = \lambda_i \mathbf{x}$, then $f(A)\mathbf{x} = f(\lambda_i)\mathbf{x}$ for any polynomial f . Since P_i evaluates to 1 at λ_i :

$$P_i \mathbf{x} = 1 \cdot \mathbf{x} = \mathbf{x} \quad \implies \quad E_{\lambda_i} \subseteq \text{Col}(P_i)$$

$$\boxed{E_{\lambda_i} = \text{Col}(P_i)}$$

Check on Our Example

$$E_1 = \text{Col}(P_1) = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}. \quad A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \mathbf{1} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \checkmark$$

$$E_3 = \text{Col}(P_3) = \text{span} \left\{ \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right\}. \quad A \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \end{pmatrix} = \mathbf{3} \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad \checkmark$$

P_i projects onto E_{λ_i} :
the subspace where A acts as **scalar multiplication by**
 λ_i .

Compatible Projections Revisited

What We Want to Show

Two main points in this section:

1. Any vector $\mathbf{v}$ can be **decomposed into eigenvectors** — one from each eigenspace E_{λ_i} .
2. Eigenvectors with **different eigenvalues** are **linearly independent**: the decomposition is unique.

The key: $P_1, \dots, P_m$ are **compatible projections** (Lecture 8–9).

Recall: Where Did P_1, P_3 Come From?

In Lecture 15, we plugged A into the **Lagrange interpolation polynomials**:

$$P_1 = \frac{A - 3I}{1 - 3}, \quad P_3 = \frac{A - I}{3 - 1}$$

This gave the **spectral decomposition**: $A = 1 \cdot P_1 + 3 \cdot P_3$.

Their **value tables** (from Lecture 15):

	$t = 1$	$t = 3$
P_1	1	0
P_3	0	1

Recall: Where Did P_1, P_3 Come From?

In Lecture 15, we plugged A into the **Lagrange interpolation polynomials**:

$$P_1 = \frac{A - 3I}{1 - 3}, \quad P_3 = \frac{A - I}{3 - 1}$$

This gave the **spectral decomposition**: $A = 1 \cdot P_1 + 3 \cdot P_3$.

Their **value tables** (from Lecture 15):

	$t = 1$	$t = 3$
P_1	1	0
P_3	0	1

We **claimed** P_1, P_3 are compatible projections that partition I .

Let's verify this using value tables — the tool we trust from Lecture 15.

Are P_1, P_3 Projections?

Check with value tables — **square each entry**:

	$t = 1$	$t = 3$
P_1	1	0
P_1^2	$1^2 = \mathbf{1}$	$0^2 = \mathbf{0}$

	$t = 1$	$t = 3$
P_3	0	1
P_3^2	$0^2 = \mathbf{0}$	$1^2 = \mathbf{1}$

Same value tables $\implies P_1^2 = P_1$ and $P_3^2 = P_3$. Both are **projections**.

✓

Compatible? Partition of /?

Compatible? Multiply value tables entry-wise:

	$t = 1$	$t = 3$
P_1	1	0
$\times P_3$	0	1
$P_1 P_3$	0	0

Value table all zeros $\implies P_1 P_3 = 0$. **Compatible!** ✓

Compatible? Partition of I ?

Compatible? Multiply value tables entry-wise:

	$t = 1$	$t = 3$
P_1	1	0
$\times P_3$	0	1
$P_1 P_3$	0	0

Value table all zeros $\implies P_1 P_3 = 0$. **Compatible!** ✓


Partition of I ? Add value tables:

	$t = 1$	$t = 3$
P_1	1	0
$+ P_3$	0	1
$P_1 + P_3$	1	1

Value table = value table of I . $P_1 + P_3 = I$. ✓

Connection to Lecture 8: Compatible Projections


From Lecture 8: $P_1 P_3 = 0$ means $\text{Col}(P_3) \subset \text{Null}(P_1)$ and $\text{Col}(P_1) \subset \text{Null}(P_3)$.



Each eigenspace lives inside the null space of the **other** projection.

Connection to Lecture 8: Compatible Projections

From Lecture 8: $P_1 P_3 = 0$ means $\text{Col}(P_3) \subset \text{Null}(P_1)$ and $\text{Col}(P_1) \subset \text{Null}(P_3)$.



Each eigenspace lives inside the null space of the **other** projection.

The Lagrange formula from L15 **automatically**
produces compatible projections.

No extra verification needed — it's built into the construction.

Eigenvectors With Different Eigenvalues Are Independent

Recall from Lecture 9 (Property 2, Independent Decompositions):

For a compatible family $\{P_1, \dots, P_m\}$, the nonzero vectors among $\{P_1\mathbf{v}, \dots, P_m\mathbf{v}\}$ are **linearly independent**.

Eigenvectors With Different Eigenvalues Are Independent

Recall from Lecture 9 (Property 2, Independent Decompositions):

For a compatible family $\{P_1, \dots, P_m\}$, the nonzero vectors among $\{P_1\mathbf{v}, \dots, P_m\mathbf{v}\}$ are **linearly independent**.

Since the spectral projections $P_1, \dots, P_m$ are compatible:

Corollary

Let $\mathbf{v}_1, \dots, \mathbf{v}_m$ be eigenvectors of A with **distinct** eigenvalues $\lambda_1, \dots, \lambda_m$.
Then $\mathbf{v}_1, \dots, \mathbf{v}_m$ are **linearly independent**.

Each $\mathbf{v}_i \in E_{\lambda_i} = \text{Col}(P_i)$, so $P_i\mathbf{v}_i = \mathbf{v}_i \neq 0$ and $P_j\mathbf{v}_i = 0$ for $j \neq i$.

This is a direct consequence of Lecture 9.

Unique Decomposition (Lecture 9 Callback)

From Lecture 9: compatible projections that partition I give a **unique decomposition** of every vector.

Since $P_1 + P_3 = I$, every $\mathbf{v} \in \mathbb{R}^n$ decomposes as:

$$\mathbf{v} = \underbrace{P_1 \mathbf{v}}_{\in E_1} + \underbrace{P_3 \mathbf{v}}_{\in E_3}$$

and by the corollary above, this decomposition is **unique**.

The eigenspaces $E_{\lambda_1}, \dots, E_{\lambda_m}$ together **span** all of $\mathbb{R}^n$.

Every vector decomposes **uniquely** into eigenvector pieces.

(This is the geometric meaning of $P_1 + P_2 + \dots + P_m = I$.)

Cross-Filling the Spectral Projections

A New Example — Repeated Eigenvalue

$$B = \underbrace{\begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix}}_{\text{eigenvalues } 2,2,5}, \quad \det(tI - B) = (t - 2)^2(t - 5)$$

Two **distinct** eigenvalues: $\lambda_1 = 2$, $\lambda_2 = 5$.

A New Example — Repeated Eigenvalue

$$B = \underbrace{\begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix}}_{\text{eigenvalues } 2,2,5}, \quad \det(tI - B) = (t-2)^2(t-5)$$

Two **distinct** eigenvalues: $\lambda_1 = 2$, $\lambda_2 = 5$.

Spectral projections (Lagrange formula):

$$P_5 = \frac{B - 2I}{5 - 2} = \frac{1}{3} \begin{pmatrix} 0 & 0 & 3 \\ 0 & 0 & 3 \\ 0 & 0 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$P_2 = I - P_5 = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

	$t = 2$	$t = 5$
B	2	5
P_2	1	0
P_5	0	1

P_2 Has Rank 2 — A Plane

$$P_2 = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{rank} = 2$$

$E_2 = \text{Col}(P_2)$ is a **plane** in $\mathbb{R}^3$, not just a line.

$$P_5 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{rank} = 1$$

$E_5 = \text{Col}(P_5)$ is a **line** in $\mathbb{R}^3$.

P_2 Has Rank 2 — A Plane

$$P_2 = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{rank} = 2$$

$E_2 = \text{Col}(P_2)$ is a **plane** in $\mathbb{R}^3$, not just a line.

$$P_5 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{rank} = 1$$

$E_5 = \text{Col}(P_5)$ is a **line** in $\mathbb{R}^3$.

To find eigenvectors, we **cross-fill** each projection (Lecture 8).

Cross-Fill P_2 (Rank 2)

$$\underbrace{P_2}_{\text{rank 2}} = \underbrace{\begin{pmatrix} \color{red}{1} & \color{blue}{0} & \color{blue}{-1} \\ \color{blue}{0} & 0 & 0 \\ \color{blue}{0} & 0 & 0 \end{pmatrix}}_{\substack{R_1 \\ \text{rank 1}}} + \underbrace{\begin{pmatrix} 0 & 0 & 0 \\ 0 & \color{red}{1} & \color{blue}{-1} \\ 0 & \color{blue}{0} & 0 \end{pmatrix}}_{\substack{R_2 \\ \text{rank 1}}}$$

Cross-Fill P_2 (Rank 2)

$$\underbrace{P_2}_{\text{rank 2}} = \underbrace{\begin{pmatrix} \color{red}{1} & \color{blue}{0} & \color{blue}{-1} \\ \color{blue}{0} & 0 & 0 \\ \color{blue}{0} & 0 & 0 \end{pmatrix}}_{\substack{R_1 \\ \text{rank 1}}} + \underbrace{\begin{pmatrix} 0 & 0 & 0 \\ 0 & \color{red}{1} & \color{blue}{-1} \\ 0 & \color{blue}{0} & 0 \end{pmatrix}}_{\substack{R_2 \\ \text{rank 1}}}$$

Two-factor form (stack columns and rows):

$$P_2 = \underbrace{\begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}}_{U_2} \underbrace{\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}}_{V_2}, \quad V_2 U_2 = I_2 \checkmark$$

Cross-Fill P_5 (Rank 1) and Eigenvectors

$$\underbrace{P_5}_{\text{rank 1}} = \underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}}_{U_5} \underbrace{\begin{pmatrix} 0 & 0 & 1 \end{pmatrix}}_{V_5}, \quad V_5 U_5 = 1 \checkmark$$

Cross-Fill P_5 (Rank 1) and Eigenvectors

$$\underbrace{P_5}_{\text{rank 1}} = \underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}}_{U_5} \underbrace{\begin{pmatrix} 0 & 0 & 1 \end{pmatrix}}_{V_5}, \quad V_5 U_5 = 1 \checkmark$$

Columns of U_i = right eigenvectors ($B\mathbf{u} = \lambda\mathbf{u}$):

$$B \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \checkmark \quad B \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = 2 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \checkmark \quad B \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 5 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \checkmark$$

Cross-Fill P_5 (Rank 1) and Eigenvectors

$$\underbrace{P_5}_{\text{rank 1}} = \underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}}_{U_5} \underbrace{\begin{pmatrix} 0 & 0 & 1 \end{pmatrix}}_{V_5}, \quad V_5 U_5 = 1 \checkmark$$

Columns of U_i = right eigenvectors ($B\mathbf{u} = \lambda\mathbf{u}$):

$$B \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \mathbf{2} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \checkmark \quad B \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \mathbf{2} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \checkmark \quad B \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \mathbf{5} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \checkmark$$

Rows of V_i = left eigenvectors ($\mathbf{v}^T B = \lambda \mathbf{v}^T$):

$$\begin{pmatrix} 1 & 0 & -1 \end{pmatrix} B = \mathbf{2} \begin{pmatrix} 1 & 0 & -1 \end{pmatrix} \checkmark \quad \begin{pmatrix} 0 & 1 & -1 \end{pmatrix} B = \mathbf{2} \begin{pmatrix} 0 & 1 & -1 \end{pmatrix} \checkmark \\ \begin{pmatrix} 0 & 0 & 1 \end{pmatrix} B = \mathbf{5} \begin{pmatrix} 0 & 0 & 1 \end{pmatrix} \checkmark$$

Cross-Filling Produces Rank-1 Projections (L08, L09)

From Lecture 9: cross-filling a projection gives **rank-1 projections** that form a compatible family.

$$P_2 = \underbrace{\begin{pmatrix} \mathbf{1} & \mathbf{0} & \mathbf{-1} \\ \mathbf{0} & 0 & 0 \\ \mathbf{0} & 0 & 0 \end{pmatrix}}_{\substack{R_1 \\ \text{rank-1 projection}}} + \underbrace{\begin{pmatrix} 0 & 0 & 0 \\ 0 & \mathbf{1} & \mathbf{-1} \\ 0 & \mathbf{0} & 0 \end{pmatrix}}_{\substack{R_2 \\ \text{rank-1 projection}}}$$

R_1, R_2 compatible: $R_1 R_2 = 0$, $R_1^2 = R_1$, $R_2^2 = R_2$. P_5 already rank 1.

Cross-Filling Produces Rank-1 Projections (L08, L09)

From Lecture 9: cross-filling a projection gives **rank-1 projections** that form a compatible family.

$$P_2 = \underbrace{\begin{pmatrix} \mathbf{1} & \mathbf{0} & \mathbf{-1} \\ \mathbf{0} & 0 & 0 \\ \mathbf{0} & 0 & 0 \end{pmatrix}}_{\substack{R_1 \\ \text{rank-1 projection}}} + \underbrace{\begin{pmatrix} 0 & 0 & 0 \\ 0 & \mathbf{1} & \mathbf{-1} \\ 0 & \mathbf{0} & 0 \end{pmatrix}}_{\substack{R_2 \\ \text{rank-1 projection}}}$$

R_1, R_2 compatible: $R_1 R_2 = 0$, $R_1^2 = R_1$, $R_2^2 = R_2$. P_5 already rank 1.

Cross-eigenspace: $V_2 U_5 = 0$, $V_5 U_2 = 0$ (from $P_2 P_5 = 0$).

The Final Spectral Decomposition

$B = 2P_2 + 5P_5$. Replace $P_2 = R_1 + R_2$: now **every piece is rank 1**:

$$B = \underbrace{2 \cdot R_1}_{\text{rank 1}} + \underbrace{2 \cdot R_2}_{\text{rank 1}} + \underbrace{5 \cdot P_5}_{\text{rank 1}}$$

Each R_i is a rank-1 projection (L09), and the eigenvalue 2 appears **twice**.

The Final Spectral Decomposition

$B = 2P_2 + 5P_5$. Replace $P_2 = R_1 + R_2$: now **every piece is rank 1**:

$$B = \underbrace{2 \cdot R_1}_{\text{rank 1}} + \underbrace{2 \cdot R_2}_{\text{rank 1}} + \underbrace{5 \cdot P_5}_{\text{rank 1}}$$

Each R_i is a rank-1 projection (L09), and the eigenvalue 2 appears **twice**.

From Lecture 8: a sum of rank-1 matrices = a **product** $\longrightarrow$

Diagonalization: Stacking the Cross-Fillings

From Sum to Product

$$B = 2 \cdot R_1 + 2 \cdot R_2 + 5 \cdot P_5$$

Write with **general variable** λ replacing the eigenvalues:

$$\begin{aligned} & \lambda \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 \end{pmatrix} + \lambda \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & -1 \end{pmatrix} + \lambda \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \end{pmatrix} \\ &= \underbrace{\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}}_P \underbrace{\begin{pmatrix} \lambda & & \\ & \lambda & \\ & & \lambda \end{pmatrix}}_{D(\lambda)} \underbrace{\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}}_Q \end{aligned}$$

Same P , same Q — only the **diagonal changes**.

The General Formula: $g(B) = P g(D) Q$

For **any** function g , replace each diagonal entry λ_i by $g(\lambda_i)$:

$$g(B) = P \begin{pmatrix} g(\lambda_1) & & \\ & g(\lambda_2) & \\ & & g(\lambda_3) \end{pmatrix} Q$$

The General Formula: $g(B) = P g(D) Q$

For **any** function g , replace each diagonal entry λ_i by $g(\lambda_i)$:

$$g(B) = P \begin{pmatrix} g(\lambda_1) & & \\ & g(\lambda_2) & \\ & & g(\lambda_3) \end{pmatrix} Q$$

Set $g(\lambda) = 1$ for all λ :

$$\underbrace{1 \cdot R_1 + 1 \cdot R_2 + 1 \cdot P_5}_{=P_2+P_5=I \text{ (value table from §2)}} = P \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} Q = PQ$$

$$\boxed{PQ = I, \quad \text{so} \quad Q = P^{-1}}$$

The General Formula: $g(B) = P g(D) Q$

For **any** function g , replace each diagonal entry λ_i by $g(\lambda_i)$:

$$g(B) = P \begin{pmatrix} g(\lambda_1) & & \\ & g(\lambda_2) & \\ & & g(\lambda_3) \end{pmatrix} Q$$

Set $g(\lambda) = 1$ for all λ :

$$\underbrace{1 \cdot R_1 + 1 \cdot R_2 + 1 \cdot P_5}_{=P_2+P_5=I \text{ (value table from §2)}} = P \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} Q = PQ$$

$$\boxed{PQ = I, \quad \text{so} \quad Q = P^{-1}}$$

Set $g(\lambda) = \lambda$ (identity): $B = PDP^{-1}$. Set $g(\lambda) = \lambda^n$: $B^n = PD^nP^{-1}$.

Diagonalization: $B = PDP^{-1}$

$$\underbrace{\begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix}}_B = \underbrace{\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}}_P \underbrace{\begin{pmatrix} 2 & & \\ & 2 & \\ & & 5 \end{pmatrix}}_D \underbrace{\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}}_{P^{-1}}$$

Verify: $PD = \begin{pmatrix} 2 & 0 & 5 \\ 0 & 2 & 5 \\ 0 & 0 & 5 \end{pmatrix}, \quad PDP^{-1} = \begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix} = B \quad \checkmark$

Diagonalization: $B = PDP^{-1}$

$$\underbrace{\begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix}}_B = \underbrace{\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}}_P \underbrace{\begin{pmatrix} 2 & & \\ & 2 & \\ & & 5 \end{pmatrix}}_D \underbrace{\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}}_{P^{-1}}$$

Verify: $PD = \begin{pmatrix} 2 & 0 & 5 \\ 0 & 2 & 5 \\ 0 & 0 & 5 \end{pmatrix}, \quad PDP^{-1} = \begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix} = B \quad \checkmark$

Columns of P = right eigenvectors. Rows of P^{-1} = left eigenvectors.

Diagonal of D = eigenvalues (with repetition).

The Cancellation Trick

Why does $g(B) = P g(D) P^{-1}$?

$$B^2 = \overbrace{(PDP^{-1})}^B \overbrace{(PDP^{-1})}^B = PD \underbrace{P^{-1}P}_{=I} DP^{-1} = PD^2 P^{-1}$$

The $P^{-1}P$ in the middle **Cancels!**

The Cancellation Trick

Why does $g(B) = P g(D) P^{-1}$?

$$B^2 = \overbrace{(PDP^{-1})}^B \overbrace{(PDP^{-1})}^B = PD \underbrace{P^{-1}P}_{=I} DP^{-1} = PD^2P^{-1}$$

The $P^{-1}P$ in the middle **cancels**!

Powers:

$$B^n = PD^nP^{-1}, \quad D^n = \begin{pmatrix} 2^n & & \\ & 2^n & \\ & & 5^n \end{pmatrix}$$

Diagonal matrices are trivial to exponentiate.

Connection to Value Tables

Two notations for the same machine:

Spectral formula (L15): $g(B) = g(2) P_2 + g(5) P_5$

Diagonalization: $g(B) = P g(D) P^{-1}$

Connection to Value Tables

Two notations for the same machine:

Spectral formula (L15): $g(B) = g(2) P_2 + g(5) P_5$

Diagonalization: $g(B) = P g(D) P^{-1}$

	$t = 2$	$t = 5$
$g(B)$	$g(2)$	$g(5)$

The value table computes $g(D)$. The spectral projections are hidden inside P and P^{-1} .

Same machine, two notations.

Eigenspace Geometry: The Dominant Eigenvalue

A^n via Value Tables

Back to our 2×2 example: $A = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix}$, eigenvalues 1, 3.

	$t = 1$	$t = 3$
A	1	3
A^2	1	9
A^3	1	27
A^n	1	3^n

A^n via Value Tables

Back to our 2×2 example: $A = \begin{pmatrix} -1 & 2 \\ -4 & 5 \end{pmatrix}$, eigenvalues 1, 3.


	$t = 1$	$t = 3$
A	1	3
A^2	1	9
A^3	1	27
A^n	1	3^n

$$\text{For } \mathbf{v} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}: \quad A^n \mathbf{v} = \underbrace{1^n}_{=1} \underbrace{\begin{pmatrix} 1 \\ 1 \end{pmatrix}}_{P_1 \mathbf{v}} + \underbrace{3^n}_{\rightarrow \infty} \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{P_3 \mathbf{v}}$$

As $n \rightarrow \infty$: the E_3 piece grows like 3^n , the E_1 piece stays = 1.

The dominant eigenvalue wins: $A^n \mathbf{v}$ aligns with $E_3 = \text{span}\left\{\begin{pmatrix} 1 \\ 2 \end{pmatrix}\right\}$.

Visualizing A^n



Each step: the E_3 piece stretches $\times 3$, the E_1 piece is unchanged.

Eigenvalues tell you **how much** each direction stretches.

Eigenspaces tell you **which** directions.

Together, they completely describe what A does.

Our Restriction: The Minimal Polynomial

Eigenvalues Are Roots of Any Annihilating Polynomial

Theorem

If $p(A) = 0$ and λ is an eigenvalue of A , then $p(\lambda) = 0$.

Proof: Let $A\mathbf{x} = \lambda\mathbf{x}$ with $\mathbf{x} \neq \mathbf{0}$. Then:

$$\mathbf{0} = p(A)\mathbf{x} = p(\lambda)\mathbf{x}$$

Since $\mathbf{x} \neq \mathbf{0}$: $p(\lambda) = 0$.



Eigenvalues Are Roots of Any Annihilating Polynomial

Theorem

If $p(A) = 0$ and λ is an eigenvalue of A , then $p(\lambda) = 0$.

Proof: Let $A\mathbf{x} = \lambda\mathbf{x}$ with $\mathbf{x} \neq \mathbf{0}$. Then:

$$\mathbf{0} = p(A)\mathbf{x} = p(\lambda)\mathbf{x}$$

Since $\mathbf{x} \neq \mathbf{0}$: $p(\lambda) = 0$. □

Consequence: Every annihilating polynomial must have **all eigenvalues** as roots.

Our spectral decomposition uses an annihilating polynomial with **distinct linear factors**. If the characteristic polynomial $\det(tI - A)$ has repeated factors, we need a **simpler** annihilating polynomial.

A Repeated Root That Worked

B from §3 had $\det(tI - B) = (t - 2)^2(t - 5)$ — a **repeated** root!

Yet the spectral machinery worked perfectly:

- Two distinct projections P_2, P_5 ✓
- Compatible, partition of I ✓
- Cross-filling, eigenvectors ✓
- $B = PDP^{-1}$ ✓

A Repeated Root That Worked

B from §3 had $\det(tI - B) = (t - 2)^2(t - 5)$ — a **repeated** root!

Yet the spectral machinery worked perfectly:

- Two distinct projections P_2, P_5 ✓
- Compatible, partition of I ✓
- Cross-filling, eigenvectors ✓
- $B = PDP^{-1}$ ✓

Why? There exists a simpler annihilating polynomial with distinct factors.

The Minimal Polynomial

Definition (Minimal Polynomial)

The **minimal polynomial** $m(t)$ is the lowest-degree monic polynomial such that $m(A) = 0$.

It divides every annihilating polynomial (including $\det(tI - A)$).

The Minimal Polynomial

Definition (Minimal Polynomial)

The **minimal polynomial** $m(t)$ is the lowest-degree monic polynomial such that $m(A) = 0$.

It divides every annihilating polynomial (including $\det(tI - A)$).

Check on B :

$$(B - 2I)(B - 5I) = \begin{pmatrix} 0 & 0 & 3 \\ 0 & 0 & 3 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} -3 & 0 & 3 \\ 0 & -3 & 3 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

So $m(t) = (t - 2)(t - 5)$ — **distinct** linear factors. ✓

Another Fine Example: $A = 3I$

$$A = \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}, \quad \det(tI - A) = (t - 3)^2 \quad (\text{repeated!})$$

But $A - 3I = 0$ already, so $m(t) = t - 3$ (degree 1, one distinct factor).

Spectral decomposition: $A = 3 \cdot I$. Just one projection $P_1 = I$.

The whole space $\mathbb{R}^2$ is one eigenspace. Fine!

Beyond Our Assumption: The Shear

$$J = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}, \quad \det(tI - J) = (t - 2)^2$$

$$J - 2I = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq 0. \quad (J - 2I)^2 = 0.$$


So $m(t) = (t - 2)^2$ — **repeated** factor!

Beyond Our Assumption: The Shear

$$J = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}, \quad \det(tI - J) = (t - 2)^2$$

$$J - 2I = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq 0. \quad (J - 2I)^2 = 0.$$

So $m(t) = (t - 2)^2$ — **repeated** factor!



Vectors on E_2 scale by 2. Vectors off E_2 get **sheared** — not just scaled.

This situation needs **new tools** — a topic for a future lecture.

The Criterion for Diagonalizability

Theorem

A is **diagonalizable** if and only if the minimal polynomial $m(t)$ factors into **distinct linear factors**:

$$m(t) = (t - \lambda_1)(t - \lambda_2) \cdots (t - \lambda_k), \quad \lambda_i \text{ all distinct}$$

The Criterion for Diagonalizability

Theorem

A is **diagonalizable** if and only if the minimal polynomial $m(t)$ factors into **distinct linear factors**:

$$m(t) = (t - \lambda_1)(t - \lambda_2) \cdots (t - \lambda_k), \quad \lambda_i \text{ all distinct}$$

($\Leftarrow$): Distinct linear factors $\implies$ spectral decomposition works $\implies$ diagonalizable. (This is exactly what we did in §1–§4.)

($\Rightarrow$): If $A = PDP^{-1}$, then

$(A - \lambda_1 I) \cdots (A - \lambda_k I) = P(D - \lambda_1 I) \cdots (D - \lambda_k I)P^{-1} = 0$, because every diagonal entry of D is one of the λ_i 's.

The Real Restriction

	Char. poly	Min. poly	Diag.?
$B = \begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix}$	$(t-2)^2(t-5)$	$(t-2)(t-5)$	✓
$3I_2$	$(t-3)^2$	$(t-3)$	✓
$J = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$	$(t-2)^2$	$(t-2)^2$	✗

The characteristic polynomial CAN have repeated roots — that's fine.

The **minimal polynomial** is what matters:

A is diagonalizable $\iff m(t)$ has **distinct linear factors**.

Repeated factor in $m(t)$: needs new tools (future lecture).



Summary

1. $E_{\lambda_i} = \text{Col}(P_i)$: eigenspace = column space of spectral projection.
2. Compatible projections (value tables verify instantly): eigenvectors with different eigenvalues are independent; every vector decomposes uniquely.
3. Cross-fill $P_i = U_i V_i$: columns of U_i = right eigenvectors, rows of V_i = left eigenvectors. $V_i U_j = \delta_{ij} I$.
4. Stack: $A = PDP^{-1}$, $g(A) = Pg(D)P^{-1}$ reduces every computation to plugging eigenvalues into g .
5. The **minimal polynomial** is the real restriction: distinct linear factors $\implies$ diagonalizable.

We can compute A^n via eigenvalues. But what about e^A ?

$$e^A = \lim_{n \rightarrow \infty} \left(I + \frac{A}{n} \right)^n$$

Next: The exponential function, matrix exponential, and differential equations.