

Task 3

Task 3: Compute posterior summaries and 95% credible intervals of γ , the fixed effects associated with the covariates in the model. Using the estimated Bayesian model, answer the following questions: (1) are there seasonal differences in hurricane wind speeds, and (2) is there evidence to support the claim that hurricane wind speeds have been increasing over the years?

Import gamma from task 1 and 2.

```
library(MASS)
library(tidyverse)
```

```
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
## v dplyr      1.1.1      v readr      2.1.4
## v forcats    1.0.0      v stringr   1.5.0
## v ggplot2    3.4.2      v tibble    3.2.1
## v lubridate  1.9.2      v tidyr     1.3.0
## v purrr      1.0.1
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
## x dplyr::select() masks MASS::select()
## i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become errors
```


We are mainly interested in γ_{Season} which is about year change, and γ_{Active} which is about active season change compared to inactive season.

```
gamma_list <- read.csv("./data/gamma_list.csv")
summary(gamma_list, digit = 1) %>% knitr::kable(digits = 3, col.names = c("gamma_year", "gamma_active", "gamma_ET", "gamma_NR", "gamma_SS", "gamma_TS"))
```

gamma_year	gamma_active	gamma_ET	gamma_NR	gamma_SS	gamma_TS
Min. :-1e-02	Min. :-0.196	Min. :-0.177	Min. :-2e-01	Min. :-0.196	Min. :-0.194
1st Qu.: -2e-03	1st Qu.: -0.041	1st Qu.: -0.031	1st Qu.: -3e-02	1st Qu.: -0.029	1st Qu.: -0.027
Median :	Median	Median :	Median	Median :	Median :
2e-03	:-0.008	0.002	:-1e-03	0.004	0.007
Mean : 1e-04	Mean :-0.008	Mean : 0.003	Mean :-9e-04	Mean : 0.005	Mean : 0.007
3rd Qu.: 3e-03	3rd Qu.: 0.026	3rd Qu.: 0.037	3rd Qu.: 3e-02	3rd Qu.: 0.038	3rd Qu.: 0.040
Max. : 4e-03	Max. : 0.199	Max. : 0.181	Max. : 2e-01	Max. : 0.182	Max. : 0.176


See the time-series plot to make decision about burn in

```
plot(gamma_list[,1], type = "l", ylab = "gamma_year")
abline(v = 8000, col = "red", lty = "dashed")# e.g. plot gamma1 (year) seems not converge,
```



```
plot(gamma_list[,2], type = "l", ylab = "gamma_active") # plot gamma2(active season)
abline(v = 8000, col = "red", lty = "dashed")
```

seems cor



```
# set burn in = 8000
gamma_list = slice(gamma_list, 8001:10000)
```

```
# see the histogram of year, some kind of normal
quantiles <- apply(gamma_list, 2, quantile, c(0.025, 0.975)) # use 2.5 and 97.5 quantile to get 95% cre
quantiles %>% knitr::kable(digits = 3,col.names = c("gamma_year","gamma_active","gamma_ET","gamma_NR","
```

	gamma_year	gamma_active	gamma_ET	gamma_NR	gamma_SS	gamma_TS
2.5%	0.003	-0.105	-0.095	-0.093	-0.089	-0.081
97.5%	0.004	0.088	0.099	0.098	0.103	0.103

From the result, we can see Season's 95% Credible Interval does not contain 0. However, it's still not enough to support that hurricane wind speeds have been increasing over the years because γ_{year} does not converge. As for Active season vs Inactive season, since the 95% Credible Interval contains 0, we may conclude that there are no seasonal difference in hurricane wind speeds.

What about use less burn in to get the 95% Credible Interval?

```
gamma_list <- read.csv("../data/gamma_list.csv")[,1:2]
gamma_list = slice(gamma_list, 2001:10000)
quantiles <- apply(gamma_list, 2, quantile, c(0.025, 0.975)) # use 2.5 and 97.5 quantile to get 95% cre
quantiles
```

```
##           Season ActiveActive
## 2.5% -0.002400719 -0.10570771
## 97.5% 0.003669652 0.09002346
```

The result does not change much.

So for this task, the answer is that there are no seasonal difference in hurricane wind speeds, and also there is no evidence to support the claim that hurricane wind speeds have been increasing over the years.

This is somehow against the common sense of mankind. Because we would naturally think that, because of global warming, the hurricanes are bound to get worse so that the wind speeds should have been increasing. And active seasons, we define as from August to October. The wind speeds of hurricanes in active season should be larger than those in inactive season, but it is not the case.