Task 2

Task 2: Compute posterior summaries and 95% credible intervals of γ , the fixed effects associated with the covariates in the model. Using the estimated Bayesian model, answer the following questions: (1) are there seasonal differences in hurricane wind speeds, and (2) is there evidence to support the claim that hurricane wind speeds have been increasing over the years?

Import gamma from task 1 and 2.

```
library(MASS)
library(tidyverse)
```


```
## -- Attaching packages ------ tidyverse 1.3.2 --
## v ggplot2 3.4.0
                     v purrr
                              0.3.4
## v tibble 3.1.8
                     v dplyr
                            1.0.10
## v tidyr
          1.2.1
                     v stringr 1.5.0
## v readr
           2.1.3
                     v forcats 1.0.0
## Warning: package 'ggplot2' was built under R version 4.2.2
## Warning: package 'stringr' was built under R version 4.2.2
## Warning: package 'forcats' was built under R version 4.2.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()
                  masks stats::lag()
## x dplyr::select() masks MASS::select()
```

We are mainly interested in γ_{Season} which is about year change, and γ_{Active} which is about active season change compared to inactive season.

```
gamma_list <- read.csv("./data/gamma_list.csv")[,1:2]
summary(gamma_list)</pre>
```


```
##
       Season
                       ActiveActive
          :0.0001799
   Min.
                      Min.
                             :-0.192543
                      1st Qu.:-0.040679
##
   1st Qu.:0.0025282
## Median :0.0032784
                      Median :-0.006659
## Mean :0.0031620
                      Mean :-0.007029
   3rd Qu.:0.0037629
                       3rd Qu.: 0.026579
##
   Max.
          :0.0059009
                      Max.
                             : 0.168031
```

See the time-series plot to make decision about burn in



plot(gamma_list[,2], type = "1") # plot gamma2(active season)


seems converge, keep the number



```
# set burn in = 8000
gamma_list = slice(gamma_list, 8001:10000)


# see the histogram of year, some kind of normal
ggplot(gamma_list, aes(x = Season))+geom_histogram()
```

'stat_bin()' using 'bins = 30'. Pick better value with 'binwidth'.



qqnorm(gamma_list\$Season)
qqline(gamma_list\$Season) # maybe normal

Normal Q-Q Plot



```
shapiro.test(gamma_list$Season) # not normal

##

## Shapiro-Wilk normality test

##

## data: gamma_list$Season

## W = 0.98714, p-value = 2.097e-12

quantiles <- apply(gamma_list, 2, quantile, c(0.025, 0.975)) # use 2.5 and 97.5 quantile to get 95% cre
quantiles

## Season ActiveActive</pre>
```

From the result, we can see Season's 95% Credible Interval does not contain 0. However, it's still not enough to support that hurricane wind speeds have been increasing over the years because γ_{year} does not converge. As for Active season vs Inactive season, since the 95% Credible Interval contains 0, we may conclude that there are no seasonal difference in hurricane wind speeds.

What about use less burn in to get the 95% Credible Interval?

0.08510004

2.5% 0.002901574 -0.10281169

97.5% 0.004152531

```
gamma_list <- read.csv("./data/gamma_list.csv")[,1:2]
gamma_list = slice(gamma_list, 2001:10000)
quantiles <- apply(gamma_list, 2, quantile, c(0.025, 0.975)) # use 2.5 and 97.5 quantile to get 95% cre
quantiles</pre>
```

```
## Season ActiveActive
## 2.5% 0.001850649 -0.10281169
## 97.5% 0.005320907 0.09062648
```

The result does not change much.

So for this task, the answer is that there are no seasonal difference in hurricane wind speeds, and also there is no evidence to support the claim that hurricane wind speeds have been increasing over the years.

This is somehow against the common sense of mankind. Because we would naturally think that, because of global warming, the hurricanes are bound to get worse so that the wind speeds should have been increasing. And active seasons, we define as from August to October. The wind speeds of hurricanes in active season should be larger than those in inactive season, but it is not the case.