

Development and ocular histology

From tissue origins to the microscopic lesion

Build the eye sequence and origins

The core sequence

- Week 4: forebrain neuroectoderm evaginates into optic vesicles
↓
- Contact with surface ectoderm induces a lens placode
↓
- Optic vesicle invaginates into a two-layered optic cup
↓
- Lens placode invaginates; lens vesicle separates by week 5
↓
- Optic fissure closes during week 7

Know the tissue of origin

Origin	High-yield derivatives
Optic cup neuroectoderm	Neural retina, retinal pigment epithelium; iris and ciliary epithelium; sphincter and dilator pupillae
Surface ectoderm	Lens and corneal epithelium
Periocular mesenchyme	Corneal stroma/endothelium, sclera, choroid and ciliary muscle
Mesoderm	Extraocular muscle fibres

A useful exception

The iris sphincter and dilator are smooth muscles derived from neuroectoderm. Do not infer their embryological origin from the usual origin of muscle; distinguish these muscles from the mesenchymal ciliary muscle.

Link development to adult anatomy

- Outer optic cup becomes the pigment epithelium; inner optic cup becomes the neural retina. Their apposition leaves a vulnerable potential plane.
- Retinal ganglion-cell axons grow through the optic stalk, establishing the optic nerve connection to the brain.
- The hyaloid artery enters through the optic fissure to supply the developing lens. Its distal lens-directed portion regresses; the proximal portion contributes to the central retinal artery.

Sources VS Development, slides 3-11, 14-15 | VS TBL 1, Development transcript | Gage et al.: Ocular lineage study

Developmental defects infer the failed step

Three mechanisms worth separating

Finding	Developmental mechanism
Iris cleft, possibly extending posteriorly	Coloboma: incomplete closure of the embryonic optic fissure
Lens absent	Congenital aphakia: failure of lens induction or subsequent lens development
Lens present but opaque	Congenital cataract: disruption of lens transparency during development

Why a cleft can cross several tissues

The optic fissure runs along the ventral developing optic cup and stalk. Failure of closure can involve iris, ciliary body, choroid, retina or optic nerve. A defect involving several of these structures is one developmental event, rather than several independent acquired lesions.

Lens maturation

- Posterior lens-vesicle cells elongate anteriorly
↓
- Primary lens fibres fill the vesicle lumen
↓
- Anterior cells remain as cuboidal epithelium
↓
- Further fibres are added as the lens grows

Use the stem, not the label

- An intact but cloudy lens supports cataract; failure to induce a lens supports aphakia. A present lens does not exclude coloboma.
- Maternal rubella is a classic association with congenital cataract. Early embryonic exposure is relevant; avoid treating a particular gestational week as an absolute safe cutoff.
- Retina develops into three main nuclear/cell-body bands: outer nuclear, inner nuclear and ganglion-cell layers. The ten-layer microscopic description includes their processes and boundaries.
- The lens vesicle separates from surface ectoderm; the remaining ectoderm forms corneal epithelium.

Cornea and lens transparency by structure

Cornea anterior to posterior

Layer	Recognise the function
Epithelium	Non-keratinised stratified squamous barrier; regenerates
Bowman layer	Acellular anterior stromal layer
Stroma	Bulk of cornea; organised collagen lamellae
Descemet membrane	Basement membrane of the endothelium
Endothelium	Fluid transport maintains relative stromal dehydration

Predict the effect of injury

The cornea is avascular and transparent. Regular stromal organisation and controlled hydration minimise light scattering. Endothelial failure allows stromal swelling and reduces clarity, even when the epithelium remains intact; an epithelial abrasion disrupts a different barrier.

Recognise the lens on a section


- A biconvex, avascular structure enclosed by a thick basement-membrane capsule.
- Cuboidal epithelium lies beneath the anterior capsule. The adult posterior surface is not covered by a matching epithelial layer.
- Long, closely packed lens fibres contain crystallins. Mature fibres lose their nuclei, reducing structures that scatter light.
- Lens-fibre injury produces opacity: cataract. The capsule and the epithelial lining are different structures.

Exam pitfall

Descemet membrane supports the posterior endothelium; Bowman layer sits immediately behind the anterior epithelium. Neither is the lens capsule. Match the location before choosing a histological label.

Retina map the cell to the layer

Outer retina to vitreous



Outer means nearer the choroid. Inner means nearer the vitreous. Nuclear layers hold cell bodies; plexiform layers hold synapses.

Two opposite directions

- Light: vitreous side → inner retina → photoreceptors
- Signal: photoreceptor → bipolar cell → ganglion cell
- Ganglion axons: nerve-fibre layer → optic nerve

Cells that change the answer

- Rods support dim-light sensitivity; cones support colour and fine detail. Photoreceptor segments and cell bodies occupy different layers.
- Horizontal cells modulate photoreceptor-to-bipolar communication. Amacrine cells modulate signalling in the inner plexiform layer.
- Müller cells support retinal structure and function. Plexiform layers contain connections; nuclear layers contain cell bodies.
- RPE absorbs stray light, supports the visual-pigment cycle and forms the outer blood-retinal barrier.


Detachment is a boundary problem

Retinal detachment separates neural retina from RPE at their embryological apposition plane. Photoreceptors lose essential contact with RPE and nearby choroidal support; this is distinct from vitreous separation at the inner retinal surface.

Sources VS Histology, slides 9-15 | VS Anatomy, retinal layers | Senior Eye notes, acute painless visual loss


Picture the tissue and its origin

1 One globe, different tissues



Original schematic. The corneal inset enlarges the conventional five layers; their thicknesses are illustrative. Lens opacity and corneal oedema affect different clear tissues.

2 Follow the wall, predict the derivative



Original developmental schematic. The cup lip is continuous between its inner and outer walls. The apposition plane between neural retina and RPE remains vulnerable to detachment.

Use the pictures to reason

A missing lens points to failed induction; a present opaque lens points to cataract. The cup rim also forms iris/ciliary epithelia and the neuroectodermal iris muscles.