

Cataract surgery: planning, fluidics and complication control

Part 2 | Crowding changes the operative risk model


A short eye needs more than a high-power implant calculation.

Constraint	Planning consequence
Small anterior segment	Limited working space
Thick sclera or effusion tendency	Assess posterior pressure risk
High-power lens requirement	Prediction and positioning errors matter
Poor visual history	Clarify amblyopia and posterior potential

Distinguish simple short axial length from the broader anatomy of a very small eye. Imaging can reveal subtle uveal effusion or structural constraints. Plan anesthesia, fluidics and contingencies with the anatomy in mind. A narrow chamber should not be treated by repeatedly increasing infusion without identifying its cause.

Pseudoexfoliation combines pupil and support risk

Removing the cataract does not remove the underlying zonulopathy.



Original map: pupil access, capsule support and pressure must each be planned.

Evaluate dilation, phacodonesis, chamber asymmetry, pressure and endothelial status. Prepare suitable support devices and alternatives if the bag cannot safely retain an implant. Long-term contraction or bag-IOL instability can emerge after an initially successful operation. A technically quiet early result does not end surveillance of the support system.

An unstable iris responds to flow as well as diameter

Anticipation is safer than repeated rescue through a compromised pupil.

Feature	Planning consequence
Billowing in fluid currents	Modify fluidics and maintain control
Progressive miosis	Adequate access may need an expansion device
Iris prolapse	Reassess gradients and wound configuration
Past alpha-blocker exposure	Risk may persist after discontinuation

Mechanical stretching alone may not control a floppy iris. Consider suitable pharmacological and mechanical strategies with attention to preparation safety. Review medication history before surgery and coordinate systemic medication decisions. A drug-free interval does not reliably guarantee elimination of IFIS risk.

The vitrectomized eye has altered support dynamics

A deep chamber can coexist with unstable lens-capsule behaviour.

Altered feature	Consequence
Reduced posterior support	Chamber depth may fluctuate
Possible zonular damage	Assess capsule stability
Prior retinal pathology	Limits potential visual recovery
Prior intraocular gas or oil	Review current anatomy and implant planning

Review the retinal operative history and posterior visual potential. Adjust technique to the actual chamber and capsule response rather than assuming that a wide space is easy surgery. Coordinate combined versus staged approaches where relevant; avoiding a second procedure is not automatically the best strategy for every retina and cornea.

A complication changes the remaining plan

Stabilization, safe support and follow-up are parts of the operation.



Original ordered map: recognize, stabilize, reassess support, then choose the safe completion path.

After an unexpected event, reassess vitreous, residual lens material, IOL support and the need for another specialist. Document what changed and explain the follow-up implications. Infection prevention and processing standards continue to apply during rescue. A completed lens extraction is not the only endpoint; tissue safety and an appropriate implant position matter.