

Mitochondrial dynamics in yeast cell death and aging

Ralf J. Braun¹ and Benedikt Westermann

Institut für Zellbiologie, Universität Bayreuth, 95440 Bayreuth, Germany

Abstract

Mitochondria play crucial roles in programmed cell death and aging. Different stimuli activate distinct mitochondrion-dependent cell death pathways, and aging is associated with a progressive increase in mitochondrial damage, culminating in oxidative stress and cellular dysfunction. Mitochondria are highly dynamic organelles that constantly fuse and divide, forming either interconnected mitochondrial networks or separated fragmented mitochondria. These processes are believed to provide a mitochondrial quality control system and enable an effective adaptation of the mitochondrial compartment to the metabolic needs of the cell. The baker's yeast, *Saccharomyces cerevisiae*, is an established model for programmed cell death and aging research. The present review summarizes how mitochondrial morphology is altered on induction of cell death or on aging and how this correlates with the induction of different cell death pathways in yeast. We highlight the roles of the components of the mitochondrial fusion and fission machinery that affect and regulate cell death and aging.

Introduction

Mitochondria are semi-autonomous organelles that contain their own genome [mtDNA (mitochondrial DNA)], encoding a small subset of mitochondrial proteins. They are bounded by two membranes: the outer and the inner membrane. Mitochondria are essential for various metabolic processes, including oxidative phosphorylation, amino acid metabolism and the formation of iron-sulfur clusters. In addition, mitochondria play crucial roles in the regulation of cell death and aging in various organisms, including humans and yeast [1–4]. In the budding yeast *Saccharomyces cerevisiae*, mitochondrial dysfunction, mutations in mtDNA and ROS (reactive oxygen species) produced by mitochondria have been proposed to determine both chronological lifespan (i.e. the survival time of a post-mitotic yeast culture) and replicative lifespan (i.e. the number of cell divisions of an individual mother cell) [3–5]. In particular, production of detrimental ROS by the NADH:ubiquinone oxidoreductase Ndi1 and the cytochrome *bc*₁ complex of damaged mitochondria was shown to be an important factor promoting cell death [6–9]. Mitochondrial permeabilization and the release of mitochondrial proteins into the cytosol define distinct mitochondrion-dependent cell death pathways [10–13]. Release of cytochrome *c* correlates with activation of the cell death protease Yca1 (yeast caspase 1), which ultimately executes cell death [13,13a]. In Yca1-independent pathways, Aif1 (apoptosis-inducing factor 1) or endonuclease G (Nuc1) are released from mitochondria and translocate into the nucleus, resulting in the fragmentation of the nuclear

genome [10,12]. Mitochondria are also involved in cell death pathways mediated by the alternative cell death protease Kex1 or Ysp1 (yeast suicide protein 1) and Ysp2 [9,14–16]. However, the molecular mechanisms of these cell death scenarios remain poorly understood.

Mitochondria are highly dynamic organelles that continuously fuse and divide [17]. In many organisms, including worms and mammals, mitochondrial division promotes the release of cytochrome *c* to trigger apoptosis [1]. Furthermore, mitochondrial dynamics is thought to counteract aging and constitute an organellar quality control mechanism. Fusion allows complementation and repair processes of damaged mitochondria, whereas fission separates defective organelles from the mitochondrial network, which are then subjected to degradation through mitophagy [17]. In addition to their role in apoptosis and mitochondrial quality control, mitochondrial fusion and fission are needed to optimally adapt mitochondria to the metabolic needs of the cell [17]. In yeast, inhibition of fusion leads to damage and loss of mtDNA [18–20]. In contrast, mitochondrial fusion increases mtDNA copy number and ensures high accuracy of the encoded genetic information [21]. While respiratory-active yeast cells harbour an extended tubular mitochondrial network, resting non-dividing cells contain mostly fragmented mitochondria [22].

A conserved molecular machinery mediates mitochondrial fusion and fission in humans and yeast [17]. In yeast, the large GTPase Fzo1, a member of the mitofusin protein family, promotes fusion of the mitochondrial outer membrane, whereas the dynamin-related large GTPase Mgm1 enables fusion of the inner membrane [18,19,23]. Outer and inner membrane fusion is co-ordinated by the outer membrane protein Ugo1, which physically interacts with both Fzo1 and Mgm1 [24]. Division of the outer membrane is performed by the dynamin-related large GTPase Dnm1, which binds via


Key words: aging, cell death, mitochondrial dynamics, mitochondrial fission, mitochondrial fusion, *Saccharomyces cerevisiae*.

Abbreviations used: Aif1, apoptosis-inducing factor 1; mtDNA, mitochondrial DNA; ROS, reactive oxygen species; Yca1, yeast caspase 1; Ysp, yeast suicide protein.

¹To whom correspondence should be addressed (email ralf.braun@uni-bayreuth.de).

Figure 1 | Mitochondrial fragmentation and aggregation during programmed cell death and aging in yeast

See the main text for details.



the adaptor proteins Mdv1 or Caf4 to the outer membrane receptor Fis1 and forms a contractile ring that eventually promotes outer membrane division [25–27]. An increasing number of accessory and regulatory components are being identified to regulate the fusion and fission processes. These include the rhomboid-related membrane protease Pcp1, which processes Mgm1 [28], and the F-box protein Mdm30, a ubiquitin ligase subunit mediating the turnover of Fzo1 [29].

In the present review, we summarize abnormal mitochondrial morphologies that were observed in yeast models for programmed cell death and that link these morphologies with different cell death pathways. We focus on the effect of the mitochondrial fusion and fission balance on yeast cell death and aging, and we outline how these processes may be relevant under deleterious conditions.

Changes in mitochondrial morphology during yeast cell death and aging

Fragmentation of the mitochondrial network into multiple small organelles has been observed on treatment of yeast cells with multiple different stressors inducing programmed cell death (Figure 1; Table 1). Mitochondrial fragmentation occurs on (i) acidic stress, including acetic acid [16,30,31], propionic acid [16] and formic acid [32], (ii) oxidative stress (H_2O_2) [33], (iii) treatment with drugs and fungicides (amiodarone [9,15,16], bostrycin [34] and trichothecene [35]), (iv) yeast pheromones (α -factor) [9] and (v) ethanol [36]. A stringent correlation between mitochondrial fragmentation and yeast cell death was further observed in a variety of mutant yeast strains (Table 1), including strains with mutations in (i) rRNA genes (HsTnII) [37], (ii) mRNA turnover genes (*lsm4*) [38,39], (iii) genes involved in glycoprotein biosynthesis (*wbp1-1*) [14] and (iv) stress response genes ($\Delta wbi2$) [40]. Furthermore, cell death concomitant with mitochondrial fragmentation was observed in yeast cells overexpressing signalling kinases (*TPK3*)

[41], sphingolipid-metabolizing enzymes (*YDC1*) [42] or overexpressing human proteins, including the pro-apoptotic protein BAX [43], and a Huntington's disease-causing variant of huntingtin [44]. Finally, yeast cells that enter the stationary phase and undergo chronological aging contain fragmented mitochondria [22,45]. The wide variety of death-inducing conditions that correlate with disruption of the mitochondrial network indicates that mitochondrial morphology is strongly influenced by the state of health of the cell. Conversely, conditions that trigger mitochondrial fragmentation frequently result in a decreased chronological lifespan [37–39,42].

In some of the cell death scenarios described above, mitochondrial fragmentation is superimposed by aberrant mitochondrial aggregation [40,41,44], which is an active process requiring the actin cytoskeleton [31,40,41]. Aggregated mitochondrial fragments accumulate on disruption of the mitochondrion degradation pathway [31,43], suggesting that most of these organelles are damaged and destined for degradation. Indeed, fragmented and/or aggregated mitochondria were found to be physically and functionally impaired during cell death (Figure 1; Table 1). Mitochondrial deterioration during cell death is characterized by the following events: loss of the mitochondrial membrane potential [32,37], which can be preceded by a strong increase [9], permeabilization of the mitochondrial outer membrane [31], release of cytochrome *c* [9,31] and loss of mtDNA [40]. As a consequence, cells undergoing cell death frequently become respiratory deficient [37,41]. Notably, the accumulation of ROS appears to be a general hallmark of cell death in cells with fragmented and/or aggregated mitochondria [9,14–16,32,34,36,38,39,41,44]. This suggests that damaged mitochondria are a major source of ROS during cell death and thereby actively contribute to the cellular demise (Figure 1).

Cell death correlated with mitochondrial fragmentation either depends on the activity of the yeast cell death protease

Table 1 | Yeast cell death and aging associated with abnormal mitochondrial morphologies

	Inducers of cell death	Mitochondrial morphology	Yeast cell death pathway	Reference(s)
Stressors	Acetic acid	Fragmentation; <i>FIS1</i> is not required for fragmentation; actin-dependent formation of mitochondrial aggregates in $\Delta pep4$ and $\Delta aac1 \Delta aac2 \Delta aac3$	Mitochondrial outer membrane permeabilization; cytochrome <i>c</i> release; <i>YCA1</i> , <i>DNM1</i> and <i>MDV1</i> promote cell death; <i>FIS1</i> relieves cell death	[30,31]
	α -Factor, amiodarone, acetic acid, propionic acid	Fragmentation; fragmentation depends on <i>Ysp1</i> and <i>Ysp2</i>	Cell death depends on <i>YSP1</i> and <i>YSP2</i> ; increased mitochondrial membrane potential; ROS accumulation; cytochrome <i>c</i> release	[9,15,16]
	Bostrycin	Fragmentation	<i>AIF1</i> -dependent and <i>YCA1</i> -independent cell death; ROS	[34]
	Ethanol	Fragmentation; fragmentation depends on <i>FIS1</i> but is independent of <i>DNM1</i> and <i>MDV1</i> ; at high concentrations, <i>FIS1</i> is not required	ROS accumulation depends on <i>FIS1</i> ; cell death is independent of <i>YCA1</i> , <i>AIF1</i> , cytochrome <i>c</i> , <i>MDV1</i> and <i>DNM1</i>	[36]
	Formic acid	Fragmentation	Decreased mitochondrial membrane potential; <i>YCA1</i> -independent cell death; ROS accumulation	[32]
	H ₂ O ₂	Fragmentation	Decreased viability	[33]
	Trichothecene	Fragmentation	Mutants of the mitochondrial fusion machinery are resistant to trichothecene-triggered growth deficit	[35]
Gene deletion/mutation	HsTnII	Fragmentation	Decreased mitochondrial membrane potential; respiratory deficiency; decreased chronological lifespan	[37]
	<i>Ism4</i>	<i>YCA1</i> -dependent fragmentation	<i>YCA1</i> -dependent cell death; ROS accumulation; decreased chronological lifespan	[38,39]
	<i>wbp1-1</i>	<i>KEX1</i> -dependent fragmentation on abnormal N-glycosylation	<i>KEX1</i> -dependent and <i>YCA1</i> -independent cell death; ROS accumulation	[14]
	$\Delta whi2$	Fragmentation and aggregation; actin-dependent mitochondrial aggregation; mtDNA loss	Deletion of <i>WHI2</i> results in apoptosis triggered by actin aggregation	[40]
Overexpression	BAX	Fragmentation in wild-type strain; aggregation in mitophagy-deficient $\Delta uth1$ strain	<i>AIF1</i> - and <i>YCA1</i> -independent clonogenic cell death; plasma membrane permeabilization on <i>UTH1</i> deletion	[43]
	Disease-associated human huntingtin	Fragmentation and aggregation	ROS accumulation; caspase activation; <i>YCA1</i> -dependent nuclear localization of huntingtin aggregates	[44]
	<i>TPK3</i>	Aggregation	Respiratory deficiency and ROS accumulation depends on <i>Tpk3</i> activity	[41]
	<i>YDC1</i>	Fragmentation	Decreased chronological lifespan and increased apoptosis	[42]

Table 2 | Effect of gene deletions on yeast cell death and aging

	Gene deletion	Inducers of cell death	Cell death pathway on induction	Influence of gene deletion on viability/cell death/lifespan	Reference(s)	
Mitochondrial fission	$\Delta dnm1$	H ₂ O ₂ , acetate, BAR0329, M1 killer virus	YCA1-dependent cell death; ROS accumulation	Decreased cytotoxicity	[30,33,46,47]	
		Ethanol	YCA1- and AIF1-independent cell death; ROS accumulation	Unaltered cytotoxicity	[36]	
		Aging	–	Increased chronological and replicative lifespan	[33,39,45]	
	$\Delta fis1$ (<i>whi2</i>)	H ₂ O ₂ , acetate, BAR0329, M1 killer virus	YCA1-dependent cell death; ROS accumulation	Increased cytotoxicity; WHI2 mutation but not FIS1 deletion confers sensitivity for cell death	[30,33,46–48]	
		Ethanol	YCA1- and AIF1-independent cell death; ROS accumulation	Increased cytotoxicity	[36]	
		Aging	–	Increased chronological and replicative lifespan	[33,39,45]	
	$\Delta mdv1$	H ₂ O ₂ , acetate, BAR0329	YCA1-dependent cell death; ROS accumulation	Decreased cytotoxicity	[30,46]	
		Ethanol	YCA1- and AIF1-independent cell death; ROS accumulation	Unaltered cytotoxicity	[36]	
	Mitochondrial fusion	$\Delta fzo1$	Aging	–	Increased chronological lifespan	[39]
			Trichothecene	–	Growth deficit relieved	[35]
$\Delta mdm30$		H ₂ O ₂	–	Decreased cytotoxicity	[33]	
		Aging	–	Increased chronological lifespan	[33,39]	
$\Delta pcg1$		Trichothecene	–	Growth deficit relieved	[35]	

Yca1 [30,38,39,44] or is promoted independently of Yca1 by factors including the alternative cell death protease Kex1 [14], the mitochondrial cell death factors Aif1, Ysp1 and Ysp2 [9,15,16,34] or by so far unknown factors. Thus it appears that mitochondrial fragmentation concomitant with mitochondrial dysfunction and ROS accumulation are common and crucial events during cell death, whereas the cell death executing pathways may be very different and specified by the inducer of cell death.

Impact of the molecular machinery of mitochondrial fusion and fission on yeast cell death and aging

The correlation of mitochondrial fragmentation and yeast cell death suggests that preventing fragmentation by inhibiting

mitochondrial fission (or increasing mitochondrial fusion) might be beneficial for cell survival. Indeed, deletion of the mitochondrial fission factors Dnm1 or Mdv1 results in increased resistance against various cell death stressors, including acetate [30], H₂O₂ [30,33], the fungicide BAR0329 [46] and the M1 killer virus [47] (Table 2). These observations suggest that Dnm1 and Mdv1 promote cell death. However, deletion of the *FIS1* gene encoding the Dnm1 receptor was surprisingly found to have an opposite effect. Cells lacking Fis1 are highly susceptible to ethanol [36] and all kinds of stressors, against which $\Delta dnm1$ and $\Delta mdv1$ mutants are more resistant [30,33,46,47] (Table 2). Intriguingly, later, it was demonstrated that deletion of the *FIS1* gene reproducibly results in the spontaneous acquisition of a secondary mutation in the stress-response gene *WHI2* [48].

Yeast cells lacking functional *Whi2* are highly sensitive to undergo cell death [40,48], and their mitochondrial network is prone to fragmentation [40] (Table 1). Δ *fis1* mutants that have acquired secondary *whi2* mutations are as sensitive to stress as *whi2* mutants in a *FIS1* wild-type background, even though they contain a highly interconnected mitochondrial network like Δ *dnm1* or Δ *mdv1* strains [48]. Thus the increased susceptibility of Δ *fis1* cells to undergo cell death is due to mutations in *WHI2* rather than inhibition of mitochondrial fission [48]. Therefore it is reasonable to assume that mitochondrial fragmentation by the fission machinery indeed promotes cell death in yeast.

Remarkably, Dnm1- and Mdv1-promoted cell death depends on Yca1 in all the scenarios tested so far [30,46,47]. Thus it appears that a cell death pathway executed by the dynamin-related mitochondrial fission machinery and caspase-related factors has been conserved from yeast to worms and mammals [1]. On the other hand, stressing yeast cells with high concentrations of ethanol or acetic acid results in mitochondrial fragmentation that is independent of the mitochondrial fission factors Dnm1, Mdv1 and/or Fis1 [30,36] (Table 1). In this pathway, cell death is mediated by alternative cell death factors, such as Aif1, Kex1, Ysp1 and Ysp2 [9,14–16,34] (Figure 1; Table 1). These observations suggest that detrimental environmental conditions may induce mitochondrial fragmentation and cell death in a way that is independent of the known fission machinery and Yca1.

As mitochondrial fragmentation facilitates cell death, it can be expected that inhibition of mitochondrial fusion might have a similar effect (Figure 1). However, a genome-wide screening for resistance against the mycotoxin trichothecene demonstrated that yeast strains with the mitochondrial fusion genes *FZO1* and *PCP1* deleted are highly resistant to the toxin, rather than more susceptible to it [35] (Table 2). Similarly, a yeast strain with *MDM30* deleted demonstrated an increased resistance to H₂O₂ treatment [33] (Table 2). These unexpected phenotypes might be explained by the fact that fusion-deficient yeast strains have a high tendency to lose their mtDNA [18–20]. Since cells depleted of mtDNA were shown to be completely resistant to trichothecene treatment [35], these results suggest that the beneficial effect of the deletion of mitochondrial fusion factors can be ascribed to a loss of mtDNA rather than a blocking of fusion.

Conditions that promote mitochondrial fragmentation and cell death lead to a decreased chronological lifespan [37–39,42] (Table 1). On the other hand, caloric restriction is associated with an increased chronological lifespan in yeast [3,49,50] and leads to the down-regulation of the fission factors Fis1, Mdv1 and Caf4 concomitant with up-regulation of the fusion factor Mgm1 [51]. Thus inhibition of mitochondrial fragmentation may be associated with a prolonged lifespan. Indeed, deletion of the mitochondrial fission gene *DNM1* was shown to significantly increase both the chronological and the replicative lifespan of yeast cells [33,39,45] (Table 2). Consistent with this, deletion of the *FIS1* gene also resulted in increased chronological and replicative lifespan, in spite of the presence of its pro-

death secondary mutation in the stress response gene *WHI2* [33,39,45] (Table 2). These observations suggest that during aging the pro-survival phenotype of the Δ *fis1* deletion overrules the pro-death function of the *whi2* mutation that appears to be dominant only in younger cultures.

Surprisingly, deletion of the fusion gene *FZO1* resulted in a slightly prolonged chronological lifespan [33,39], and cells with *MDM30* deleted demonstrated even a 60% increase in chronological lifespan [33,39]. Future experiments will have to show whether the fusion-incompetence or the lack of respiratory capacity and decreased mtDNA stability are responsible for longevity in these mutants.

Conclusions

A variety of different cell death-inducing conditions, including cell stress, gene mutations and overexpression, result in mitochondrial dysfunction accompanied by a boost of ROS and the release of mitochondrial pro-death factors, such as cytochrome *c* and Aif1. In addition to the deterioration of mitochondrial functions, fragmentation of the mitochondrial network appears to be a hallmark of various cell death pathways, including *YCA1*-dependent and -independent pathways. Whereas inhibition of mitochondrial fission was shown to result in increased stress resistance and prolonged chronological and replicative lifespan, the role of mitochondrial fusion is less clear. Future studies are required to determine whether the known components of the mitochondrial fusion and fission machineries play a general role in yeast cell death and aging or whether additional factors exist.

Funding

We are grateful to the Deutsche Forschungsgemeinschaft for funding our work.

References

- 1 Suen, D.F., Norris, K.L. and Youle, R.J. (2008) Mitochondrial dynamics and apoptosis. *Genes Dev.* **22**, 1577–1590
- 2 Cheng, W.C., Leach, K.M. and Hardwick, J.M. (2008) Mitochondrial death pathways in yeast and mammalian cells. *Biochim. Biophys. Acta* **1783**, 1272–1279
- 3 Fabrizio, P. and Longo, V.D. (2008) Chronological aging-induced apoptosis in yeast. *Biochim. Biophys. Acta* **1783**, 1280–1285
- 4 Laun, P., Heeren, G., Rinnerthaler, M., Rid, R., Kössler, S., Koller, L. and Breitenbach, M. (2008) Senescence and apoptosis in yeast mother cell-specific aging and in higher cells: a short review. *Biochim. Biophys. Acta* **1783**, 1328–1334
- 5 Herker, E., Jungwirth, H., Lehmann, K.A., Maldener, C., Fröhlich, K.U., Wissing, S., Büttner, S., Fehr, M., Sigrist, S. and Madeo, F. (2004) Chronological aging leads to apoptosis in yeast. *J. Cell Biol.* **164**, 501–507
- 6 Braun, R.J., Sommer, C., Carmona-Gutierrez, D., Khoury, C.M., Ring, J., Büttner, S. and Madeo, F. (2011) Neurotoxic 43-kDa TAR DNA-binding protein (TDP-43) triggers mitochondrion-dependent programmed cell death in yeast. *J. Biol. Chem.* **286**, 19958–19972

- 7 Braun, R.J., Zischka, H., Madeo, F., Eisenberg, T., Wissing, S., Büttner, S., Engelhardt, S.M., Bühringer, D. and Ueffing, M. (2006) Crucial mitochondrial impairment upon *CDC48* mutation in apoptotic yeast. *J. Biol. Chem.* **281**, 25757–25767
- 8 Li, W., Sun, L., Liang, Q., Wang, J., Mo, W. and Zhou, B. (2006) Yeast AMID homologue Ndi1p displays respiration-restricted apoptotic activity and is involved in chronological aging. *Mol. Biol. Cell* **17**, 1802–1811
- 9 Pozniakovsky, A.I., Knorre, D.A., Markova, O.V., Hyman, A.A., Skulachev, V.P. and Severin, F.F. (2005) Role of mitochondria in the pheromone- and amiodarone-induced programmed death of yeast. *J. Cell Biol.* **168**, 257–269
- 10 Büttner, S., Eisenberg, T., Carmona-Gutierrez, D., Ruli, D., Knauer, H., Ruckenstein, C., Sigrist, C., Wissing, S., Kollroser, M., Fröhlich, K.U. et al. (2007) Endonuclease G regulates budding yeast life and death. *Mol. Cell* **25**, 233–246
- 11 Pereira, C., Camougrand, N., Manon, S., Sousa, M.J. and Corte-Real, M. (2007) ADP/ATP carrier is required for mitochondrial outer membrane permeabilization and cytochrome *c* release in yeast apoptosis. *Mol. Microbiol.* **66**, 571–582
- 12 Wissing, S., Ludovico, P., Herker, E., Büttner, S., Engelhardt, S.M., Decker, T., Link, A., Proksch, A., Rodrigues, F., Corte-Real, M. et al. (2004) An AIF orthologue regulates apoptosis in yeast. *J. Cell Biol.* **166**, 969–974
- 13 Ludovico, P., Rodrigues, F., Almeida, A., Silva, M.T., Barrientos, A. and Corte-Real, M. (2002) Cytochrome *c* release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **13**, 2598–2606
- 13a Madeo, F., Herker, E., Maldener, C., Wissing, S., Lächelt, S., Herlan, M., Fehr, M., Lauber, K., Sigrist, S.J., Wesselborg, S. and Fröhlich, K.U. (2002) A caspase-related protease regulates apoptosis in yeast. *Mol. Cell* **9**, 911–917
- 14 Hauptmann, P. and Lehle, L. (2008) Kex1 protease is involved in yeast cell death induced by defective N-glycosylation, acetic acid, and chronological aging. *J. Biol. Chem.* **283**, 19151–19163
- 15 Knorre, D.A., Ojovan, S.M., Saprunova, V.B., Sokolov, S.S., Bakeeva, L.E. and Severin, F.F. (2008) Mitochondrial matrix fragmentation as a protection mechanism of yeast *Saccharomyces cerevisiae*. *Biochemistry* **73**, 1254–1259
- 16 Sokolov, S., Knorre, D., Smirnova, E., Markova, O., Pozniakovsky, A., Skulachev, V. and Severin, F. (2006) Ysp2 mediates death of yeast induced by amiodarone or intracellular acidification. *Biochim. Biophys. Acta* **1757**, 1366–1370
- 17 Westermann, B. (2010) Mitochondrial fusion and fission in cell life and death. *Nat. Rev. Mol. Cell Biol.* **11**, 872–884
- 18 Hermann, G.J., Thatcher, J.W., Mills, J.P., Hales, K.G., Fuller, M.T., Nunnari, J. and Shaw, J.M. (1998) Mitochondrial fusion in yeast requires the transmembrane GTPase Fzo1p. *J. Cell Biol.* **143**, 359–373
- 19 Rapaport, D., Brunner, M., Neupert, W. and Westermann, B. (1998) Fzo1p is a mitochondrial outer membrane protein essential for the biogenesis of functional mitochondria in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **273**, 20150–20155
- 20 Merz, S. and Westermann, B. (2009) Genome-wide deletion mutant analysis reveals genes required for respiratory growth, mitochondrial genome maintenance and mitochondrial protein synthesis in *Saccharomyces cerevisiae*. *Genome Biol.* **10**, R95
- 21 Hori, A., Yoshida, M. and Ling, F. (2011) Mitochondrial fusion increases the mitochondrial DNA copy number in budding yeast. *Genes Cells* **16**, 527–544
- 22 Merz, S., Hammermeister, M., Altmann, K., Dürr, M. and Westermann, B. (2007) Molecular machinery of mitochondrial dynamics in yeast. *Biol. Chem.* **388**, 917–926
- 23 Wong, E.D., Wagner, J.A., Gorsich, S.W., McCaffery, J.M., Shaw, J.M. and Nunnari, J. (2000) The dynamin-related GTPase, Mgm1p, is an intermembrane space protein required for maintenance of fusion competent mitochondria. *J. Cell Biol.* **151**, 341–352
- 24 Sesaki, H. and Jensen, R.E. (2004) Ugo1p links the Fzo1p and Mgm1p GTPases for mitochondrial fusion. *J. Biol. Chem.* **279**, 28298–28303
- 25 Mozdy, A.D., McCaffery, J.M. and Shaw, J.M. (2000) Dnm1p GTPase-mediated mitochondrial fission is a multi-step process requiring the novel integral membrane component Fis1p. *J. Cell Biol.* **151**, 367–380
- 26 Tieu, Q. and Nunnari, J. (2000) Mdv1p is a WD repeat protein that interacts with the dynamin-related GTPase, Dnm1p, to trigger mitochondrial division. *J. Cell Biol.* **151**, 353–366
- 27 Griffin, E.E., Graumann, J. and Chan, D.C. (2005) The WD40 protein Caf4p is a component of the mitochondrial fission machinery and recruits Dnm1p to mitochondria. *J. Cell Biol.* **170**, 237–248
- 28 Herlan, M., Vogel, F., Bornhövd, C., Neupert, W. and Reichert, A.S. (2003) Processing of Mgm1 by the rhomboid-type protease Pcp1 is required for maintenance of mitochondrial morphology and of mitochondrial DNA. *J. Biol. Chem.* **278**, 27781–27788
- 29 Fritz, S., Weinbach, N. and Westermann, B. (2003) Mdm30 is an F-box protein required for maintenance of fusion-competent mitochondria in yeast. *Mol. Biol. Cell* **14**, 2303–2313
- 30 Fannjiang, Y., Cheng, W.C., Lee, S.J., Qi, B., Pevsner, J., McCaffery, J.M., Hill, R.B., Basanez, G. and Hardwick, J.M. (2004) Mitochondrial fission proteins regulate programmed cell death in yeast. *Genes Dev.* **18**, 2785–2797
- 31 Pereira, C., Chaves, S., Alves, S., Salin, B., Camougrand, N., Manon, S., Sousa, M.J. and Corte-Real, M. (2010) Mitochondrial degradation in acetic acid-induced yeast apoptosis: the role of Pep4 and the ADP/ATP carrier. *Mol. Microbiol.* **76**, 1398–1410
- 32 Du, L., Su, Y., Sun, D., Zhu, W., Wang, J., Zhuang, X., Zhou, S. and Lu, Y. (2008) Formic acid induces Yca1p-independent apoptosis-like cell death in the yeast *Saccharomyces cerevisiae*. *FEMS Yeast Res.* **8**, 531–539
- 33 Palermo, V., Falcone, C. and Mazzoni, C. (2007) Apoptosis and aging in mitochondrial morphology mutants of *S. cerevisiae*. *Folia Microbiol.* **52**, 479–483
- 34 Xu, C., Wang, J., Gao, Y., Lin, H., Du, L., Yang, S., Long, S., She, Z., Cai, X., Zhou, S. and Lu, Y. (2010) The anthracenedione compound bostrycin induces mitochondria-mediated apoptosis in the yeast *Saccharomyces cerevisiae*. *FEMS Yeast Res.* **10**, 297–308
- 35 McLaughlin, J.E., Bin-Umer, M.A., Tortora, A., Mendez, N., McCormick, S. and Tumer, N.E. (2009) A genome-wide screen in *Saccharomyces cerevisiae* reveals a critical role for the mitochondria in the toxicity of a trichothecene mycotoxin. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 21883–21888
- 36 Kitagaki, H., Araki, Y., Funato, K. and Shimoi, H. (2007) Ethanol-induced death in yeast exhibits features of apoptosis mediated by mitochondrial fission pathway. *FEBS Lett.* **581**, 2935–2942
- 37 Aerts, A.M., Zabrocki, P., Govaert, G., Mathys, J., Carmona-Gutierrez, D., Madeo, F., Winderickx, J., Cammue, B.P. and Thevissen, K. (2009) Mitochondrial dysfunction leads to reduced chronological lifespan and increased apoptosis in yeast. *FEBS Lett.* **583**, 113–117
- 38 Mazzoni, C., Herker, E., Palermo, V., Jungwirth, H., Eisenberg, T., Madeo, F. and Falcone, C. (2005) Yeast caspase 1 links messenger RNA stability to apoptosis in yeast. *EMBO Rep.* **6**, 1076–1081
- 39 Palermo, V., Falcone, C., Calvani, M. and Mazzoni, C. (2010) Acetyl-L-carnitine protects yeast cells from apoptosis and aging and inhibits mitochondrial fission. *Aging Cell* **9**, 570–579
- 40 Leadsham, J.E., Miller, K., Ayscough, K.R., Colombo, S., Martegani, E., Sudbery, P. and Gourlay, C.W. (2009) Whi2p links nutritional sensing to actin-dependent Ras-cAMP-PKA regulation and apoptosis in yeast. *J. Cell Sci.* **122**, 706–715
- 41 Leadsham, J.E. and Gourlay, C.W. (2010) cAMP/PKA signaling balances respiratory activity with mitochondria dependent apoptosis via transcriptional regulation. *BMC Cell Biol.* **11**, 92
- 42 Aerts, A.M., Zabrocki, P., Francois, I.E., Carmona-Gutierrez, D., Govaert, G., Mao, C., Smets, B., Madeo, F., Winderickx, J., Cammue, B.P. and Thevissen, K. (2008) Ydc1p ceramidase triggers organelle fragmentation, apoptosis and accelerated ageing in yeast. *Cell. Mol. Life Sci.* **65**, 1933–1942
- 43 Kissova, I., Plamondon, L.T., Brisson, L., Priault, M., Renouf, V., Schaeffer, J., Camougrand, N. and Manon, S. (2006) Evaluation of the roles of apoptosis, autophagy, and mitophagy in the loss of plating efficiency induced by Bax expression in yeast. *J. Biol. Chem.* **281**, 36187–36197
- 44 Sokolov, S., Pozniakovsky, A., Bocharova, N., Knorre, D. and Severin, F. (2006) Expression of an expanded polyglutamine domain in yeast causes death with apoptotic markers. *Biochim. Biophys. Acta* **1757**, 660–666
- 45 Scheckhuber, C.Q., Erjavec, N., Tinazli, A., Hamann, A., Nystrom, T. and Osiewacz, H.D. (2007) Reducing mitochondrial fission results in increased life span and fitness of two fungal ageing models. *Nat. Cell Biol.* **9**, 99–105
- 46 Bink, A., Govaert, G., Francois, I.E., Pellens, K., Meerpoel, L., Borgers, M., Van Minnebruggen, G., Vroome, V., Cammue, B.P. and Thevissen, K. (2010) A fungicidal piperazine-1-carboxamide induces mitochondrial fission-dependent apoptosis in yeast. *FEMS Yeast Res.* **10**, 812–818

-
- 47 Ivanovska, I. and Hardwick, J.M. (2005) Viruses activate a genetically conserved cell death pathway in a unicellular organism. *J. Cell Biol.* **170**, 391–399
- 48 Cheng, W.C., Teng, X., Park, H.K., Tucker, C.M., Dunham, M.J. and Hardwick, J.M. (2008) Fis1 deficiency selects for compensatory mutations responsible for cell death and growth control defects. *Cell Death Differ.* **15**, 1838–1846
- 49 Rockenfeller, P. and Madeo, F. (2010) Ageing and eating. *Biochim. Biophys. Acta* **1803**, 499–506
- 50 Fontana, L., Partridge, L. and Longo, V.D. (2010) Extending healthy life span: from yeast to humans. *Science* **328**, 321–326
- 51 Goldberg, A.A., Bourque, S.D., Kyryakov, P., Gregg, C., Boukh-Viner, T., Beach, A., Burstein, M.T., Machkalyan, G., Richard, V., Rampersad, S. et al. (2009) Effect of calorie restriction on the metabolic history of chronologically aging yeast. *Exp. Gerontol.* **44**, 555–571
-

Received 8 June 2011
doi:10.1042/BST0391520