

Lecture 13

Compactness for first-order logic

The compactness theorem, non-standard models of arithmetic

Introduction to Logic for Computer Science

Prof Hongseok Yang
KAIST

These slides are minor variants of those made by Prof Worrell and Dr Haase for their logic course at Oxford.

The compactness theorem

Theorem

Let S be a countably infinite set of first-order sentences. Then S is satisfiable if and only if every finite subset of S is satisfiable.

The compactness theorem

Theorem

Let S be a countably infinite set of first-order sentences. Then S is satisfiable if and only if every finite subset of S is satisfiable.

Ex1: Which direction is easy to prove?

The compactness theorem

Theorem

Let S be a countably infinite set of first-order sentences. Then S is satisfiable if and only if every finite subset of S is satisfiable.

Ex1: Which direction is easy to prove?

Ex2: Prove the if direction. Assume that we can skolemise S to \mathcal{T} .

Hint: Consider the Herbrand expansion \mathcal{E} of \mathcal{T} . Use the compactness theorem for prop. logic and the Herbrand theorem.

The compactness theorem

Theorem

Let S be a countably infinite set of first-order sentences. Then S is satisfiable if and only if every finite subset of S is satisfiable.

Ex1: Which direction is easy to prove?

Ex2: Prove the if direction. Assume that we can skolemise S to \mathcal{T} .

Hint: Consider the Herbrand expansion \mathcal{E} of \mathcal{T} . Use the compactness theorem for prop. logic and the Herbrand theorem.

Answer:

- (1) all finite subsets of S are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow S$ is satisfiable.

The compactness theorem

Theorem

Let S be a countably infinite set of first-order sentences. Then S is satisfiable if and only if every finite subset of S is satisfiable.

Ex1: Which direction is easy to prove?

Ex2: Prove the if direction. Assume that we can skolemise S to \mathcal{T} .

Hint: Consider the Herbrand expansion \mathcal{E} of \mathcal{T} . Use the compactness theorem for prop. logic and the Herbrand theorem.

Answer:

- (1) all finite subsets of S are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow S$ is satisfiable.

Justification of individual proof steps

Assumed that we can skolemise \mathcal{S} to \mathcal{T} .

But the assumption may fail. Consider the case that \mathcal{S} could use up all function symbols f_1, f_2, \dots

Justification of individual proof steps

Assumed that we can skolemise \mathcal{S} to \mathcal{T} .

But the assumption may fail. Consider the case that \mathcal{S} could use up all function symbols f_1, f_2, \dots

Trick: Rename f_i to f_{2i} to ensure that infinitely many unused function symbols f_{2i+1} are available.

Justification of individual proof steps

Assumed that we can skolemise \mathcal{S} to \mathcal{T} .

But the assumption may fail. Consider the case that \mathcal{S} could use up all function symbols f_1, f_2, \dots

Trick: Rename f_i to f_{2i} to ensure that infinitely many unused function symbols f_{2i+1} are available.

Ex: Explain why it is ok to rename. You need to prove the following.

- Every finite subset of \mathcal{S} before renaming is satisfiable if and only if every finite subset of \mathcal{S} after renaming is satisfiable.
- The whole of \mathcal{S} before renaming is satisfiable if and only if the whole of \mathcal{S} after renaming is satisfiable.

Justification of individual proof steps

- (1) all finite subsets of \mathcal{S} are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow \mathcal{S}$ is satisfiable.

Justification of individual proof steps

- (1) all finite subsets of \mathcal{S} are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow \mathcal{S}$ is satisfiable.

(1) \Rightarrow (2) since skolemisation preserves satisfiability.

Justification of individual proof steps

- (1) all finite subsets of \mathcal{S} are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow \mathcal{S}$ is satisfiable.

(1) \Rightarrow (2) since skolemisation preserves satisfiability.

(2) \Rightarrow (3) by the Herbrand theorem.

Justification of individual proof steps

- (1) all finite subsets of \mathcal{S} are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow \mathcal{S}$ is satisfiable.

(1) \Rightarrow (2) since skolemisation preserves satisfiability.

(2) \Rightarrow (3) by the Herbrand theorem.

(3) \Rightarrow (4) since propositional logic is compact.

Justification of individual proof steps

- (1) all finite subsets of \mathcal{S} are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow \mathcal{S}$ is satisfiable.

(1) \Rightarrow (2) since skolemisation preserves satisfiability.

(2) \Rightarrow (3) by the Herbrand theorem.

(3) \Rightarrow (4) since propositional logic is compact.

(4) \Rightarrow (5) since a prop. model \mathcal{A} for \mathcal{E} induces a Herbrand model \mathcal{H} :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \iff \mathcal{A} \models P(t_1, \dots, t_k).$$

Justification of individual proof steps

- (1) all finite subsets of \mathcal{S} are satisfiable
- (2) \Rightarrow all finite subsets of \mathcal{T} are satisfiable
- (3) \Rightarrow all finite subsets of \mathcal{E} are satisfiable
- (4) $\Rightarrow \mathcal{E}$ is satisfiable
- (5) $\Rightarrow \mathcal{T}$ is satisfiable
- (6) $\Rightarrow \mathcal{S}$ is satisfiable.

(1) \Rightarrow (2) since skolemisation preserves satisfiability.

(2) \Rightarrow (3) by the Herbrand theorem.

(3) \Rightarrow (4) since propositional logic is compact.

(4) \Rightarrow (5) since a prop. model \mathcal{A} for \mathcal{E} induces a Herbrand model \mathcal{H} :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \iff \mathcal{A} \models P(t_1, \dots, t_k).$$

(5) \Rightarrow (6). Ex: Why does this step hold?

Applications of the compactness theorem

Can use the theorem to show models with specific properties exist.

Applications of the compactness theorem

Can use the theorem to show models with specific properties exist.

Lemma

Let F be a σ -sentence over some signature σ such that F has a model \mathcal{A}_n with $|U_{\mathcal{A}_n}| = n$ for every $n > 1$. Then, F has a model with an infinite universe.

Applications of the compactness theorem

Can use the theorem to show models with specific properties exist.

Lemma

Let F be a σ -sentence over some signature σ such that F has a model \mathcal{A}_n with $|U_{\mathcal{A}_n}| = n$ for every $n > 1$. Then, F has a model with an infinite universe.

Proof: Introduce a fresh binary predicate R . For $n > 1$, define

$$G_n = \forall x \neg R(x, x) \wedge \exists x_1 \dots \exists x_n \bigwedge_{1 \leq i < j \leq n} R(x_i, x_j).$$

Then, $B \models G_n$ implies $|U_B| \geq n$.

Ex1: Using the G_n and the compactness theorem, complete the proof.

Applications of the compactness theorem

Can use the theorem to show models with specific properties exist.

Lemma

Let F be a σ -sentence over some signature σ such that F has a model \mathcal{A}_n with $|U_{\mathcal{A}_n}| = n$ for every $n > 1$. Then, F has a model with an infinite universe.

Proof: Introduce a fresh binary predicate R . For $n > 1$, define

$$G_n = \forall x \neg R(x, x) \wedge \exists x_1 \dots \exists x_n \bigwedge_{1 \leq i < j \leq n} R(x_i, x_j).$$

Then, $B \models G_n$ implies $|U_B| \geq n$.

Ex1: Using the G_n and the compactness theorem, complete the proof.

Let

$$F_n := F \wedge G_n \quad \text{and} \quad \mathcal{S} := \bigcup_{n>1} \{F_n\}.$$

Every finite subset of \mathcal{S} is satisfiable. (Ex2: Why?)

Applications of the compactness theorem

Can use the theorem to show models with specific properties exist.

Lemma

Let F be a σ -sentence over some signature σ such that F has a model \mathcal{A}_n with $|U_{\mathcal{A}_n}| = n$ for every $n > 1$. Then, F has a model with an infinite universe.

Proof: Introduce a fresh binary predicate R . For $n > 1$, define

$$G_n = \forall x \neg R(x, x) \wedge \exists x_1 \dots \exists x_n \bigwedge_{1 \leq i < j \leq n} R(x_i, x_j).$$

Then, $B \models G_n$ implies $|U_B| \geq n$.

Ex1: Using the G_n and the compactness theorem, complete the proof.

Let

$$F_n := F \wedge G_n \quad \text{and} \quad \mathcal{S} := \bigcup_{n>1} \{F_n\}.$$

Every finite subset of \mathcal{S} is satisfiable. (Ex2: Why?) By compactness, \mathcal{S} has a model \mathcal{B} . Then, $|U_{\mathcal{B}}|$ is infinite, and $\mathcal{B} \models F$. (Ex3: Why?)

Peano axioms

Let $\sigma = \langle 0, s, +, \cdot, = \rangle$ be the sig. of arithmetic. Can we find a possibly infinite set of σ -formulas whose only model up to isomorphism is the classical arithmetic, i.e. the standard structure on \mathbb{N} ?

Peano axioms

Let $\sigma = \langle 0, s, +, \cdot, = \rangle$ be the sig. of arithmetic. Can we find a possibly infinite set of σ -formulas whose only model up to isomorphism is the classical arithmetic, i.e. the standard structure on \mathbb{N} ?



Figure: Giuseppe Peano (1858 - 1932)

Peano axioms

Let $\sigma = \langle 0, s, +, \cdot, = \rangle$. Peano axioms axiomatise elementary facts about arithmetic in first-order logic with equality:

$$\begin{array}{ll} \forall x \neg (s(x) = 0), & \forall x \forall y (s(x) = s(y) \rightarrow x = y), \\ \forall x (x + 0 = x), & \forall x \forall y (x + s(y) = s(x + y)), \\ \forall x (x \cdot 0 = 0), & \forall x \forall y (x \cdot s(y) = (x \cdot y) + x). \end{array}$$

Peano axioms

Let $\sigma = \langle 0, s, +, \cdot, = \rangle$. Peano axioms axiomatise elementary facts about arithmetic in first-order logic with equality:

$$\begin{array}{ll} \forall x \neg(s(x) = 0), & \forall x \forall y (s(x) = s(y) \rightarrow x = y), \\ \forall x (x + 0 = x), & \forall x \forall y (x + s(y) = s(x + y)), \\ \forall x (x \cdot 0 = 0), & \forall x \forall y (x \cdot s(y) = (x \cdot y) + x). \end{array}$$

Induction over natural numbers:

$$\forall Y ((0 \in Y \wedge \forall x (x \in Y \rightarrow s(x) \in Y)) \rightarrow \forall x (x \in Y)).$$

Peano axioms

Let $\sigma = \langle 0, s, +, \cdot, = \rangle$. Peano axioms axiomatise elementary facts about arithmetic in first-order logic with equality:

$$\begin{array}{ll} \forall x \neg (s(x) = 0), & \forall x \forall y (s(x) = s(y) \rightarrow x = y), \\ \forall x (x + 0 = x), & \forall x \forall y (x + s(y) = s(x + y)), \\ \forall x (x \cdot 0 = 0), & \forall x \forall y (x \cdot s(y) = (x \cdot y) + x). \end{array}$$

Induction over natural numbers:

$$\forall Y ((0 \in Y \wedge \forall x (x \in Y \rightarrow s(x) \in Y)) \rightarrow \forall x (x \in Y)).$$

But, cannot quantify over sets in first-order logic. We instead resort to infinitely many axioms, one for each formula $\phi(x, y_1, \dots, y_k)$:

$$\forall y_1 \dots y_k \left(\left(\phi(0) \wedge \forall x (\phi(x) \rightarrow \phi(s(x))) \right) \rightarrow \forall x \phi(x) \right).$$

Peano axioms

Let $\sigma = \langle 0, s, +, \cdot, = \rangle$. Peano axioms axiomatise elementary facts about arithmetic in first-order logic with equality:

$$\begin{array}{ll} \forall x \neg (s(x) = 0), & \forall x \forall y (s(x) = s(y) \rightarrow x = y), \\ \forall x (x + 0 = x), & \forall x \forall y (x + s(y) = s(x + y)), \\ \forall x (x \cdot 0 = 0), & \forall x \forall y (x \cdot s(y) = (x \cdot y) + x). \end{array}$$

Induction over natural numbers:

$$\forall Y ((0 \in Y \wedge \forall x (x \in Y \rightarrow s(x) \in Y)) \rightarrow \forall x (x \in Y)).$$

But, cannot quantify over sets in first-order logic. We instead resort to infinitely many axioms, one for each formula $\phi(x, y_1, \dots, y_k)$:

$$\forall y_1 \dots y_k \left(\left(\phi(0) \wedge \forall x (\phi(x) \rightarrow \phi(s(x))) \right) \rightarrow \forall x \phi(x) \right).$$

Let S_{PA} be the union of all formulas above. Then, “classical arithmetic” is a model of S_{PA} .

Non-standard models of arithmetic

Theorem (without proof)

*The compactness theorem holds for first-order logic **with equality**.*

Ex1: Using the theorem, show that S_{PA} has a model not isomorphic to the classic arithmetic.

Non-standard models of arithmetic

Theorem (without proof)

*The compactness theorem holds for first-order logic **with equality**.*

Ex1: Using the theorem, show that S_{PA} has a model not isomorphic to the classic arithmetic.

Answer1: Introduce a new constant symbol c , and set

$$\mathcal{C} = \{\neg(c = s^i(0)) : i \in \mathbb{N}\}.$$

Then, every finite subset of $S_{PA} \cup \mathcal{C}$ is satisfiable. (Ex2: Why?)

Non-standard models of arithmetic

Theorem (without proof)

*The compactness theorem holds for first-order logic **with equality**.*

Ex1: Using the theorem, show that \mathcal{S}_{PA} has a model not isomorphic to the classic arithmetic.

Answer1: Introduce a new constant symbol c , and set

$$\mathcal{C} = \{\neg(c = s^i(0)) : i \in \mathbb{N}\}.$$

Then, every finite subset of $\mathcal{S}_{PA} \cup \mathcal{C}$ is satisfiable. (Ex2: Why?)

Thus, $\mathcal{S}_{PA} \cup \mathcal{C}$ has a model \mathcal{A} by the compactness theorem.

The model \mathcal{A} is not isomorphic to the “classical” model of arithmetic.
(Ex3: Why?)

Non-standard models of arithmetic

Theorem (without proof)

*The compactness theorem holds for first-order logic **with equality**.*

Ex1: Using the theorem, show that \mathcal{S}_{PA} has a model not isomorphic to the classic arithmetic.

Answer1: Introduce a new constant symbol c , and set

$$\mathcal{C} = \{\neg(c = s^i(0)) : i \in \mathbb{N}\}.$$

Then, every finite subset of $\mathcal{S}_{PA} \cup \mathcal{C}$ is satisfiable. (Ex2: Why?)

Thus, $\mathcal{S}_{PA} \cup \mathcal{C}$ has a model \mathcal{A} by the compactness theorem.

The model \mathcal{A} is not isomorphic to the “classical” model of arithmetic. (Ex3: Why?)

Answer3: Because $c_{\mathcal{A}} \neq s_{\mathcal{A}}^i(0_{\mathcal{A}})$ for all $i \in \mathbb{N}$.

Upward Löwenheim-Skolem theorem

Ex: What if we add more axioms to S_{PA} ? Can we axiomatise the classic arithmetic then?

Upward Löwenheim-Skolem theorem

Ex: What if we add more axioms to S_{PA} ? Can we axiomatise the classic arithmetic then?

Answer: No. Because the following theorem says that we can make an arbitrary big model.

Theorem (Upward Löwenheim-Skolem theorem)

If a set of formulas S over a finite signature σ has an infinite model \mathcal{A} , then for any cardinal κ , it has a model \mathcal{B} with a universe of cardinality κ .

Downward Löwenheim-Skolem theorem

Let σ be the signature

$$\langle 0, 1, +, \cdot, <, = \rangle.$$

Let \mathcal{S} be the set of first-order σ -sentences that holds for \mathbb{R} .

Ex: Can \mathcal{S} have a model with a countable universe?

Downward Löwenheim-Skolem theorem

Let σ be the signature

$$\langle 0, 1, +, \cdot, <, = \rangle.$$

Let \mathcal{S} be the set of first-order σ -sentences that holds for \mathbb{R} .

Ex: Can \mathcal{S} have a model with a countable universe?

Answer: Yes. Because of the following theorem.

Theorem (Downward Löwenheim-Skolem theorem)

If a set of formulas \mathcal{S} over a finite signature σ has an infinite model \mathcal{A} , then it has a model \mathcal{B} with a countable universe.

Downward Löwenheim-Skolem theorem

Let σ be the signature

$$\langle 0, 1, +, \cdot, <, = \rangle.$$

Let \mathcal{S} be the set of first-order σ -sentences that holds for \mathbb{R} .

Ex: Can \mathcal{S} have a model with a countable universe?

Answer: Yes. Because of the following theorem.

Theorem (Downward Löwenheim-Skolem theorem)

If a set of formulas \mathcal{S} over a finite signature σ has an infinite model \mathcal{A} , then it has a model \mathcal{B} with a countable universe.

Corollary

\mathcal{S} has a model with a countable universe but not isomorphic to the classic arithmetic.