

# Lecture 12

## Resolution for first-order logic

Unification, resolution

*Introduction to Logic for Computer Science*

Prof Hongseok Yang  
KAIST

These slides are minor variants of those made by Prof Worrell and Dr Haase for their logic course at Oxford.

## Drawbacks of ground resolution

Ground resolution good for showing semi-decidability, but bad for practical purposes.

Requires “looking ahead” to see which ground terms will be needed.

Want to instantiate ground terms “by need.”

## Topic of this lecture

First-order-logic version of resolution.

Uses so called unification.

Supports by-need grounding.

Forms basis of programming language Prolog.

## Substitution

Used to replace variables by  $\sigma$ -terms.

More generally, a substitution is a function  $\theta$  mapping  $\sigma$ -terms to  $\sigma$ -terms such that

$$\begin{aligned}c\theta &= c, \\ f(t_1, \dots, t_k)\theta &= f(t_1\theta, \dots, t_k\theta).\end{aligned}$$

## Substitution

Used to replace variables by  $\sigma$ -terms.

More generally, a substitution is a function  $\theta$  mapping  $\sigma$ -terms to  $\sigma$ -terms such that

$$\begin{aligned}c\theta &= c, \\ f(t_1, \dots, t_k)\theta &= f(t_1\theta, \dots, t_k\theta).\end{aligned}$$

Ex1: Show that if  $\theta$  and  $\theta'$  work the same for variables, they are equal.

## Substitution

Used to replace variables by  $\sigma$ -terms.

More generally, a substitution is a function  $\theta$  mapping  $\sigma$ -terms to  $\sigma$ -terms such that

$$\begin{aligned}c\theta &= c, \\ f(t_1, \dots, t_k)\theta &= f(t_1\theta, \dots, t_k\theta).\end{aligned}$$

Ex1: Show that if  $\theta$  and  $\theta'$  work the same for variables, they are equal.

Denote by  $\theta \cdot \theta'$  substitution first performing  $\theta$  and then  $\theta'$ .

## Substitution

Used to replace variables by  $\sigma$ -terms.

More generally, a substitution is a function  $\theta$  mapping  $\sigma$ -terms to  $\sigma$ -terms such that

$$\begin{aligned}c\theta &= c, \\f(t_1, \dots, t_k)\theta &= f(t_1\theta, \dots, t_k\theta).\end{aligned}$$

Ex1: Show that if  $\theta$  and  $\theta'$  work the same for variables, they are equal.

Denote by  $\theta \cdot \theta'$  substitution first performing  $\theta$  and then  $\theta'$ .

Ex2: Extend  $\theta$  to arbitrary formulas in first-order logic.

Ex3: Let  $\theta = [f(y)/x]$ ,  $\theta' = [g(c, z)/y]$ , and  $P(x, c)$  be an atomic formula. Compute the following three:

$$P(x, c)\theta, \quad \theta \cdot \theta', \quad P(x, c)(\theta \cdot \theta').$$

Ex4: Do you think  $F(\theta \cdot \theta') = (F\theta)\theta'$  always?

## Unifier and most general unifier

For a set of literals  $D = \{L_1, \dots, L_k\}$ , define  $D\theta := \{L_1\theta, \dots, L_k\theta\}$ .

$\theta$  **unifies**  $D$  if  $D\theta = \{L\}$  for some literal  $L$ .



## Unifier and most general unifier

For a set of literals  $D = \{L_1, \dots, L_k\}$ , define  $D\theta := \{L_1\theta, \dots, L_k\theta\}$ .

$\theta$  **unifies**  $D$  if  $D\theta = \{L\}$  for some literal  $L$ .

### Example

We have that  $\theta = [f(a)/x][a/y]$  unifies  $\{P(x), P(f(y))\}$  since

$$\{P(x)\theta, P(f(y))\theta\} = \{P(f(a)), P(f(a))\} = \{P(f(a))\},$$

but  $\theta' = [f(y)/x]$  is also unifier.

## Unifier and most general unifier

For a set of literals  $D = \{L_1, \dots, L_k\}$ , define  $D\theta := \{L_1\theta, \dots, L_k\theta\}$ .

$\theta$  **unifies**  $D$  if  $D\theta = \{L\}$  for some literal  $L$ .

### Example

We have that  $\theta = [f(a)/x][a/y]$  unifies  $\{P(x), P(f(y))\}$  since

$$\{P(x)\theta, P(f(y))\theta\} = \{P(f(a)), P(f(a))\} = \{P(f(a))\},$$

but  $\theta' = [f(y)/x]$  is also unifier. Note that  $\theta = \theta' \cdot [a/y]$ .

## Unifier and most general unifier

For a set of literals  $D = \{L_1, \dots, L_k\}$ , define  $D\theta := \{L_1\theta, \dots, L_k\theta\}$ .

$\theta$  **unifies**  $D$  if  $D\theta = \{L\}$  for some literal  $L$ .

### Example

We have that  $\theta = [f(a)/x][a/y]$  unifies  $\{P(x), P(f(y))\}$  since

$$\{P(x)\theta, P(f(y))\theta\} = \{P(f(a)), P(f(a))\} = \{P(f(a))\},$$

but  $\theta' = [f(y)/x]$  is also unifier. Note that  $\theta = \theta' \cdot [a/y]$ .

### Definition

We call  $\theta$  a **most general unifier (mgu)** of  $D$  if  $\theta$  is a unifier and for all other unifiers  $\theta'$  there is a substitution  $\theta''$  such that  $\theta' = \theta \cdot \theta''$ .

## Unifier and most general unifier

For a set of literals  $D = \{L_1, \dots, L_k\}$ , define  $D\theta := \{L_1\theta, \dots, L_k\theta\}$ .

$\theta$  **unifies**  $D$  if  $D\theta = \{L\}$  for some literal  $L$ .

### Example

We have that  $\theta = [f(a)/x][a/y]$  unifies  $\{P(x), P(f(y))\}$  since

$$\{P(x)\theta, P(f(y))\theta\} = \{P(f(a)), P(f(a))\} = \{P(f(a))\},$$

but  $\theta' = [f(y)/x]$  is also unifier. Note that  $\theta = \theta' \cdot [a/y]$ .

### Definition

We call  $\theta$  a **most general unifier (mgu)** of  $D$  if  $\theta$  is a unifier and for all other unifiers  $\theta'$  there is a substitution  $\theta''$  such that  $\theta' = \theta \cdot \theta''$ .

Ex1: Is an mgu unique?

Ex2: When doesn't it exist? Find examples.

Ex3: Assume that  $D$  is unifiable. Does an mgu exist in this case?

### Theorem (Unification Theorem)

*A unifiable set of literals  $D$  has a most general unifier.*

### Theorem (Unification Theorem)

*A unifiable set of literals  $D$  has a most general unifier.*

Proof by an algorithm.

## Theorem (Unification Theorem)

*A unifiable set of literals  $D$  has a most general unifier.*

Proof by an algorithm.

### Unification Algorithm

**Input:** Set of literals  $D$

**Output:** Either a most general unifier  $\theta$  of  $D$  or “fail”

$\theta :=$  identity substitution

**while**  $\theta$  is not a unifier of  $D$  **do**

    pick two distinct literals in  $D\theta$  and

        find the left-most positions at which they differ

**if** one of the corresponding sub-terms is variable  $x$  and  
        the other term  $t$  does not contain  $x$

**then**  $\theta := \theta \cdot [t/x]$  **else** output “fail” and halt

**od**

output  $\theta$

## Theorem (Unification Theorem)

*A unifiable set of literals  $D$  has a most general unifier.*

Proof by an algorithm.

### Unification Algorithm

**Input:** Set of literals  $D$

**Output:** Either a most general unifier  $\theta$  of  $D$  or “fail”

$\theta :=$  identity substitution

**while**  $\theta$  is not a unifier of  $D$  **do**

    pick two distinct literals in  $D\theta$  and

        find the left-most positions at which they differ

**if** one of the corresponding sub-terms is variable  $x$  and  
        the other term  $t$  does not contain  $x$

**then**  $\theta := \theta \cdot [t/x]$  **else** output “fail” and halt

**od**

output  $\theta$

Ex1: Run the algo. for  $\{P(x), P(f(y))\}$  and  $\{P(x, y), P(f(z), x)\}$ .

Ex2: Why correct? Why terminate? Loop invariant?



## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur.

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ .

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ . Ex: Complete the rest.

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ . Ex: Complete the rest.
- Since  $\theta'$  unifies  $D$  and  $\theta' = \theta \cdot \theta'$  by ind. hypo., we have  $t\theta' = x\theta'$ .

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ . Ex: Complete the rest.
- Since  $\theta'$  unifies  $D$  and  $\theta' = \theta \cdot \theta'$  by ind. hypo., we have  $t\theta' = x\theta'$ .
- This implies  $\theta' = [t/x] \cdot \theta'$ .

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ . Ex: Complete the rest.
- Since  $\theta'$  unifies  $D$  and  $\theta' = \theta \cdot \theta'$  by ind. hypo., we have  $t\theta' = x\theta'$ .
- This implies  $\theta' = [t/x] \cdot \theta'$ .
- Thus,  $\theta' = \theta \cdot \theta' = \theta \cdot ([t/x] \cdot \theta') = (\theta \cdot [t/x]) \cdot \theta'$ .



## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ . Ex: Complete the rest.
- Since  $\theta'$  unifies  $D$  and  $\theta' = \theta \cdot \theta'$  by ind. hypo., we have  $t\theta' = x\theta'$ .
- This implies  $\theta' = [t/x] \cdot \theta'$ .
- Thus,  $\theta' = \theta \cdot \theta' = \theta \cdot ([t/x] \cdot \theta') = (\theta \cdot [t/x]) \cdot \theta'$ .

Ex: Use the invariant and prove the theorem.

## Proof of unification theorem

Termination: if we do not halt, a variable  $x$  gets replaced by a term in which  $x$  does not occur. Ex: Why does this imply termination?

Loop invariant: for any unifier  $\theta'$  of  $D$ , have  $\theta' = \theta \cdot \theta'$ .

- Clearly holds before entering loop.
- Suppose the algo. finds  $x$  and  $t$  in  $D\theta$ . Ex: Complete the rest.
- Since  $\theta'$  unifies  $D$  and  $\theta' = \theta \cdot \theta'$  by ind. hypo., we have  $t\theta' = x\theta'$ .
- This implies  $\theta' = [t/x] \cdot \theta'$ .
- Thus,  $\theta' = \theta \cdot \theta' = \theta \cdot ([t/x] \cdot \theta') = (\theta \cdot [t/x]) \cdot \theta'$ .

Ex: Use the invariant and prove the theorem.

At abnormal termination, the loop inv. implies that  $D$  is not unifiable.

At normal termination,  $\theta$  is a unifier. The loop inv. implies  $\theta$  is an mgu.

## First-order-logic resolution

For a set of literals  $D$ ,  $\overline{D}$  denotes the complements of all literals in  $D$ .

## First-order-logic resolution

For a set of literals  $D$ ,  $\overline{D}$  denotes the complements of all literals in  $D$ .

### Definition (First-order-logic resolution)

Let  $C_1, C_2$  be clauses with no variables in common.

$R$  is a **resolvent** of  $C_1$  and  $C_2$  if there are  $D_1 \subseteq C_1$  and  $D_2 \subseteq C_2$  s.t.

- $D_1 \cup \overline{D_2}$  has an mgu  $\theta$ , and
- $R = (C_1\theta \setminus \{L\}) \cup (C_2\theta \setminus \{\overline{L}\})$  with  $L = D_1\theta$  and  $\overline{L} = D_2\theta$ .

## First-order-logic resolution

For a set of literals  $D$ ,  $\overline{D}$  denotes the complements of all literals in  $D$ .

### Definition (First-order-logic resolution)

Let  $C_1, C_2$  be clauses with no variables in common.

$R$  is a **resolvent** of  $C_1$  and  $C_2$  if there are  $D_1 \subseteq C_1$  and  $D_2 \subseteq C_2$  s.t.

- $D_1 \cup \overline{D_2}$  has an mgu  $\theta$ , and
- $R = (C_1\theta \setminus \{L\}) \cup (C_2\theta \setminus \{\overline{L}\})$  with  $L = D_1\theta$  and  $\overline{L} = D_2\theta$ .

If  $C_1, C_2$  have variables in common,  $R$  is a **resolvent** if there are renamings  $\theta_1, \theta_2$  s.t.

- $C_1\theta_1, C_2\theta_2$  have no variables in common, and
- $R$  is a resolvent of  $C_1\theta_1$  and  $C_2\theta_2$ .

## Example


Ex: Given the signature with constant symbol  $e$ , unary function symbols  $f$  and  $g$ , and ternary predicate symbol  $P$ , compute a resolvent of

$$C_1 = \{\neg P(f(e), x, f(g(e)))\} \text{ and } C_2 = \{\neg P(x, y, z), P(f(x), y, f(z))\}.$$

## Example

Ex: Given the signature with constant symbol  $e$ , unary function symbols  $f$  and  $g$ , and ternary predicate symbol  $P$ , compute a resolvent of

$$C_1 = \{\neg P(f(e), x, f(g(e)))\} \text{ and } C_2 = \{\neg P(x, y, z), P(f(x), y, f(z))\}.$$



**Figure:** First-order-logic resolution example.

## Resolution derivation

Use resolution in order to derive a clause  $C$  from a set of clauses  $F$ .

A **derivation** of  $C$  is a sequence of clauses  $C_1, \dots, C_m$  such that

- $C = C_m$ ; and
- each  $C_i$  is either a clause from  $F$  possibly with variable renaming or obtained from resolution of  $C_j$  and  $C_k$  for some  $j, k < i$ .

$\text{Res}^*(F)$  is the set of all clauses derivable from  $F$ .



## Putting it all together

Let

$$F_1 = \forall x A(e, x, x),$$

$$F_2 = \forall x \forall y \forall z (\neg A(x, y, z) \vee A(s(x), y, s(z))),$$

$$F_3 = \forall x \exists y A(s(s(e)), x, y).$$

Ex: Prove that  $F_1 \wedge F_2 \models F_3$ , i.e.  $F_1 \wedge F_2 \wedge \neg F_3$  is unsatisfiable. Use Skolemisation and first-order-logic resolution.

## Putting it all together

$$F_1 = \forall x A(e, x, x),$$

$$F_2 = \forall x \forall y \forall z (\neg A(x, y, z) \vee A(s(x), y, s(z))),$$

$$F_3 = \forall x \exists y A(s(s(e)), x, y).$$

Step 1: Skolemise each  $F_i$  separately.

$$\neg F_3 = \exists y \forall z \neg A(s(s(e)), y, z) \rightsquigarrow G_3 := \forall z \neg A(s(s(e)), c, z)$$

Step 2: Use resolution to derive the empty clause.

- |  |  |
|--|--|
| 1. $\{\neg A(s(s(e)), c, z_1)\}$                       | clause of $G_3$  |
| 2. $\{\neg A(x_2, y_2, z_2), A(s(x_2), y_2, s(z_2))\}$ | clause of $F_2$  |
| 3. $\{\neg A(s(e), c, z_3)\}$                          | 1,2 Res. with sub.<br>$[s(e)/x_2][c/y_2][s(z_2)/z_1][z_3/z_2]$ |
| 4. $\{\neg A(e, c, z_4)\}$                             | 2,3 Res. with sub.<br>$[e/x_2][c/y_2][s(z_2)/z_3][z_4/z_2]$    |
| 5. $\{A(e, y_5, y_5)\}$                                | clause of $F_1$  |
| 6. $\square$   | 4,5 Res. with sub. $[c/y_5][c/z_4]$                            |

### Lemma (Resolution Lemma)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>*

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

### Lemma (Resolution Lemma)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>*

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

Ex: Prove the lemma.

### Lemma (Resolution Lemma)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>*

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

Ex: Prove the lemma.

### Proof.

Clearly  $\forall^*(G \cup \{R\}) \models F$ .

### Lemma (Resolution Lemma)

Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

Ex: Prove the lemma.

### Proof.

Clearly  $\forall^*(G \cup \{R\}) \models F$ .

For converse direction, show that  $F \models R$ . Suppose  $R$  is a resolvent of clauses  $C_1, C_2 \in G$ , with  $R = (C_1\theta \setminus \{L\}) \cup (C_2\theta' \setminus \{\bar{L}\})$  for some substitutions  $\theta, \theta'$  and complementary literals  $L \in C_1\theta$  and  $\bar{L} \in C_2\theta'$ .

### Lemma (Resolution Lemma)

Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

Ex: Prove the lemma.

### Proof.

Clearly  $\forall^*(G \cup \{R\}) \models F$ .

For converse direction, show that  $F \models R$ . Suppose  $R$  is a resolvent of clauses  $C_1, C_2 \in G$ , with  $R = (C_1\theta \setminus \{L\}) \cup (C_2\theta' \setminus \{\bar{L}\})$  for some substitutions  $\theta, \theta'$  and complementary literals  $L \in C_1\theta$  and  $\bar{L} \in C_2\theta'$ .

Let  $\mathcal{A}$  be an assignment that satisfies  $F = \forall^* G$ . Since  $C_1, C_2 \in G$ , by the Translation Lemma  $\mathcal{A} \models C_1\theta$  and  $\mathcal{A} \models C_2\theta'$ .

### Lemma (Resolution Lemma)

Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

Ex: Prove the lemma.

### Proof.

Clearly  $\forall^*(G \cup \{R\}) \models F$ .

For converse direction, show that  $F \models R$ . Suppose  $R$  is a resolvent of clauses  $C_1, C_2 \in G$ , with  $R = (C_1\theta \setminus \{L\}) \cup (C_2\theta' \setminus \{\bar{L}\})$  for some substitutions  $\theta, \theta'$  and complementary literals  $L \in C_1\theta$  and  $\bar{L} \in C_2\theta'$ .

Let  $\mathcal{A}$  be an assignment that satisfies  $F = \forall^* G$ . Since  $C_1, C_2 \in G$ , by the Translation Lemma  $\mathcal{A} \models C_1\theta$  and  $\mathcal{A} \models C_2\theta'$ .

Moreover, since  $\mathcal{A}$  satisfies at most one of the complementary literals  $L$  and  $\bar{L}$ ,  $\mathcal{A}$  satisfies at least one of  $C_1\theta \setminus \{L\}$  and  $C_2\theta' \setminus \{\bar{L}\}$ .



### Lemma (Resolution Lemma)

Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

Ex: Prove the lemma.

### Proof.

Clearly  $\forall^*(G \cup \{R\}) \models F$ .

For converse direction, show that  $F \models R$ . Suppose  $R$  is a resolvent of clauses  $C_1, C_2 \in G$ , with  $R = (C_1\theta \setminus \{L\}) \cup (C_2\theta' \setminus \{\bar{L}\})$  for some substitutions  $\theta, \theta'$  and complementary literals  $L \in C_1\theta$  and  $\bar{L} \in C_2\theta'$ .

Let  $\mathcal{A}$  be an assignment that satisfies  $F = \forall^* G$ . Since  $C_1, C_2 \in G$ , by the Translation Lemma  $\mathcal{A} \models C_1\theta$  and  $\mathcal{A} \models C_2\theta'$ .

Moreover, since  $\mathcal{A}$  satisfies at most one of the complementary literals  $L$  and  $\bar{L}$ ,  $\mathcal{A}$  satisfies at least one of  $C_1\theta \setminus \{L\}$  and  $C_2\theta' \setminus \{\bar{L}\}$ .

Thus,  $\mathcal{A}$  satisfies  $R$ , as required. □

### Lemma (Resolution Lemma)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form, with  $G$  quantifier-free and CNF. Let  $R$  be a resolvent of two clauses in  $G$ . Then  $F \equiv \forall^*(G \cup \{R\})$ .<sup>a</sup>*

---

<sup>a</sup>For a first-order formula  $G'$ , we write  $\forall^* G'$  for  $\forall y_1 \dots \forall y_n G'$  where  $y_1, \dots, y_n$  are all the free variables of  $G'$ .

### Theorem (Soundness)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form with  $G$  quantifier-free and CNF. If there is a resolution derivation of  $\square$  from  $G$ , then  $F$  is unsatisfiable.*

Ex: Prove the theorem using the Resolution Lemma.

# Completeness

## Theorem (Completeness)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form with  $G$  quantifier-free and CNF. If  $F$  is unsatisfiable, there is a resolution derivation of  $\square$  from  $G$ .*

# Completeness

## Theorem (Completeness)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form with  $G$  quantifier-free and CNF. If  $F$  is unsatisfiable, there is a resolution derivation of  $\square$  from  $G$ .*

Ex1: Prove the Completeness Theorem.

# Completeness

## Theorem (Completeness)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form with  $G$  quantifier-free and CNF. If  $F$  is unsatisfiable, there is a resolution derivation of  $\square$  from  $G$ .*

Ex1: Prove the Completeness Theorem.

Hint: Use the Lifting Lemma below.

## Lemma (Lifting Lemma)

*Let  $C_1$  and  $C_2$  be clauses with respective ground instances  $D_1$  and  $D_2$ . Suppose that  $R$  is a propositional resolvent of  $D_1$  and  $D_2$ . Then,  $C_1$  and  $C_2$  have a first-order-logic resolvent  $R'$  such that  $R$  is a ground instance of  $R'$ .*

# Completeness

## Theorem (Completeness)

*Let  $F = \forall x_1 \dots \forall x_n G$  be a closed formula in Skolem form with  $G$  quantifier-free and CNF. If  $F$  is unsatisfiable, there is a resolution derivation of  $\square$  from  $G$ .*

Ex1: Prove the Completeness Theorem.

Hint: Use the Lifting Lemma below.

## Lemma (Lifting Lemma)

*Let  $C_1$  and  $C_2$  be clauses with respective ground instances  $D_1$  and  $D_2$ . Suppose that  $R$  is a propositional resolvent of  $D_1$  and  $D_2$ . Then,  $C_1$  and  $C_2$  have a first-order-logic resolvent  $R'$  such that  $R$  is a ground instance of  $R'$ .*

Ex2: Prove the Lifting Lemma.