

# Lecture 10

## Herbrand's theorem and ground resolution

*Introduction to Logic for Computer Science*

Prof Hongseok Yang  
KAIST

These slides are minor variants of those made by Prof Worrell and Dr Haase for their logic course at Oxford.

## Recap

Let  $F$  be a quantifier-free formula.

Prenex form:  $Q_1 x_1 Q_2 x_2 \cdots Q_n x_n F$ , where  $Q_i \in \{\forall, \exists\}$ .

Skolem form:  $\forall x_1 \forall x_2 \cdots \forall x_n F$ .

Every first-order formula can be translated into an equi-satisfiable formula in Skolem form in poly. time.

## Ground terms

### Definition

Given a signature  $\sigma$ , a **ground term** is a  $\sigma$ -term in which no variable symbol appears.

## Ground terms

### Definition

Given a signature  $\sigma$ , a **ground term** is a  $\sigma$ -term in which no variable symbol appears.

### Example

Let  $\sigma = \langle c, d, f, g, P, Q \rangle$  with unary  $f$  and binary  $g$ , the set of ground terms is

$$\{c, d, f(c), f(d), g(c, c), g(c, d), g(d, c), g(d, d), f(f(c)), f(f(d)), \dots\}.$$

## Ground terms

### Definition

Given a signature  $\sigma$ , a **ground term** is a  $\sigma$ -term in which no variable symbol appears.

### Example

Let  $\sigma = \langle c, d, f, g, P, Q \rangle$  with unary  $f$  and binary  $g$ , the set of ground terms is

$$\{c, d, f(c), f(d), g(c, c), g(c, d), g(d, c), g(d, d), f(f(c)), f(f(d)), \dots\}.$$

Ex: Find a signature  $\sigma$  such that its set of ground terms is isomorphic to  $\mathbb{N}$ .

## Herbrand structures

### Definition

Let  $\sigma$  be a signature with at least one constant symbol. A  $\sigma$ -structure  $\mathcal{H}$  is a **Herbrand structure** if the following hold:

- The universe  $U_{\mathcal{H}}$  is the set of ground terms over  $\sigma$ .
- For every constant symbol  $c$ , we have  $c_{\mathcal{H}} = c$ .
- For every function symbol  $f$ ,  $f_{\mathcal{H}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ .

## Herbrand structures

### Definition

Let  $\sigma$  be a signature with at least one constant symbol. A  $\sigma$ -structure  $\mathcal{H}$  is a **Herbrand structure** if the following hold:

- The universe  $U_{\mathcal{H}}$  is the set of ground terms over  $\sigma$ .
- For every constant symbol  $c$ , we have  $c_{\mathcal{H}} = c$ .
- For every function symbol  $f$ ,  $f_{\mathcal{H}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ .

**Proposition:** For any Herbrand structure  $\mathcal{H}$  and ground term  $t$ ,  $\mathcal{H}(t) = t$ .

## Herbrand structures

### Definition

Let  $\sigma$  be a signature with at least one constant symbol. A  $\sigma$ -structure  $\mathcal{H}$  is a **Herbrand structure** if the following hold:

- The universe  $U_{\mathcal{H}}$  is the set of ground terms over  $\sigma$ .
- For every constant symbol  $c$ , we have  $c_{\mathcal{H}} = c$ .
- For every function symbol  $f$ ,  $f_{\mathcal{H}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ .

**Proposition:** For any Herbrand structure  $\mathcal{H}$  and ground term  $t$ ,  $\mathcal{H}(t) = t$ .

Ex1: Prove the proposition.



## Herbrand structures

### Definition

Let  $\sigma$  be a signature with at least one constant symbol. A  $\sigma$ -structure  $\mathcal{H}$  is a **Herbrand structure** if the following hold:

- The universe  $U_{\mathcal{H}}$  is the set of ground terms over  $\sigma$ .
- For every constant symbol  $c$ , we have  $c_{\mathcal{H}} = c$ .
- For every function symbol  $f$ ,  $f_{\mathcal{H}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ .

**Proposition:** For any Herbrand structure  $\mathcal{H}$  and ground term  $t$ ,  $\mathcal{H}(t) = t$ .

Ex1: Prove the proposition.

**Translation Lemma:** For all ground  $t$ ,  $\mathcal{H} \models F[t/x]$  iff  $\mathcal{H}_{[x \mapsto t]} \models F$ .

## Herbrand structures

### Definition

Let  $\sigma$  be a signature with at least one constant symbol. A  $\sigma$ -structure  $\mathcal{H}$  is a **Herbrand structure** if the following hold:

- The universe  $U_{\mathcal{H}}$  is the set of ground terms over  $\sigma$ .
- For every constant symbol  $c$ , we have  $c_{\mathcal{H}} = c$ .
- For every function symbol  $f$ ,  $f_{\mathcal{H}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ .

**Proposition:** For any Herbrand structure  $\mathcal{H}$  and ground term  $t$ ,  $\mathcal{H}(t) = t$ .

Ex1: Prove the proposition.

**Translation Lemma:** For all ground  $t$ ,  $\mathcal{H} \models F[t/x]$  iff  $\mathcal{H}_{[x \mapsto t]} \models F$ .

Ex2: Prove the lemma.

## Jaques Herbrand (1908 – 1931)



## Herbrand's theorem

### Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Herbrand's theorem

### Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

Ex: Prove the theorem.

## Herbrand's theorem

### Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

Ex: Prove the theorem.

Hint1: It is sufficient to consider closed  $F$  only.

## Herbrand's theorem

### Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

Ex: Prove the theorem.

Hint1: It is sufficient to consider closed  $F$  only.

Hint2: The proof of the only-if direction goes like this.

Suppose  $\mathcal{A} \models F$ .

## Herbrand's theorem

### Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

Ex: Prove the theorem.

Hint1: It is sufficient to consider closed  $F$  only.

Hint2: The proof of the only-if direction goes like this.

Suppose  $\mathcal{A} \models F$ . Define a Herbrand model  $\mathcal{H}$  by setting the interpretation of each predicate symbol  $P$  as follows:

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$



## Herbrand's theorem

### Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

Ex: Prove the theorem.

Hint1: It is sufficient to consider closed  $F$  only.

Hint2: The proof of the only-if direction goes like this.

Suppose  $\mathcal{A} \models F$ . Define a Herbrand model  $\mathcal{H}$  by setting the interpretation of each predicate symbol  $P$  as follows:

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

Now show that for all closed  $G$  in Skolem form, if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . Use induction on the number of  $\forall$  in  $G$ .

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

We show that for all *closed*  $G = \forall x_1 \forall x_2 \dots \forall x_m G^*$  with quantifier free  $G^*$ , if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . We use induction on  $m$ .

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

We show that for all *closed*  $G = \forall x_1 \forall x_2 \dots \forall x_m G^*$  with quantifier free  $G^*$ , if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . We use induction on  $m$ .

For  $m = 0$ ,  $G$  is closed. Thus, it is a Boolean combination of  $P(t_1, \dots, t_k)$ . By definition,  $\mathcal{H} \models G$ .

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

We show that for all *closed*  $G = \forall x_1 \forall x_2 \dots \forall x_m G^*$  with quantifier free  $G^*$ , if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . We use induction on  $m$ .

For  $m = 0$ ,  $G$  is closed. Thus, it is a Boolean combination of  $P(t_1, \dots, t_k)$ . By definition,  $\mathcal{H} \models G$ .

For  $m > 0$ , suppose  $\mathcal{A} \models \forall x G'$ . Note that  $x$  may appear free in  $G'$ .

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

We show that for all *closed*  $G = \forall x_1 \forall x_2 \dots \forall x_m G^*$  with quantifier free  $G^*$ , if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . We use induction on  $m$ .

For  $m = 0$ ,  $G$  is closed. Thus, it is a Boolean combination of  $P(t_1, \dots, t_k)$ . By definition,  $\mathcal{H} \models G$ .

For  $m > 0$ , suppose  $\mathcal{A} \models \forall x G'$ . Note that  $x$  may appear free in  $G'$ .

Translation Lemma gives  $\mathcal{A} \models G'[t/x]$  iff  $\mathcal{A}_{[x \mapsto \mathcal{A}(t)]} \models G'$  for all ground terms  $t$ .

## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

We show that for all *closed*  $G = \forall x_1 \forall x_2 \dots \forall x_m G^*$  with quantifier free  $G^*$ , if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . We use induction on  $m$ .

For  $m = 0$ ,  $G$  is closed. Thus, it is a Boolean combination of  $P(t_1, \dots, t_k)$ . By definition,  $\mathcal{H} \models G$ .

For  $m > 0$ , suppose  $\mathcal{A} \models \forall x G'$ . Note that  $x$  may appear free in  $G'$ .

Translation Lemma gives  $\mathcal{A} \models G'[t/x]$  iff  $\mathcal{A}_{[x \mapsto \mathcal{A}(t)]} \models G'$  for all ground terms  $t$ . So,  $\mathcal{H} \models G'[t/x]$  for all ground  $t$  by the ind. hypothesis.



## Theorem

*Let  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$  be a formula in Skolem form. Then  $F$  is satisfiable if and only if  $F$  has a Herbrand model.*

## Proof.

Assume that  $F$  is closed. Suppose  $\mathcal{A} \models F$ .

Define  $P_{\mathcal{H}}$  for each predicate symbol  $P$ :

$$(t_1, \dots, t_k) \in P_{\mathcal{H}} \text{ iff } \mathcal{A} \models P(t_1, \dots, t_k).$$

We show that for all *closed*  $G = \forall x_1 \forall x_2 \dots \forall x_m G^*$  with quantifier free  $G^*$ , if  $\mathcal{A} \models G$ , then  $\mathcal{H} \models G$ . We use induction on  $m$ .

For  $m = 0$ ,  $G$  is closed. Thus, it is a Boolean combination of  $P(t_1, \dots, t_k)$ . By definition,  $\mathcal{H} \models G$ .

For  $m > 0$ , suppose  $\mathcal{A} \models \forall x G'$ . Note that  $x$  may appear free in  $G'$ .

Translation Lemma gives  $\mathcal{A} \models G'[t/x]$  iff  $\mathcal{A}_{[x \mapsto \mathcal{A}(t)]} \models G'$  for all ground terms  $t$ . So,  $\mathcal{H} \models G'[t/x]$  for all ground  $t$  by the ind. hypothesis.

But then  $\mathcal{H} \models \forall x G'$ . □

## Ground resolution

**Herbrand expansion** of  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$ :

$$E(F) := \{F^*[t_1/x_1][t_2/x_2] \dots [t_n/x_n] : t_1, \dots, t_n \text{ ground terms}\}.$$

## Ground resolution

**Herbrand expansion** of  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$ :

$$E(F) := \{F^*[t_1/x_1][t_2/x_2] \dots [t_n/x_n] : t_1, \dots, t_n \text{ ground terms}\}.$$

### Theorem (Ground Resolution)

*A closed formula in Skolem form is unsatisfiable if and only if there is a propositional resolution proof of  $\square$  from  $E(F)$ .*

## Ground resolution

**Herbrand expansion** of  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$ :

$$E(F) := \{F^*[t_1/x_1][t_2/x_2] \dots [t_n/x_n] : t_1, \dots, t_n \text{ ground terms}\}.$$

### Theorem (Ground Resolution)

*A closed formula in Skolem form is unsatisfiable if and only if there is a propositional resolution proof of  $\square$  from  $E(F)$ .*

Ex1: What does this imply in terms of developing a satisfiability checker for first-order formulas?

## Ground resolution

**Herbrand expansion** of  $F = \forall x_1 \forall x_2 \dots \forall x_n F^*$ :

$$E(F) := \{F^*[t_1/x_1][t_2/x_2] \dots [t_n/x_n] : t_1, \dots, t_n \text{ ground terms}\}.$$

### Theorem (Ground Resolution)

*A closed formula in Skolem form is unsatisfiable if and only if there is a propositional resolution proof of  $\square$  from  $E(F)$ .*

Ex1: What does this imply in terms of developing a satisfiability checker for first-order formulas?

Ex2: Prove the theorem.

Hint: Prove that Herbrand' theorem implies the following theorem. Then, use the Compactness theorem for propositional logic.

### Theorem

*A closed formula  $F$  in Skolem form is satisfiable iff  $E(F)$  is satisfiable when considered as a set of propositional formulas.*

## Theorem

*A closed formula  $F$  in Skolem form is satisfiable iff  $E(F)$  is satisfiable when considered as a set of propositional formulas.*

## Proof.

By Herbrand's theorem,  $F$  is satisfiable if and only if  $F$  has a Herbrand model. Now

$$\begin{aligned}\mathcal{H} \models F &\text{ iff } \mathcal{H}_{[x_1 \mapsto t_1] \dots [x_n \mapsto t_n]} \models F^* \text{ for all ground terms } t_1, \dots, t_n \\ &\text{ iff } \mathcal{H} \models F^*[t_1/x_1] \dots [t_n/x_n] \text{ for all ground } t_i \text{ (by Trans. Lemma)} \\ &\text{ iff } \mathcal{H} \models E(F) \\ &\text{ iff } E(F) \text{ is satisfiable as prop. formula.}\end{aligned}$$



### Theorem (Ground Resolution)

*A closed formula in Skolem form is unsatisfiable if and only if there is a propositional resolution proof of  $\square$  from  $E(F)$ .*

### Proof.

By the Compactness theorem,  $E(F)$  is unsatisfiable if and only if some finite subset of  $E(F)$  is unsatisfiable. The latter happens if and only if  $\square$  can be derived from  $E(F)$  using resolution.  $\square$