STUDENT REFERENCE & INTERNAL USE NOTICE

This protocol documents standard in-vitro DPPH radical-scavenging assay procedures for estimating chemical antioxidant activity in HOPE Lab research and IS PROVIDED ONLY FOR HOPLE LAB MEMBERS for training and internal reference. External users should independently assess method suitability, verify current standards and chemical safety for reagents, and obtain required biosafety authorization and other essential approvals. This document does not replace an institutionally approved SOP or method-specific risk assessment.

TITLE

DPPH Radical Scavenging Assay for the Estimation of In-Vitro Antioxidant Activity

PRINCIPLE

The DPPH assay estimates the capacity of a test material to donate hydrogen atoms or electrons to reduce the stable organic nitrogen radical 1,1-diphenyl-2-picrylhydrazyl (DPPH•). In its radical state, DPPH exhibits a deep violet color with a strong visible absorption band centered near 517 nm. Upon reduction by an antioxidant compound or microbial metabolite (such as exopolysaccharides or cell-free extracts), the unpaired electron becomes paired, resulting in a stoichiometric decolorization from deep violet to pale yellow and a measurable decrease in absorbance at 517 nm.

Measured scavenging responses depend heavily on test-material concentration, solvent polarity, reaction pH, temperature, light exposure, optical path length, and incubation duration. Concentration-matched sample-only blanks and solvent-matched reagent controls are mandatory, particularly for pigmented or slightly turbid biological extracts. The DPPH assay is strictly an in-vitro chemical radical-scavenging test and does not establish antioxidant efficacy in living cells, animal models, humans, or complex food matrices.

Learning Outcomes

- Explain the chemical reduction mechanism and analytical boundaries of the DPPH free-radical assay.
- Distinguish the solvent-matched DPPH reagent control from concentration-specific sample-only blanks.
- Apply blank-corrected absorbance equations consistently without double-blanking errors.
- Evaluate concentration-response curves and apply valid interpolation rules for estimating IC_{50} values.
- Separate technical repeat measurements from independent biological replicate preparations.

OBJECTIVE & SCOPE

Objective: To standardize the experimental setup, blank-corrected calculation formulas, quality controls, and reporting conventions for estimating in-vitro DPPH radical-scavenging activity.

Scope: Applicable to characterized bacterial exopolysaccharides (EPS), cell-free culture supernatants, metabolite fractions, and pure reference compounds evaluated in compatible solvent systems. Samples that precipitate in alcoholic solvents or exhibit extreme turbidity require solvent compatibility validation. Results represent in-vitro chemical radical scavenging and must not be extrapolated to in-vivo biological antioxidant protection.

CORE CONCEPTS & CALCULATIONS

- **Blank-Corrected Scavenging Formula:** Define the three required absorbance measurements at 517 nm:
  • A_control: Absorbance of DPPH working solution mixed with the exact sample solvent (no test sample).
  • A_sample+reagent: Absorbance of the test sample mixed with DPPH working solution.
  • A_sample blank: Absorbance of the identical test-sample concentration mixed with pure solvent (no DPPH reagent), correcting for intrinsic sample color or scattering.
  DPPH Radical Scavenging (%) = [1 − (A_sample+reagent − A_sample blank) / A_control] × 100
  Ensure instrument zeroing (solvent baseline) is applied consistently so blank corrections are never subtracted twice.
- **Positive Antioxidant Reference:** Include a recognized standard antioxidant (e.g., Ascorbic Acid or Trolox) across a defined concentration series to confirm reagent reactivity and provide an internal benchmark.
- **Concentration-Response & IC_{50} Estimation:** IC_{50} represents the sample concentration achieving 50% radical scavenging under the stated assay conditions. Estimate IC_{50} via regression/interpolation only when empirical data points clearly bracket the 50% response threshold. Never extrapolate IC_{50} beyond the highest tested concentration; if 50% scavenging is not reached, report IC_{50} as 'not determined within the tested concentration range'.

MATERIALS & EQUIPMENT

Reagents & Consumables

- DPPH free-radical reagent (analytical grade, stored desiccated at -20°C in dark)
- Analytical-grade solvent system (e.g., absolute methanol or ethanol, or validated aqueous-alcohol matrix)
- Test samples (purified EPS solutions, fractions, or cell-free extracts across serial dilutions)
- Positive reference standard (e.g., L-Ascorbic Acid or Trolox freshly prepared)
- Amber glass bottles/foil-wrapped tubes & matched optical cuvettes or 96-well microplates

Equipment & Instrumentation

- Calibrated UV-Vis Spectrophotometer or Microplate Reader capable of reading at 517 nm
- Analytical Balance (0.1 mg readability) for accurate reagent and sample weighing
- Calibrated Micropipettes & Vortex Mixer
- Dark Incubation Chamber / Light-shielded Workstation at controlled temperature (e.g., 25°C)
- Precision Timer & Chemical Fume Hood

STEP-BY-STEP PROCEDURE

Step 1: Fresh Preparation of DPPH Working Solution

Weigh DPPH reagent accurately and dissolve in analytical-grade methanol or ethanol (typically 0.1–0.2 mM working concentration, as defined by the validated SOP) in an amber or foil-wrapped vessel. Mix thoroughly until completely dissolved. Verify that the initial absorbance of the DPPH control mixture at 517 nm falls within the instrument's optimal linear range (typically 0.70–1.00 AU). Prepare fresh daily and protect from light.

Step 2: Preparation of Test Sample Dilutions & Reference Standards

Prepare serial dilutions of the test material (e.g., 0.1 to 5.0 mg/mL for EPS or extracts) and positive reference standard (e.g., Ascorbic Acid, 1–100 µg/mL) in a compatible solvent. Ensure that the final solvent composition (e.g., water-to-methanol ratio) is identical across all sample reactions, sample blanks, and reagent controls.

Step 3: Reaction Setup (Samples, Sample Blanks & Reagent Controls)

In light-protected tubes or microplate wells, combine defined volumes of test sample and DPPH working solution (e.g., 1:1 v/v ratio, such as 1.0 mL sample + 1.0 mL DPPH, or 100 µL + 100 µL in microplates). Concurrently prepare: (a) Reagent Control (solvent + DPPH solution), and (b) Concentration-Matched Sample Blanks (each sample dilution + pure solvent without DPPH). Vortex or mix gently.

Step 4: Dark Incubation at Controlled Temperature

Incubate all reaction mixtures, blanks, and controls in complete darkness at a standardized, monitored temperature (e.g., 25 ± 2°C) for a fixed validated reaction endpoint (typically 30 minutes).

Step 5: Spectrophotometric Absorbance Measurement

Immediately at the 30-minute endpoint, measure the absorbance of each mixture at 517 nm against the appropriate solvent baseline. Record raw absorbance values for A_control, A_sample+reagent, and A_sample blank across all concentrations.

Step 6: Calculation, IC_{50} Determination & Replicate Reporting

Compute blank-corrected Scavenging (%) for each concentration using the standard equation. Plot scavenging percentage versus concentration across at least three independent biological replicates (with technical duplicates/triplicates). Determine IC_{50} only if experimental data bracket 50% scavenging.

CRITICAL FACTORS AFFECTING RESULTS

Variable	Effect of Inadequate Control	Required Control Standard
DPPH Reagent Stability	Thermal or photolytic decay lowers control absorbance	Store stock at -20°C; prepare working solution fresh & dark
Linear Absorbance Range	Initial A_control > 1.2 AU or < 0.4 AU distorts linearity	Adjust DPPH working conc. so A_control is ~0.70–1.00 at 517 nm
Light Exposure	Ambient light accelerates non-specific DPPH bleaching	Conduct preparation and 30 min incubation in total darkness
Sample Color & Turbidity	Intrinsic pigment or precipitation inflates/depresses OD_{517}	Include concentration-matched sample-only blanks for every level
Solvent Compatibility	High alcohol % can precipitate EPS or proteins causing scatter	Match final aqueous/organic solvent ratio across all tubes
Reaction Endpoint Timing	Slow-reacting antioxidants continue scavenging over time	Read absorbance at exact validated endpoint (e.g., 30 min)

QUALITY CONTROL & TROUBLESHOOTING

Quality Control (QC): Verify that A_control remains stable throughout the run and that the positive reference standard (e.g., Ascorbic Acid) achieves its expected dose-dependent scavenging response. Retain all raw absorbances, blank values, solvent ratios, and regression models.

Observation	Possible Explanation	Corrective Action
Positive standard shows weak response	Degraded DPPH stock, expired standard, or	Prepare fresh DPPH & ascorbic acid stocks;

	wrong wavelength	verify 517 nm
Calculated scavenging > 100%	Double-blanking error, sample precipitation, or unstable control	Audit blank subtraction math; check for sample turbidity/clumping
Negative scavenging values	Sample turbidity/precipitation upon mixing with methanol	Adjust aqueous/alcohol solvent ratio; verify sample solubility
50% scavenging not reached	Tested sample concentrations are below effective potency range	Do not extrapolate IC_{50}; report IC_{50} as 'not determined'
High technical variability	Pipetting error with volatile alcohol or evaporation	Use positive-displacement/pre-wetted tips; keep wells covered

BIOSAFETY & ADDITIONAL CONSIDERATIONS

- **Chemical Safety & Waste Disposal:** DPPH is a stable free radical and potential irritant; methanol and ethanol are flammable and toxic. Prepare reagent solutions inside a chemical fume hood wearing nitrile gloves and eye protection. Dispose of DPPH-organic solvent waste in designated hazardous chemical waste containers.
- **Scientific Interpretation Boundary:** Report DPPH scavenging strictly as an in-vitro chemical electron/hydrogen-transfer assay under the stated solvent, wavelength, and endpoint conditions. Do not interpret chemical radical scavenging alone as evidence of cellular, dietary, or clinical antioxidant efficacy.

REFERENCES & RELATED DOCUMENTS

Tarannum N, Ali F, Khan MS, Alhumaidan OS, Zawad ANMS, Hossain TJ. (2024). Bioactive exopolysaccharide from Limosilactobacillus fermentum LAB-1: antioxidant, anti-inflammatory, antibacterial and antibiofilm properties. Bioactive Carbohydrates and Dietary Fibre, 31, 100409.

Tarannum N, Hossain TJ, Ali F, Das T, Dhar K, Nafiz IH. (2023). Antioxidant, antimicrobial and emulsification properties of exopolysaccharides from lactic acid bacteria of bovine milk: insights from biochemical and genomic analysis. LWT - Food Science and Technology, 186, 115263.