Commutative Algebra for Algebraic Geometry

Bailey Arm

$\mathrm{May}\ 31,\ 2025$

Contents

1	Alg	gebraic Geometry, Hartshorne	9
1		ieties	9
	1.1	Affine Varieties	11
	1.2	Projective Varieties	11
	1.3	Morphisms	11
	1.4	Rational Maps	11
	1.5	Nonsingular Varieties	11
	1.6	Nonsingular Curves	11
	1.7	Intersections in Projective Space	11
2	Sch	emes	11
	2.1	Sheaves	11
	2.2	Schemes	11
	2.3	First Properties of Schemes	11
	2.4	Separated and Proper Morphisms	11
	2.5	Sheaves of Modules	11
	2.6	Divisors	11
	2.7	Projective Morphisms	11
	2.8	Differentials	11
	2.9	Formal Schemes	11
3	Coh	omology	11
Ü	3.1	Derived Functors	11
	3.2	Cohomology of Sheaves	11
	3.3	Cohomology of a Noetherian Affine Scheme	11
	3.4	Cech Cohomology	11
	3.5	The Cohomology of Projective Space	11
	3.6	Ext Groups and Sheaves	11
	3.7	The Serre Duality Theorem	11
	3.8	Higher Direct Images of Sheaves	11
	3.9	Flat Morphisms	11
		Smooth Morphisms	11
		The Theorem on Formal Functions	11
		The Semicontinuity Theorem	11
4	Cur		11
	4.1	Riemann-Roch Theorem	11
	4.2	Hurwitz's Theorem	11
	4.3	Embeddings in Projective Space	11
	4.4	Elliptic Curves	11

	4.5 4.6			11 11
5	Sur	faces	1	L1
	5.1			11
	5.2		· ·	11
	5.3			11
	5.4			11
	5.5			12
	5.6			12
	0.0	Classii	leading of Surfaces	-
II bı		ommut	tative Geometry with a View Toward Algebraic Geometry, Eisen-	13
6	Ras	ic Con	structions	L3
U	6.1			13
	0.1	6.1.1		13
		6.1.1	v	13
		6.1.2	v v	13
		6.1.4	· · · · · · · · · · · · · · · · · · ·	13
		6.1.4		13
		6.1.6		13
		6.1.0		13
		6.1.8	· · · · · · · · · · · · · · · · · · ·	13
			o de la companya de	13
		6.1.9	· ·	13
	6.0		V V 00	
	6.2			13
		6.2.1		13
		6.2.2		15
		6.2.3		15
		6.2.4		15
	0.0	6.2.5		15
	6.3		· · · · · · · · · · · · · · · · · · ·	15
		6.3.1		15
		6.3.2		15
		6.3.3	· · · · · · · · · · · · · · · · · · ·	15
		6.3.4		15
		6.3.5	· · · · · · · · · · · · · · · · · · ·	15
		6.3.6	v .	15
		6.3.7		15
		6.3.8		15
		6.3.9	· O	15
	6.4	_	•	15
		6.4.1		15
		6.4.2		15
		6.4.3	Normalisation in the Analytic Case	15
		6.4.4		15
		6.4.5		15
	6.5	Filtrat	ions and the Artin-Rees Lemma	15
		6.5.1	Associated Graded Rings and Modules	15
		6.5.2	The Blowup Algebra	15
		6.5.3	The Krull Intersection Theorem	15
		6.5.4	The Tangent Cone	15

	6.6	Flat Families	15
		6.6.1 Elementary Examples	15
		6.6.2 Introduction to Tor	15
		6.6.3 Criteria for Flatness	15
		6.6.4 The Local Criterion for Flatness	15
		6.6.5 The Rees Algebra	15
	6.7	Completions and Hensel's Lemma	15
		6.7.1 Examples and Definitions	15
		6.7.2 The Utility of Completions	15
		6.7.3 Lifting Idempotents	15
		6.7.4 Cohen Structure Theory and Coefficient Fields	15
		6.7.5 Basic Properties of Completion	15
		6.7.6 Maps from Power Series Rings	15
7	Dim	nension Theory	15
	7.1	Introduction to Dimension Theory	15
	7.2	Fundamental Definitions of Dimension Theory	15
	7.3		15
	7.4	Dimension and Codimension One	15
	7.5		15
	7.6		15
	7.7		15
	7.8		15
	7.9	Modules of Differentials	15
8	Hon	nological Methods	15
	8.1	<u> </u>	15
	8.2		15
	8.3		15
	8.4		15
	8.5		15
	T 0		
Η	1 C	Commutative Ring Theory, Matsumura	16
9			1 7
	9.1		17
	9.2		$\frac{17}{17}$
	9.3	Chain Conditions	17
10			17
		1 0	17
			17
	10.3	Associated Primes and Primary Decomposition	17
11			17
			17
			17
	11.3	Integral Extensions	17
12		O Company of the comp	17
			17
			17
	12.3	Krull Rings	17

13 Dimension Theory	17
13.1 Graded Rings, the Hilbert Function and the Samuel Function	17
13.2 Systems of Parameters and Multiplicity	17
13.3 The Dimension of Extension Rings	17
14 Regular Sequences	17
14.1 Regular Sequences and the Koszul Complex	17
14.2 Cohen-Macualay Rings	17
14.3 Gorenstein Rings	
<u></u>	
15 Regular Rings	17
15.1 Regular Rings	17
15.2 UFDs	
15.3 Complete Intersection Rings	
15.5 Complete intersection rungs	11
16 Flatness Revisited	17
16.1 The Local Flatness Criterion	
16.2 Flatness and Fibres	
16.3 Generic Freeness and Open Loci Results	17
17 Derivations	17
17.1 Derivations and Differentials	17
17.2 Separability	17
17.3 Higher Derivations	
18 I-Smoothness	17
18.1 <i>I</i> -Smoothness	17
18.2 The Structure Theorems for Complete Local Rings	
18.3 Connections with Derivations	
18.5 Connections with Derivations	11
19 Applications of Complete Local Rings	17
19.1 Chains of Prime Ideals	
19.2 The Formal Fibre	
19.3 Other Applications	17
IV Algebraic Geometry I: Schemes, Gortz-Wedhorn	18
V Algebraic Geometry II: Cohomology of Schemes, Gortz-Wedhorn	19
VI Sheaf Theory, Bredon	20
20 Sheaves and Presheaves	21
20.1 Definitions	21
20.2 Homomorphisms, Subsheaves and Quotient Sheaves	21
20.3 Direct and Inverse Images	
20.4 Cohomomorphisms	
20.5 Algebraic Constructions	
20.6 Supports	
20.7 Classical Cohomology Theories	21

21	Snear Conmology	21
	21.1 Differential Sheaves and Resolutions	21
	21.2 The Canonical Resolution and Sheaf Cohomology	21
	21.3 Injective Sheaves	21
	21.4 Acyclic Sheaves	21
	21.5 Flabby Sheaves	21
	21.6 Connected Sequences of Functors	21
		21
		21
		21
		21
	·	21
	• •	21
	₩	$\frac{-}{21}$
		21
	v	21
		21
		$\frac{21}{21}$
	· ·	21
		$\frac{21}{21}$
		21
	·	
	21.21The Steenrod Operations	21
22	Comparison with Other Cohomology Theories	21
		21
	0	21
	\mathbf{I}	21
	\circ	21
	22.4 Cech Cohomology	21
23	Applications of Spectral Sequences	21
		21
		21
		21
		21
		21
		$\frac{-}{21}$
		21
		21
		21
		21
		21
		$\frac{21}{21}$
		21
		$\frac{21}{21}$
	25.141 ne Onver Transfer and the Conner Conjecture	4 1
24	Borel-Moore Homology	21
		$\frac{1}{21}$
		$\frac{-}{21}$
		21
		21
		$\frac{21}{21}$
		21
	7 17 0 1	21
	<u>.</u>	21
	4 TO THE DWIE DPECHAL DEQUEED TO THE TOTAL TO THE TOTAL	∠ ⊥

	24.9 Poincaré Duality	21
	24.10The Cap Product	21
	24.11Intersection Theory	21
	24.12Uniqueness Theorems	21
	24.13Uniqueness Theorems for Maps and Relative Homology	21
	24.14The Künneth Formula	21
	24.15Change of Rings	21
	24.16Generalised Manifolds	21
	24.17Locally Homogenous Spaces	21
	24.18Homological Fibrations and <i>p</i> -adic Transformation Groups	21
	24.19The Transfer Homomorphism on Homology	21
	24.20Smith Theory in Homology	21
25	Cosheaves and Cech Homology	21
	25.1 Theory of Cosheaves	21
	25.2 Local Triviality	21
	25.3 Local Isomorphisms	21
	25.4 Chech Homology	21
	25.5 The Reflector	21
	25.6 Spectral Sequences	21
	25.7 Coresolutions	21
	25.8 Relative Cech Homology	21
	25.9 Locally Paracompact Spaces	21
	25.10Borel-Moore Homology	21
	25.11Modified Barel-Moore Homology	21
	25.12Singular Homology	21
	25.13Acyclic Coverings	21
	25.14Applications to Maps	21
V	I Introduction to Algebraic K-Theory	22
26	Projective Modules and $K_0\Lambda$	22
27	Constructing Projective Modules	22
28	The Whitehead Group $K_1\Lambda$	22
2 9	The Exact Sequence Associated with an Ideal	22
30	Steinberg Groups and the Functor K_2	22
31	Extending the Extact Sequences	22
32	The Case of a Commutative Banch Algebra	22
	The Product $K_1\Lambda \otimes K_1\Lambda \to K_2\Lambda$	22
	Computations in the Steinberg Group	22
	Computation of K_2Z	22
26		
	Matsumoto's Computation of K_2 of a Field Proof of Matsumoto's Theorem	$\frac{22}{22}$

38 More about Dedekind Domains	22
39 The Transfer Homomorphism	22
40 Power Norm Residue Theorems	22
41 Number Fields	22
VIII Formal Knot Theory, Kauffman	23
42 Introduction	23
43 States, Trails and the Clock Theorem	23
44 State Polynomials and the Duality Conjecture	23
45 Knots and Links	23
46 Axiomatic Link Calculations	23
47 Curliness and the Alexander Polynomial	23
48 The Coat of Many Colours	23
49 Spanning Surfaces	23
50 The Genus of Alternative Links	23
51 Ribbon Knot and the Arf Invariant	23
IV An Introduction to Invariants and Maduli Mulai	24
IX An Introduction to Invariants and Moduli, Mukai	
52 Invariants and Moduli	24
53 Rings and Polynomials	24
54 Algebraic Varieties	24
55 Algebraic Groups and Rings of Invariants	24
56 The Construction of Quotient Varieties	24
57 The Projective Quotients	24
58 The Numerical Criterion and Some Applications	24
59 Grassmannians and Vector Bundles	24
60 Curves and their Jacobians	24
61 Stable Vector Bundles on Curves	24
62 Moduli Functors	24
63 Intersection Numbers and the Verlinde Formula	24

X	Simplicial and Dendroidal Homotopy Theory	25
64	Operads	25
	64.1 Operads	25
	64.2 Algebras for Operads	26
	64.3 Trees	27
	64.4 Alternative Definitions for Operads	27
	64.5 Free Operads	27
	64.6 The Tensor Product of Operads	27
	64.7 The Boardman-Vogt Resolution of an Operad	27
	64.8 Configuration Spaces and the Fulton-MacPherson Operad	27
	64.9 Configuration Spaces and the Operad of Little Cubes	27
	one comparation spaces and the operat of Entire Cases	
65	Simplicial Sets	27
	65.1 The Simplex Category Δ	27
	65.1.1 Limits and Colimits of The Simplicial Category	28
	ob.1.1 Elimits and Commits of The Simplicial Category	20
66	Dendroidal Sets	28
67	Tensor Products of Dendroidal Sets	28
68	Kan Conditions for Simplicial Sets	28
69	Kan Conditions for Dendroidal Sets	28
70	Model Categories	28
7 1	Model Structures on the Category of Simplicial Sets	28
72	Three Model Structures on the Category of Dendroidal Sets	28
73	Reedy Categories and Diagrams of Spaces	28
74	Mapping Spaces and Bousfield Localisations	28
75	Dendroidal Spaces and ∞ -Operads	28
7 6	Left Fibrations and the Covariant Model Structure	28
77	Simplical Operads and ∞-Operads	28

Part I

Algebraic Geometry, Hartshorne

In this part, I will be going through the work of Hartshorne as he laid out in [Har13]. The book is sectioned into 5 distinct chapters - 'Varieties', 'Schemes', 'Cohomology', 'Curves' and 'Surfaces'. We study them in order as they flow nicely.

1 Varieties

Definition 1 Affine n-Space over k: Let k be a field and $n \in \mathbb{N}$. Then the affine n-space over k is defined

$$\mathbb{A}_{k}^{n} = \{(k_{1}, ..., k_{n}) \in k^{n}\}$$

It seems that this is a silly definition, but later on we will see that it is useful to have a distinction between the variety \mathbb{A}^n_k and the set of points in k^n . We later view \mathbb{A}^n_k as an affine variety - an object in some arbitrary space rather than the set of k-tuples.

Let $A = k[x_1, ..., x_n]$ be the polynomial ring over k in n variables. Then $f \in A$ is a map $f : k^n \to k$. We define the vanishing locus of this function in the following way:

Definition 2 Vanishing Locus of a Polynomial: Let $f \in A = k[x_1, ..., x_n]$, then the vanishing locus is


$$\mathbb{V}(f) := \{ p \in \mathbb{A}^n_k : f(p) = 0 \}$$

We can develop a more advanced analogue of this:

Definition 3 Vanishing Locus of a Set of Polynomials: Let $T = \{f_i\}_{i \in I} \subset A = k[x_1, ..., x_n]$, then the vanishing locus of T is

$$\mathbb{V}(T) := \{ p \in \mathbb{A}^n_k : f(p) = 0, \quad \forall f \in T \}$$

Some examples:






Figure 1: The vanishing locus of two separate polynomials plotted in \mathbb{R}^2

- 1.1 Affine Varieties
- 1.2 Projective Varieties
- 1.3 Morphisms
- 1.4 Rational Maps
- 1.5 Nonsingular Varieties
- 1.6 Nonsingular Curves
- 1.7 Intersections in Projective Space
- 2 Schemes
- 2.1 Sheaves
- 2.2 Schemes
- 2.3 First Properties of Schemes
- 2.4 Separated and Proper Morphisms
- 2.5 Sheaves of Modules
- 2.6 Divisors
- 2.7 Projective Morphisms
- 2.8 Differentials
- 2.9 Formal Schemes
- 3 Cohomology
- 3.1 Derived Functors
- 3.2 Cohomology of Sheaves
- 3.3 Cohomology of a Noetherian Affine Scheme
- 3.4 Cech Cohomology
- 3.5 The Cohomology of Projective Space
- 3.6 Ext Groups and Sheaves
- 3.7 The Serre Duality Theorem
- 3.8 Higher Direct Images of Sheaves
- 3.9 Flat Morphisms
- 3.10 Smooth Morphisms
- 3.11 The Theorem on Formal Functions
- 3.12 The Semicontinuity Theorem
- 4 Curves
- 4.1 Riemann-Roch Theorem
- 4.2 Hurwitz's Theorem
- 4.3 Embeddings in Projective Space
- 4.4 Elliptic Curves
- 4.5 The Canonical Embedding

- 5.5 Birational Transformations
- 5.6 Classification of Surfaces

Part II

Commutative Geometry with a View Toward Algebraic Geometry, Eisenbud

6 Basic Constructions

- 6.1 Roots of Commutative Algebra
- 6.1.1 Number Theory
- 6.1.2 Algebraic Curves and Function Theory
- 6.1.3 Invariant Theory
- 6.1.4 The Basis Theorem
- 6.1.5 Graded Rings
- 6.1.6 Algebra and Rings The Nullstellensatz
- 6.1.7 Geometric Invariant Theory
- 6.1.8 Projective Varieties
- 6.1.9 Hilbert Functions and Polynomials
- 6.1.10 Free Resolutions and the Syzygy THeorem

6.2 Localisation

A local ring is a ring with a single unique maximal ideal. The technique of localisation reduces many problems in commutative algebra to problems on commutative rings. The idea of localisation is as follows. Given a point p in an algebraic set $X \subset \mathbb{A}^n_k$, we want to investigate what X looks like near p, that is we want to investigate arbitrarily small open neighbourhoods of p in the Zariski topology. The Zariski open neighbourhoods of p are sets of the form $X \setminus Y$ for $p \not ! nY \subset X$.

6.2.1 Fractions

Definition 4 Localisation of an R-Module via a Multiplicatively Closed Subset U: Let R be a ring, M an R-module and $U \subset R$ a multiplicatively closed subset. The localisation of M at U, $M[U^{-1}]$ is the set of equivalent classes of pairs $(m, u) \sim (m', u')$ where $m, m' \in M$ and $u, u' \in U$ are related if there is some $v \in U$ such that v(mu' - m'u) = 0.

Proposition 1 Let U be multiplicatively closed set of R and let M be an R-module. An element $m \in M$ goes to 0 in $M[U^{-1}]$ under the map $\pi : M \to M[U^{-1}], m \to m/1$ if m is annihilated by an element $u \in U$. In particular, if M is finitely generated then $M[U^{-1}] = 0$ iff M is annihilated by an element of U.

proof

- 6.2.2 Hom and Tensor
- 6.2.3 The Construction of Primes
- 6.2.4 Rings and Modules of Finite Length
- 6.2.5 Products of Dom
- 6.3 Associated Primes and Primary Decomposition
- 6.3.1 Associated Primes
- 6.3.2 Prime Avoidance
- 6.3.3 Primary Decomposition
- 6.3.4 Primary Decomposition and Factorality
- 6.3.5 Primary Decomposition in the Graded Case
- 6.3.6 Extracting Information form Primary Decomposition
- 6.3.7 Why Primary Decomposition is not Unique
- 6.3.8 Geometric Interpretation of Primary Decomposition
- 6.3.9 Symbolic Powers and Functions Vanishing to High Order
- 6.4 Integral Dependence and the Nullstellensatz
- 6.4.1 The Cayley-Hamilton Theorem and Nakayama's Lemma
- 6.4.2 Normal Domains and the Normalisation Process
- 6.4.3 Normalisation in the Analytic Case
- 6.4.4 Primes in an Integral Extension
- 6.4.5 The Nullstellensatz
- 6.5 Filtrations and the Artin-Rees Lemma
- 6.5.1 Associated Graded Rings and Modules
- 6.5.2 The Blowup Algebra
- 6.5.3 The Krull Intersection Theorem
- 6.5.4 The Tangent Cone
- 6.6 Flat Families
- 6.6.1 Elementary Examples
- 6.6.2 Introduction to Tor
- 6.6.3 Criteria for Flatness
- 6.6.4 The Local Criterion for Flatness
- 6.6.5 The Rees Algebra
- 6.7 Completions and Hensel's Lemma
- 6.7.1 Examples and Definitions
- 6.7.2 The Utility of Completions
- 6.7.3 Lifting Idempotents
- 6.7.4 Cohen Structure Theory and Coefficient Fields

15

- 6.7.5 Basic Properties of Completion
- 6.7.6 Maps from Power Series Rings

Part III

Commutative Ring Theory, Matsumura		
9 Commutative Rings and Modules		
9.1 Ideals		
9.2 Modules		
9.3 Chain Conditions		
10 Prime Ideals		
10.1 Localisation and Spec of a Ring		
10.2 The Hilbert Nullstellensatz and First Steps in Dimension Theor		
10.3 Associated Primes and Primary Decomposition		
11 Properties of Extension Rings		
11.1 Flatness		
11.2 Completion and the Artin-Rees Lemma		
11.3 Integral Extensions		
12 Valuation Rings		
12.1 General Valuations		
12.2 DVRs amd Dedekind Rings		
12.3 Krull Rings		
13 Dimension Theory		
13.1 Graded Rings, the Hilbert Function and the Samuel Function		
13.2 Systems of Parameters and Multiplicity		
13.3 The Dimension of Extension Rings		
14 Regular Sequences		
14.1 Regular Sequences and the Koszul Complex		
14.2 Cohen-Macualay Rings		
14.3 Gorenstein Rings		
15 Regular Rings		
15.1 Regular Rings		
15.2 UFDs		
15.3 Complete Intersection Rings		

- The Local Flatness Criterion 16.1
- Flatness and Fibres 16.2 Generic Freeness and Open Loci Results 16.3

Flatness Revisited

16

Part IV
Algebraic Geometry I: Schemes,
Gortz-Wedhorn

Part V
Algebraic Geometry II: Cohomology of Schemes, Gortz-Wedhorn

Part VI

Sheaf Theory, Bredon

Singular Cohomology

de Rham Cohmomology

Alexander-Spanier Cohomology 21

22.1

22.2

22.3

20 Sheaves and Presheaves
20.1 Definitions
20.2 Homomorphisms, Subsheaves and Quotient Sheaves
20.3 Direct and Inverse Images
20.4 Cohomomorphisms
20.5 Algebraic Constructions
20.6 Supports
20.7 Classical Cohomology Theories
21 Sheaf Cohmology
21.1 Differential Sheaves and Resolutions
21.2 The Canonical Resolution and Sheaf Cohomology
21.3 Injective Sheaves
21.4 Acyclic Sheaves
21.5 Flabby Sheaves
21.6 Connected Sequences of Functors
21.7 Axioms for Cohmomology and the Cup Product
21.8 Maps of Spaces
21.9 Φ -Soft and Φ -Fine Sheaves
21.10 Subspaces
21.11 The Vietoris Maping Theorem and Homotopy Invariance
21.12 Relative Cohomology
21.13 Mayer-Vietoris Theorems
21.14 Continuity
21.15 The Künneth and Universal Coefficient Theorems
21.16 Dimension
21.17 Local Connectivity
21.18 Change pf Supports and Local Cohomology Groups
21.19 The Transfer Homomorphism and the Smith Sequences
21.20 Steenrod's Cyclic Reduced Powers
21.21 The Steenrod Operations
22 Comparison with Other Cohomology Theories

Part VII

Introduction to Algebraic K-Theory

- **26** Projective Modules and $K_0\Lambda$
- 27 Constructing Projective Modules
- 28 The Whitehead Group $K_1\Lambda$
- 29 The Exact Sequence Associated with an Ideal
- 30 Steinberg Groups and the Functor K_2
- 31 Extending the Extact Sequences
- 32 The Case of a Commutative Banch Algebra
- **33** The Product $K_1\Lambda \otimes K_1\Lambda \to K_2\Lambda$
- 34 Computations in the Steinberg Group
- 35 Computation of K_2Z
- 36 Matsumoto's Computation of K_2 of a Field
- 37 Proof of Matsumoto's Theorem
- 38 More about Dedekind Domains
- 39 The Transfer Homomorphism
- 40 Power Norm Residue Theorems
- 41 Number Fields

Part VIII

Formal Knot Theory, Kauffman

- 42 Introduction
- 43 States, Trails and the Clock Theorem
- 44 State Polynomials and the Duality Conjecture
- 45 Knots and Links
- 46 Axiomatic Link Calculations
- 47 Curliness and the Alexander Polynomial
- 48 The Coat of Many Colours
- 49 Spanning Surfaces
- 50 The Genus of Alternative Links
- 51 Ribbon Knot and the Arf Invariant

Part IX

An Introduction to Invariants and Moduli, Mukai

52	Invariants and Moduli
53	Rings and Polynomials
54	Algebraic Varieties
55	Algebraic Groups and Rings of Invariants
56	The Construction of Quotient Varieties
57	The Projective Quotients
58	The Numerical Criterion and Some Applications
59	Grassmannians and Vector Bundles
60	Curves and their Jacobians
61	Stable Vector Bundles on Curves
62	Moduli Functors
63	Intersection Numbers and the Verlinde Formula

Part X

Simplicial and Dendroidal Homotopy Theory

64 Operads

64.1 Operads

Definition 5 Operad: An operad P consists of a set of colours C and for each $n \geq 0$ and sequence $c_1, ..., c_n, c$ of colours in C, a set $P(c_1, ..., c_n; c)$ of operations, thought of as taking n inputs of colours $c_1, ..., c_n$ and with output of colour c. Moreover there are the structure maps

- $\forall c \in C, \exists 1_c \in P(c; c),$
- For $\sigma \in \Sigma_n$ a map

$$\sigma^* : P(c_1, ..., c_n; c) \to P(c_{\sigma(1)}, ..., c_{\sigma(n)}; c)$$

denoted $\sigma^* \circ p = p \circ \sigma$

- For any sequence $c_1, ..., c_n$ and n-tuple of sequences $d_1^i, ..., d_{k_i}^i$, a composition
- •

$$\gamma: P(c_1, ..., c_n; c) \times \prod_{i=1}^n P(d_1^i, ..., d_{k_i}^i; c_i) \to P(d_1^1, ..., d_{k_n}^n; c)$$

which is written as $\gamma(p, q_1, ..., 1_n) \to p \circ (q_1, ..., q_n)$

and there are further requirements on the structure maps:

- $\forall p \in P(c_1, ..., c_n; c), \gamma(1_c, p) = p,$
- $\forall p \in P(c_1, ..., c_n; c), \gamma(p, 1_{c_1}, ..., 1_{c_n}) = p$

There are some classical notation we must be weary of and state here. Let P be an operad with a singleton colour set, i.e. $C = \{*\}$. Then we can write $P(c_1, ..., c_n; c) = P(c, ..., c; c) =: P(n)$. There is an obvious formulation for the compostion in this case:

$$P(n) \times \prod_{i=1}^{n} P(k_i) \to P(k_1 + \dots + k_n)$$

In this case we say P is uncoloured. If $C \notin \{\phi, \{*\}\}$ then we say P is a coloured operad. We can then clearly see that

$$\begin{tabular}{ll} Monoids & \longleftarrow & Categories \\ & & & & & & \\ & & & & & \\ Uncoloured & Operads & \longleftarrow & Operads \\ \end{tabular}$$

The definition of an operad allows for n = 0 in $P(c_1, ..., c_n; c)$, which we define as P(-; c). The elements of P(-; c) are called the constants of colour c, and an operad with $P(-; c) = \{*_c\}$ is called unital. An operad P is open if there are no constants for any colour, i.e. its interior P^o (the set of constants) is empty.

The most fundamental examples of operads are Com and Ass:

- Com is the commutative operad with $Com(n) = \{*\},\$
- Ass is the associative operad with $Ass(n) = \Sigma_n$
- Tree^{pl} is the planar tree operad. The n-ary operations here are the set of planar rooted trees with n numbered leaves.

$$\mathbf{Tree}^{pl}(7)\ni \tilde{T}=\ \ \overset{T_2}{\searrow}\ \ \overset{T_1}{\searrow}$$

Then in fact $\gamma(T, T_1, T_2) = \tilde{T}$. The operation of composition on the operad **Tree**^{pl} is computed as $\gamma(T \in \mathbf{Tree}^{pl}(n), T_1 \in \mathbf{Tree}^{pl}(k_1), ..., T_n \in \mathbf{Tree}^{pl}(k_n)) = \hat{T}$ where $\hat{T} \in \mathbf{Tree}^{pl}(k_1 + ... + k_n)$ is the original T but with the subtree T_i grafted onto the leaf i for all choices of i.

Definition 6 Topological Operad: A topological operad is an operad P where each set of operations $P(c_1,...,c_n;c)$ is equipped with some topology and all the structure maps are continuous with respect to this topology.

The most basic form of a topological operad is the little d-cubs operad \mathbf{E}_d . The space $\mathbf{E}_d(n)$ is the space of n numbered d-dimensional cubes inside the d-dimensional unit cube $[0,1]^d$. The operadic composition between $p \in \mathbf{E}_d(n)$ with operations $q_1, ..., q_n$ is given by substituting the rescaled q_i into the ith cube of p. Note that this is really just a topological analogue to the planar tree operad \mathbf{Tree}^{pl} , where instead of grafting trees onto leaves we are scaling and embedding cubes in some smooth way.

More specifically, a point in $\mathbf{E}_d(n)$ is an *n*-tuple of embeddings $f_1, ..., f_n : [0, 1]^d \to [0, 1]^d$ satisfying:

- Each f_i is the composition of d affine embeddings,
- The interiors of the cubes embedded by f_i are mutually disjoint

We can now observe that operads form a category in a very natural fashion. Given two operads P,Q, a morphism $\varphi:P\to Q$ is a function $f:C_p\to C_Q$ on operadic colours and for each sequence $c_1,...,c_n;c$ of C_P , we have

$$\varphi_{(c_1,...,c_n;c)}: P(c_1,...,c_n;c) \to Q(f(c_1),...,f(c_n);f(c))$$

that is compatible in the natural way with Σ_n actions

64.2 Algebras for Operads

Definition 7 Operadic Algebras: Let P be an operad. A P-algebra A is a family of sets $\{A_c\}_{c \in C_P}$ together with maps

$$P(c_1,...,c_n;c)\times A_{c_1}\times...\times A_{c_n}\to A_c$$

written $(p, a_1, ..., a_n) \rightarrow A(p)(a_1, ..., a_n)$. These maps also satisfy:

- $1_c(a) = a \quad \forall a \in A_c$
- $\sigma \in \Sigma_n, a_i \in A_{c_i}, \sigma^* p(a_{\sigma(1)}, ..., a_{\sigma(n)}) = p(a_1, ..., a_n)$

Definition 8 Morphisms of Operadic Algebras: Let A, B be two P-algebras. A morphism $f: A \rightarrow B$ is a family of maps

$$f_c: A_c \to B_c$$

which are compatible:

$$f_c(A(p)(a_1,...,a_n)) = B(p)(f_{c_1}(a_1),...,f_{c_n}(a_n))$$

Definition 9 Category of P-Algebras: Let P be an operad. Then we have a category of P-algebras Alg_P with:

- $ObAlg_P = \{P algebras \ A\}$
- $Hom_{Alg_P}(A, B) = \{f : A \to B : f \text{ is a morphism of algebras}\}$

We can now see some examples of operadic algebras. A **Com** algebra is a set A together with a map $\mu_n: A^{\times n} \to A$ for each $n \geq 0$. We can then verify that the category of algebras over the commutative operad, $\text{Alg}_{\textbf{Com}}$ is the category of commutative monoids. In a similar way $\text{Alg}_{\textbf{Ass}}$ is the category of associative monoids.

Consider the little-d cubes operad \mathbf{E}_d . Let X be a topological space with basepoint x_0 . Then the loop space of X is ΩX , the space of basepoint preserving maps $S^1 \to X$, or in otherwords $\{\omega : [0,1] \to X, \omega(\partial[0,1]) = x_0\}$. One can then inductively construct the d-fold loop space $\Omega^d X = \Omega(\Omega^{d-1}X)$. $\Omega^d X$ is very naturally a \mathbf{E}_d algebra.

- 64.3 Trees
- 64.4 Alternative Definitions for Operads
- 64.5 Free Operads
- 64.6 The Tensor Product of Operads
- 64.7 The Boardman-Vogt Resolution of an Operad
- 64.8 Configuration Spaces and the Fulton-MacPherson Operad
- 64.9 Configuration Spaces and the Operad of Little Cubes
- 65 Simplicial Sets

65.1 The Simplex Category Δ

Definition 10 The Simplex Category Δ : Δ is the category with:

- $Ob\Delta = \mathbb{N}$.
- $Hom_{\Lambda}([n],[m]) = \{order\ preserving\ maps\ [n] \to [m]\}$

There are special maps in Δ - the elementary faces $\delta^i:[m-1]\to[m]$ and elementary degeneracies $\sigma^i:[m]\to[m-1]\ 0\le i\le m-1$:

$$\delta^{i}(j) = \begin{cases} j & j < i \\ j+1 & j \ge i \end{cases}, \quad \sigma^{i}(j) = \begin{cases} j & j \le i \\ j-1 & j > i \end{cases}$$

These have some nice relations, called the cosimplicial identities:

- $\sigma_i \sigma_j = \sigma_{j-1} \sigma_i$, i < j
- $\delta_i \delta_i = \delta_i \delta_{i-1}$

65.1.1 Limits and Colimits of The Simplicial Category

$$\begin{array}{ccc} k & \xrightarrow{f} & n \\ \downarrow^g & \downarrow \\ m & \to & m+n \end{array}$$

is a pushout, where f(i) = i, g(i) = m - k + i

- 66 Dendroidal Sets
- 67 Tensor Products of Dendroidal Sets
- 68 Kan Conditions for Simplicial Sets
- 69 Kan Conditions for Dendroidal Sets
- 70 Model Categories
- 71 Model Structures on the Category of Simplicial Sets
- 72 Three Model Structures on the Category of Dendroidal Sets
- 73 Reedy Categories and Diagrams of Spaces
- 74 Mapping Spaces and Bousfield Localisations
- 75 Dendroidal Spaces and ∞ -Operads
- 76 Left Fibrations and the Covariant Model Structure
- 77 Simplical Operads and ∞ -Operads

References

 $[{\rm Har}13]$ Robin Hartshorne. Algebraic geometry, volume 52. Springer Science & Business Media, 2013.