

a
Expressed L–R repertoire per sample
Expressed L–R interactions per sample (5 NVU types)
P = 0.48
P = 0.89
Ctrl
AD
Hippocampus
Cortex
b
Hippocampus: 0 of 1,565 pairs at FDR < 0.05
(cell text: nominal P < 0.01 / tested)
Sender
Receiver
Micro
Endo
Peri
Astro
Oligo
Median log2FC
c
Cortex: 0 of 2,011 pairs at FDR < 0.05
(cell text: nominal P < 0.01 / tested)
d
Legacy APP→SORL1 claim, per donor
min FDR = 0.81
min FDR = 1
APP→SORL1 score (into microglia)
e
Top 20 hippocampal L–R changes by nominal P (all FDR ≥ 0.68); OLS with sex and yield
Micro→Oligo DSCAM→DCC
Peri→Peri LAMC1→ITGA7 ITGB1
Endo→Astro TFPI→F3
Micro→Endo PDGFC→FLT1
Endo→Peri CALM1→GRM3
Peri→Endo CNTN6→PTPRG
Peri→Endo COL18A1→ITGA5
Peri→Oligo COL4A1→ITGAV ITGB8
Endo→Peri MAML2→NOTCH3
Endo→Peri LAMA3→ITGA7 ITGB1
Peri→Oligo COL4A1→CD47
Peri→Astro COL4A1→CD47
Peri→Oligo APP→DCC
Endo→Oligo APP→DCC
Endo→Micro ADAM10→CADM1
Peri→Endo FGF1→NRP1
Peri→Peri CNTN6→PTPRG
Peri→Oligo COL4A2→ITGAV ITGB8
Peri→Peri LAMA2→ITGA7 ITGB1
Peri→Oligo IL1RAPL1→PTPRD
Hippocampus (discovery)
Cortex (same donors)
log2 fold change of interaction score (AD/Control) ± 95% CI