
a
central memory CD8-positive, alpha-beta T cell (4)
megakaryocyte (3)
naive B cell (6)
erythrocyte (3)
effector memory CD8-positive, alpha-beta T cell, terminally differentiated (5)
CD16-negative, CD56-bright natural killer cell (6)
plasmablast (5)
regulatory T cell (6)
central memory CD4-positive, alpha-beta T cell (6)
CD16-positive, CD56-dim natural killer cell (6)
mast cell (7)
non-classical monocyte (7)
endothelial cell of lymphatic vessel (6)
mucosal invariant T cell (5)
conventional dendritic cell (8)
T follicular helper cell (5)
mesothelial cell (2)
effector memory CD4-positive, alpha-beta T cell (6)
germinal center B cell (2)
effector memory CD8-positive, alpha-beta T cell (7)
plasma cell (7)
endothelial cell of artery (4)
vein endothelial cell (4)
neutrophil (3)
fat cell (2)
classical monocyte (7)
goblet cell (2)
gamma-delta T cell (4)
pericyte (5)
myofibroblast cell (2)
0.0
0.2
Calibrated invariance
(cosine above the organ pair's null)
b
Erythroid/Mega (n=2)
Lymphoid (n=15)
Myeloid (n=5)
Epithelial (n=2)
Endothelial (n=3)
Mesenchymal/Stromal (n=3)
0.0
0.2
Calibrated invariance
circulating lineage
tissue-resident lineage
null ± 1 SD
specific term
coarse term, comparable organs
median (specific)