
A
Microenvironment subcluster size
homeostatic microglia
TGFBI/AOAH macrophage
hypoxic/glycolytic TAM
interferon TAM
ribosomal residue (cells)
proliferating myeloid
THEMIS/CAMK4 T
CD8 T
CD4 T
NK
IL7R/KLRB1 T
endothelial (neural genes)
tip/angiogenic EC
capillary EC
capillary EC 2
pericyte
mature oligodendrocyte 1
mature oligodendrocyte 2
mature oligodendrocyte 3
mature oligodendrocyte 4
committed OPC
10^2
10^3
10^4
cells (log scale)
0
0.5
1
fraction nuclei
myeloid
vascular
T/NK
oligodendrocyte
B
Assay purity per compartment
myeloid
vascular
T/NK
oligodendrocyte
assay purity
(cells in < 10 % or > 90 % nuclei clusters)
1.0
0.8
0.6
0.4
0.2
0.0
atlas scVI
per-compartment scVI, masked
C
Composition effects per contrast
T/NK
astrocyte
malignant
malignant: AC-like
malignant: MES-like
malignant: OC-like
malignant: OPC-like
malignant: cycling
malignant: unassigned
myeloid
neuron
oligodendrocyte
vascular
nuclei vs cells
H3.1 vs H3.3
adult vs paediatric
thalamus vs pons
ONC201 vs standard
Δ logit proportion
3
2
1
0
−1
−2
−3