

A
Top genes per metaprogram
MP1 cycling
ASPM, TOP2A, CENPF, MKI67, KIF14, TPX2, CENPE, NUF2
MP2 ribosomal*
RPL32, RPS18, RPS8, RPS27, EEF1A1, RPL8, RPLP1, RPL41
MP3 OPC-like
FGF12, DSCAM, TNR, CA10, DCC, CADM2, MMP16, NXPH1
MP4 hypoxia
SLC6A6, NDRG1, AKAP12, VEGFA, HILPDA, IGFBP5, SLC2A1, PLOD2
MP5 OC-like
TNS3, BCAS1, OPCML, GPR17, PKP4, FGF12, BMPER, FERMT1
MP6 OPC-like
LHFPL3, SEZ6L, LRRC4C, KCNIP4, MMP16, SEMA5A, CA10, PCDH15
MP7 AC-like
DAAM2, NKAIN3, MGAT4C, EEPD1, DCLK1, SLC4A4, KCNN3, ROBO2
MP8 MES-like
CD44, NAMPT, HRH1, SAMD4A, TGFB2, TPST1, COLEC12, ACTN1
MP9 cilia (MP9)
ADGB, CFAP43, DNAH9, ZBBX, MAP3K19, CFAP299, SPAG17, CFAP73
MP10 AC-like
SLC4A4, EEPD1, FAT3, PTPRT, ADGRV1, SLC1A3, MMD2, SLC6A11
MP11 cycling
BRIP1, DTL, CENPP, CENPK, RFC3, MMS22L, DIAPH3, FANCI
MP12 mitochondrial*
MT-CO2, MT-ND4, MT-CO3, MT-ND3, MT-CYB, MT-ATP6, MT-ND2, MT-ND1
MP13 IEG stress*
IER2, FOS, JUNB, EGR1, JUN, FOSB, BTG2, RHOB
MP14 AC-like
DCLK1, EEPD1, SORBS1, AQP4, DAAM2, RGMA, RNF19A, CCDC85A
MP15 AC-like
EEPD1, CD44, SPARCL1, SORBS1, SLC4A4, TNC, LRIG1, GLIS3
B
Overlap with published gene sets
MP1
MP2
MP3
MP4
MP5
MP6
MP7
MP8
MP9
MP10
MP11
MP12
MP13
MP14
MP15
20.0
17.5
15.0
12.5
10.0
7.5
5.0
2.5
0.0
−log10 q (overlap)
curated AC like markers
curated Cycling markers
curated Glycolysis
curated Hypoxia
curated Dissociation
curated MES like markers
curated OC like markers
curated OPC like markers
curated Stress HSP
Filbin untrac. AC like
Filbin untrac. MES like
Filbin untrac. OPC like
Filbin untrac. cycling
Liu DE AC-like
Liu DE MES-like
Liu DE G2M
Liu DE OC-like
Liu DE OPC-like-1
Liu DE OPC-like-2
Liu DE OPC-like-3
Liu S2a AC-like
Liu S2a G2M
Liu S2a MES-like
Liu S2a OC-like
Liu S2a OPC-like-1
Liu S2a OPC-like-2
Liu S2a OPC-like-3
Liu S2a S
C
Metaprograms across dendrogram cuts
metaprograms
replicated
count
programmes assigned
dendrogram cut
0.750
0.775
0.800
0.825
0.850
0.875
0.900
0.925
D
Recovery of Jessa 2022 programmes
best usage ρ with an MP
OPC-like
AC-like
OC-like
MES-like
cycling
hypoxia
cilia (MP9)