
A
Marker genes, gene by gene
AC-like (29)
MES-like (27)
OC-like (27)
OPC-like (28)
cycling (27)
detected in mouse tumour cells
detected in human cells of that subpopulation
0.0
0.2
0.4
0.6
0.8
1.0
B
Pathway activity, human and mouse
Androgen
EGFR
Estrogen
Hypoxia
JAK-STAT
MAPK
NFkB
PI3K
TGFb
TNFa
Trail
VEGF
WNT
p53
PROGENy activity
20
10
0
−10
−20
H AC-like
H MES-like
H OC-like
H OPC-like
H cycling
H unassigned
M T/NK (normal)
M astrocyte-like (normal)
M cerebellar neuron
M myeloid (normal)
M other
M tumour
C
Transcription-factor activity
AR
E2F4
ETS1
FOS
HNF1B
HSF1
IRF1
IRF5
KLF2
MYC
NFKB1
REL
RELA
SOX4
SPI1
STAT4
TP53
ZEB1
CollecTRI activity
10
5
0
−5
−10
H AC-like
H MES-like
H OC-like
H OPC-like
H cycling
H unassigned
M T/NK (normal)
M astrocyte-like (normal)
M cerebellar neuron
M myeloid (normal)
M other
M tumour
D
Pseudobulk correlation per subpopulation
T/NK (normal)
astrocyte-like (normal)
cerebellar neuron
myeloid (normal)
other
tumour
-0.12 0.03 -0.25 -0.07 0.33
0.27 0.03 0.15 -0.05 -0.31
0.08 -0.10 0.23 0.15 -0.29
0.06 0.30 -0.18 -0.29 0.06
-0.04 0.11 0.19 -0.08 -0.18
-0.36 -0.40 -0.13 0.35 0.47
AC-like
MES-like
OC-like
OPC-like
cycling
Spearman rho
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6