
A
Marker genes, gene by gene
AC-like (29)
MES-like (27)
OC-like (27)
OPC-like (28)
cycling (27)
detected in mouse tumour cells
detected in human cells of that subpopulation
B
Pathway activity, human and mouse
Human
Mouse
Androgen
EGFR
Estrogen
Hypoxia
JAK-STAT
MAPK
NFkB
PI3K
TGFb
TNFa
Trail
VEGF
WNT
p53
AC-like
MES-like
OC-like
OPC-like
cycling
unassigned
T/NK (normal)
astrocyte-like (normal)
cerebellar neuron
myeloid (normal)
other
tumour
PROGENy activity
C
Transcription-factor activity
Human
Mouse
AR
E2F4
ETS1
FOS
HNF1B
HSF1
IRF1
IRF5
KLF2
MYC
NFKB1
REL
RELA
SOX4
SPI1
STAT4
TP53
ZEB1
CollecTRI activity
D
Pseudobulk correlation per subpopulation
T/NK (normal) -0.12 0.03 -0.25 -0.07 0.33
astrocyte-like (normal) 0.27 0.03 0.15 -0.05 -0.31
cerebellar neuron 0.08 -0.10 0.23 0.15 -0.29
myeloid (normal) 0.06 0.30 -0.18 -0.29 0.06
other -0.04 0.11 0.19 -0.08 -0.18
tumour -0.36 -0.40 -0.13 0.35 0.47
Spearman rho