

Lab#4 Extension

-1

Lab#4 is extended for another week for

- i) Midterm exams
- ii) Synchronizing the labs and lectures

Lab#4 becomes a 3-week lab and its due date is March 13, 2019.

All other remaining labs and due dates are pushed back for one week accordingly.


Make-up Class

0

For university closure on Feb. 12, 2019

Within {Feb. 26, 27, Mar. 6, 7, 8}, the only availability is:

Date	Thursday, March 7, 2019
Time	9:00-12:00
Location	VIC608



Today

2



Data Modeling
in
RDBMS

Real World Entity

Conceptual Level | Entity-Relationship Model (E/R)

| Logical Level | Relational Model

| Physical Level | SQL

Computable Entity

Welcome | Relational Algebra | Unary Op | Binary Op

Relational

3


Edgar Frank “Ted” Codd, IBM, 1969, 1970

Information Retrieval

A Relational Model of Data for Large Shared Data Banks

E. F. Codd

IBM Research Laboratory, San Jose, California



Relational Model \times Algebra

4

Given a set, defining operations on elements of the set!

Given $Z = \{\text{integers}\} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Operators & Operands:

Unary: $-(2)$, $2!$

Binary: $2+3$, $2*3$, 2^3

Closure:

Result is also an element of the set



Discrete
Mathematics

Relational Model × Algebra

5

Given relational (table) schema filled with actual data instances (rows):
Operations to **SELECT** Information **FROM** Relations

Query in Natural Language → Query in Math Formula

Relational Model \times Algebra

6

Who made 'Pulp Fiction'?

π

Director.FirstName
Director.LastName

$(\sigma$

Movie.Title='PulpFiction'
Movie.Id=MovieDirector.MovieId
Director.Id=MovieDirector.DirectorId

$(\text{Movie} \times \text{MovieDirector} \times \text{Director}))$

AND

AND

Relational Model \times Algebra

7

Who acted in 'Pulp Fiction'?

π

Actor.FirstName
Actor.LastName

$(\sigma$

(Movie \times StarIn \times Director))

Movie.Title='PulpFiction' AND
Movie.Id=StarIn.MoviId AND
Actor.Id=StarIn.ActorId

Relational Model \times Algebra

8

Given relational (table) schema filled with actual data instances (rows):
Operations to **SELECT** Information **FROM** Relations

Unary Operation

$\pi(R)$: Project

$\sigma(R)$: Select

$\rho(R)$: Rename

Binary Operation (Set Theory)

$R1 \cup R2$: Union

$R1 \cap R2$: Intersection

$R1 \setminus R2$: Set Difference

$R1 \times R2$: Cartesian Product

Relational Model \times Algebra

9

Given relational (table) schema filled with actual data instances (rows):

Operations to SELECT Information FROM Relations

Operations to write query

Unary Operation

$\pi(R)$: Project

$\sigma(R)$: Select

$\rho(R)$: Rename

Binary Operation (Set Theory)

$R1 \cup R2$: Union

$R1 \cap R2$: Intersection

$R1 \setminus R2$: Set Difference

$R1 \times R2$: Cartesian Product

Algebra × Project (π)

10

π , pi, is used to select a subset of attributes (columns) from a relation

$$A = \pi_{\langle \text{attribute list} \rangle}(R)$$



Vertical
Filtering

R is a relation

$\langle \text{attribute list} \rangle$ subset of attributes of R

A is a relation including all tuples in R with only attributes in list

Algebra × Project (π)

11

<u>Id</u>	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

What are directors' name?

$$A = \pi_{\text{FirstName, LastName}}(\text{Director})$$

Algebra \times Project (π)

12

<u>Id</u>	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

What are directors' name?

$$A = \pi_{\text{FirstName, LastName}}(\text{Director})$$

Algebra \times Project (π)

13

<u>Id</u>	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

How many movies each director made?

$$A = \pi_{\text{FirstName, LastName, MovieCount}}(\text{Director})$$

Algebra \times Project (π)

14

<u>Id</u>	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

How many movies each director made?

$$A = \pi_{\text{FirstName, LastName, MovieCount}}(\text{Director})$$

Relational Model \times Algebra

15

Given relational (table) schema filled with actual data instances (rows):

Operations to SELECT Information FROM Relations

Operations to write query

Unary Operation

$\pi(R)$: Project

$\sigma(R)$: Select

$\rho(R)$: Rename

Binary Operation (Set Theory)

$R1 \cup R2$: Union

$R1 \cap R2$: Intersection

$R1 \setminus R2$: Set Difference

$R1 \times R2$: Cartesian Product

Algebra × Selection (σ)

16

σ , sigma, is used to select a subset of tuples from a relation based on a condition (θ) over relation's attributes.

$$A = \sigma_{\theta}(R)$$



Horizontal
Filtering

R is a relation

θ is a Boolean expression on the attributes of R

A is a relation including tuples that make θ true

Algebra × Selection (σ)

17

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which director was born in US?

$A = \sigma_{\text{PlaceOfBirth}='USA'}(\text{Director})$

Algebra × Selection (σ)

18

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which director was born in US?

$A = \sigma_{\text{PlaceOfBirth}='USA'}(\text{Director})$

Algebra × Selection (σ)

19

<u>Id</u>	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which director make more than 20 movies?

$$A = \sigma_{\text{MovieCount} > 20} (\text{Director})$$

Algebra × Selection (σ)

20

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which director make more than 20 movies?

$$A = \sigma_{\text{MovieCount} > 20} (\text{Director})$$

Algebra × Selection (σ)

21

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which director has same first and last names?

$$A = \sigma_{\text{FirstName} = \text{LastName}} (\text{Director})$$

Algebra × Selection (σ)

22

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which director has same first and last names?


$$\begin{aligned} A &= \sigma_{\text{FirstName} = \text{LastName}} (\text{Director}) \\ &= \emptyset \end{aligned}$$

Algebra × Selection (σ)

23

θ can be made up of number of Boolean clauses

$$A = \sigma_{\theta}(R)$$



Algebra × Selection (σ)

24

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies?

$A = \sigma_{\text{PlaceOfBirth}='USA' \text{ AND } \text{MovieCount} > 20} (\text{Director})$

Algebra × Selection (σ)

25

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies or is not American?

A1 = $\sigma_{\text{PlaceOfBirth}='USA' \text{ AND } \text{MovieCount} > 20 \text{ OR } \text{PlaceOfBirth} \neq 'USA'}$ (Director)

A2 = $\sigma_{\text{PlaceOfBirth}='USA' \text{ AND } (\text{MovieCount} > 20 \text{ OR } \text{PlaceOfBirth} \neq 'USA')}$ (Director)

A3 = $\sigma_{(\text{PlaceOfBirth}='USA' \text{ AND } \text{MovieCount} > 20) \text{ OR } (\text{PlaceOfBirth} \neq 'USA')}$ (Director)

Algebra × Selection (σ)

26

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies or is not American?

$A2 = \sigma_{\text{PlaceOfBirth}='USA' \text{ AND } (\text{MovieCount} > 20 \text{ OR } \text{PlaceOfBirth} \neq 'USA')}$ (Director)



Algebra × Selection (σ)

27

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35


Which American director made more than 20 movies or is not American?

$A3 = \sigma_{(\text{PlaceOfBirth}='USA' \text{ AND MovieCount} > 20) \text{ OR } (\text{PlaceOfBirth} \neq 'USA')}$ (Director)




Algebra \times Selection (σ)

28



Algebra \times Selection (σ)

29



Algebra \times Selection (σ)

30

Operation Precedence

$$() > \S > + > \neq$$

Commutative Law

$$A \S B = B \S A \text{ (we say } A \text{ commutes with } B \text{ under } \S)$$

Associative Law

$$A \S (B \S C) = (A \S B) \S C = A \S B \S C$$

Distributive Law

$$A \S (B + C) = (A \S B) + (A \S C)$$

Algebra \times Selection (σ)

31

Operation Precedence for Logical Operations

() > NOT > AND = OR

Commutative Law

$A \text{ AND } B = B \text{ AND } A$ (we say A commutes with B under AND)

$A \text{ OR } B = B \text{ OR } A$ (we say A commutes with B under OR)

Associative Law

$A \text{ AND } (B \text{ AND } C) = (A \text{ AND } B) \text{ AND } C = A \text{ AND } B \text{ AND } C$

$A \text{ OR } (B \text{ OR } C) = (A \text{ OR } B) \text{ OR } C = A \text{ OR } B \text{ OR } C$

Distributive Law

$A \text{ AND } (B \text{ OR } C) = (A \text{ AND } B) \text{ OR } (A \text{ AND } C)$

$A \text{ OR } (B \text{ AND } C) = (A \text{ OR } B) \text{ AND } (A \text{ OR } C)$

de Morgan's Theorem**

$\text{NOT } (A \text{ AND } B) = \text{NOT}(A) \text{ OR } \text{NOT}(B)$

$\text{NOT } (A \text{ OR } B) = \text{NOT}(A) \text{ AND } \text{NOT}(B)$

Algebra × Selection (σ)

32

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies?

$$\begin{aligned} A &= \sigma_{\text{PlaceOfBirth}='USA' \text{ AND } \text{MovieCount} > 20} (\text{Director}) \\ &= \sigma_{\text{MovieCount} > 20 \text{ AND } \text{PlaceOfBirth}='USA'} (\text{Director}) \end{aligned}$$

Algebra × Selection (σ)

33

<u>Id</u>	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies?

$$A1 = \sigma_{\text{PlaceOfBirth}='USA'}(\text{Director})$$

$$A = \sigma_{\text{MovieCount} > 20}(A1)$$

Algebra × Selection (σ)

34

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies?

$A = \sigma_{\text{MovieCount} > 20} (\sigma_{\text{PlaceOfBirth} = \text{'USA'}} (\text{Director}))$

Algebra × Selection (σ)

35

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Which American director made more than 20 movies?

$$\begin{aligned} A &= \sigma_{\text{PlaceOfBirth}='USA'} (\sigma_{\text{MovieCount} > 20} (\text{Director})) \\ &= \sigma_{\text{PlaceOfBirth}='USA' \text{ AND } \text{MovieCount} > 20} (\text{Director}) \\ &= \sigma_{\text{MovieCount} > 20 \text{ AND } \text{PlaceOfBirth}='USA'} (\text{Director}) \\ &= \sigma_{\text{MovieCount} > 20} (\sigma_{\text{PlaceOfBirth}='USA'} (\text{Director})) \end{aligned}$$

Algebra × Selection (σ)

36

$$\sigma_{\theta}(\sigma_{\theta'}(R))$$

=

$$\sigma_{\theta \text{ AND } \theta'}(R)$$

=

$$\sigma_{\theta' \text{ AND } \theta}(R)$$

=

$$\sigma_{\theta'}(\sigma_{\theta}(R))$$



Commutative
Law

Corollary: $\sigma_{\theta}(\sigma_{\theta'}(\sigma_{\theta''}(\sigma_{\theta'''}(R))) = \sigma_{\theta \text{ AND } \theta' \text{ AND } \theta'' \dots \text{ AND } \theta'''(R)}$

Algebra \times Selection (σ) \times Project (π)

37

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Find directors' name who make more than 20 movies?

Algebra \times Selection (σ) \times Project (π)

38

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Find directors' name who make more than 20 movies?

$$A1 = \sigma_{\text{MovieCount} > 20} (\text{Director})$$

Algebra \times Selection (σ) \times Project (π)

39

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Find directors' name who make more than 20 movies?

$$A1 = \sigma_{\text{MovieCount} > 20}(\text{Director})$$

$$A = \pi_{\text{FirstName, LastName}}(A1)$$

$$= \pi_{\text{FirstName, LastName}}(\sigma_{\text{MovieCount} > 20}(\text{Director}))$$

Algebra \times Selection (σ) \times Project (π) 40

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

Find directors' name who make more than 20 movies?

 $A = \sigma_{\text{MovieCount} > 20} (\pi_{\text{FirstName, LastName}} (\text{Director}))$

Relational Model \times Algebra

41

Given relational (table) schema filled with actual data instances (rows):

Operations to SELECT Information FROM Relations

Operations to write query

Unary Operation

$\pi(R)$: Project

$\sigma(R)$: Select

$\rho(R)$: Rename

Binary Operation (Set Theory)

$R1 \cup R2$: Union

$R1 \cap R2$: Intersection

$R1 \setminus R2$: Set Difference

$R1 \times R2$: Cartesian Product

Algebra \times Rename (ρ)

42

ρ , rho, is used to rename a relation or its attributes or both

$$A = \rho_{\langle R'(a'/a, b'/b, \dots) \rangle}(R)$$

R is a relation

R' is the new name for R(a, b, ...)

a' is the new name for attribute a of R

b' is the new name for attribute b of R

...

Algebra × Rename (ρ)

43

Id	FirstName	LastName	DateOfBirth	PlaceOfBirth	BestMovieId	MovieCount
1	Stanley	Kubrick	Jul. 26, 1928	USA	1	13
2	Alfred	Hitchcock	Aug. 13, 1899	England	203	47
3	Clint	Eastwood	May 31, 1930	USA	803	35

$$A = \pi_{\text{FirstName, Name}} (\rho_{\text{ActiveDirector(Name/LastName)}} (\sigma_{\text{MovieCount} > 20}(\text{Director})))$$

We'll c
its real
use 😊

