

Results

Linear Mixed Models

```
jaspMixedModels::MixedModelsLMM(  
  data = NULL,  
  version = "0.95",  
  formula = Log_rt ~ Block * congruency + (1 | caseid),  
  contrasts = NULL,  
  fixedEffectEstimate = TRUE,  
  marginalMeansTerms = ~ Block + congruency,  
  modelSummary = TRUE,  
  plotBackgroundData = ~ caseid.nominal,  
  plotBackgroundElement = "boxplot",  
  plotEstimatesTable = TRUE,  
  plotHorizontalAxis = ~ Block,  
  plotLevelsByColor = TRUE,  
  plotLevelsByLinetype = FALSE,  
  plotSeparateLines = ~ congruency,  
  plotTheme = "apa",  
  randomEffectEstimate = TRUE,  
  setSeed = TRUE,  
  testMethod = "likelihoodRatioTest",  
  trendsContrasts = NULL)
```

ANOVA Summary

Effect	df	ChiSq	p
Block	1	1263.43	1.003×10 ⁻²⁷⁶
congruency	1	21.13	4.302×10 ⁻⁶
Block * congruency	1	38.15	6.556×10 ⁻¹⁰

Note. 463 observations were removed due to missing values.

Note. Model terms tested with likelihood ratio tests testMethod.

Note. The following variable is used as a random effects grouping factor: 'caseid'.

Note. Type III Sum of Squares

Model summary

Fit statistics

Deviance	log Lik.	df	AIC	BIC
19201	−9600	6	19213	19261

Note. The model was fitted using maximum likelihood.

Sample sizes

Observations	Levels of RE grouping factors
	caseid
22649	36

Fixed Effects Estimates

Term	Estimate	SE	df	t	p
Intercept	6.572	0.024	36.03	269.259	3.939×10^{-61}
Block (1)	−0.090	0.003	22613.02	−36.047	1.019×10^{-276}
congruency (1)	−0.012	0.003	22613.28	−4.597	4.303×10^{-6}
Block (1) * congruency (1)	−0.015	0.003	22613.20	−6.179	6.559×10^{-10}

Note. The intercept corresponds to the (unweighted) grand mean; for each factor with k levels, k - 1 parameters are estimated with sum contrast coding. Consequently, the estimates cannot be directly mapped to factor levels. Use estimated marginal means for obtaining estimates for each factor level/design cell or their differences.

Random Effect Estimates


caseid: Random Effect Estimates

caseid	(Intercept)
1	0.135
2	−0.153
3	−0.114
4	0.122
5	0.138
6	−0.101
7	−0.046
8	0.269
9	−0.082
10	−0.023
11	−0.034
12	−0.146
13	0.009
14	0.188
15	−0.171
16	0.277
17	−0.378
18	0.040
19	−0.195
20	0.115
21	0.119
22	0.067
23	−0.061
24	0.113
25	0.043
26	0.199
27	−0.065
28	−0.058
29	−0.130
30	−0.235
31	−0.028
32	0.149
33	0.012
34	0.152
35	−0.007
36	−0.121

Estimated Means and Confidence Intervals

Block	congruency	Mean	95% CI	
			Lower	Upper
3	congruent	6.455	6.406	6.504
6	congruent	6.666	6.618	6.715
3	incongruent	6.509	6.460	6.558
6	incongruent	6.659	6.610	6.707

Plot



Estimated Marginal Means

Block	congruency	Estimate	SE	95% CI	
				Lower	Upper
3	congruent	6.455	0.025	6.406	6.504
6	congruent	6.666	0.025	6.618	6.715
3	incongruent	6.509	0.025	6.460	6.558
6	incongruent	6.659	0.025	6.610	6.707

Descriptive Statistics


```
jaspDescriptives::Descriptives(  
  data = NULL,  
  version = "0.95",  
  formula = ~ Log_rt,  
  distributionPlots = TRUE,  
  qqPlot = TRUE,  
  quartiles = TRUE,  
  skewness = TRUE)
```

Descriptive Statistics

Log_rt	
Valid	22649
Missing	463
Mean	6.591
Std. Deviation	0.406
Skewness	0.817
Std. Error of Skewness	0.016
Minimum	5.342
Maximum	8.505
25th percentile	6.326
50th percentile	6.524
75th percentile	6.802

Distribution Plots

Log_rt



Q-Q Plot

Log_rt

