



MSc Thesis for Health Data Science

A Bayesian Time-Series Framework for Comparing Hospital Pressure Across Irish HSE Health Regions

Lucas Henry, Evan Bergin

Supervisor: Dr. Carl Scarrott

XXXX 2026

Declaration

We hereby certify that this material, which we now submit for assessment on the programme of study leading to the award of Master of Science is entirely our own work and had not been taken from the work of others or Artificial Intelligence, save and to the extent that such work has been cited and acknowledged within the text of our work.

ID: 25252089

Date:

Contents

1	Introduction	3
2	Data	3
2.1	Data Collection	3
3	Methods	4
3.1	Data Collection Pipeline	4
3.2	Response Scaling	4
3.3	Model Development	4
3.4	Posterior Sampling and Bayesian Ranking	5
3.5	Code and Data Availability	6
4	Results	6
4.1	Model Comparison	6
4.1.1	Regional Baseline	6
4.1.2	Week-to-Week Autocorrelation	6
4.1.3	Annual Cycle	6
4.1.4	Regional Rankings	6
5	Discussion	7
6	References	8
7	Appendix	9
7.1	Model Inheritance	9
7.2	Full Model Specifications	9
	Model v0.1	9
	Model v1.1	10
	Model v2.1	10
	Model v2.2	10
	Model v2.3	11
	Model v2.4	11
	Model v3.1	12
	Model v3.2	12
	Model v3.3	13
7.3	Alternative Response Scales	13
7.3.1	Budget scaling	13
7.3.2	Bed scaling	14
7.4	Residual Diagnostics	14
7.5	Posterior Mean Fit	14
7.6	Gelman-Rubin Convergence Summary	15
7.7	Posterior Parameter Estimates	15

1 Introduction

2 Data

2.1 Data Collection

3 Methods

3.1 Data Collection Pipeline

A pipeline was created to extract daily on trolley counts from each HSE hospital and transform them into a csv file. The code for this can be found on github (Henry & Bergin, 2026).



Figure 1: Data pipeline from source to model input.

3.2 Response Scaling

The on trolley counts were standardised to a rate of count per 10,000 people in each health region to control for catchment population size. Counts were standardised to a rate per 10,000 catchment population.

$$y_{i,t} = \frac{\text{weekly trolley count}_{i,t}}{\text{health region catchment population}_i} \times 10,000$$

3.3 Model Development

Models were developed iteratively with increasing complexity as described in Gelman (2004). Model iteration can be visualised in appendix figure 2. Components were added to the model according to the residual patterns, resulting in the base model (2).

The base model likelihood initialises at $t = 1$ and propagates an AR(1) carryover at $t > 1$.

$$y_{i,1} \sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \quad (1)$$

$$y_{i,t} \sim \mathcal{N}(\mu_{i,t} + \mathcal{E}_{i,t}^1, \tau_i^{-1}), \quad t > 1 \quad (2)$$

The AR(1) term captures week-to-week autocorrelation in the residuals.

$$\mathcal{E}_{i,t}^1 = \phi(y_{i,t-1} - \mu_{i,t-1}) \quad (3)$$

The mean function decomposes the expected trolley rate into a baseline level (α_i), an annual cycle ($\mathcal{C}_{i,t}^{52}$), a regional New Year effect ($\mathcal{R}_{i,t}$), and a Mid West funding reset ($\mathcal{W}_{i,t}$).

$$\mu_{i,t} = \alpha_i + \mathcal{C}_{i,t}^{52} + \mathcal{R}_{i,t} + \mathcal{W}_{i,t} \quad (4)$$

The annual cycle is captured with the $\mathcal{C}_{i,t}^{52}$ component.

$$\mathcal{C}_{i,t}^{52} = \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \quad (5)$$

The regional New Year effect allows each region to respond differently across the three week New Year window.

$$\begin{aligned} \mathcal{R}_{i,t} = & \delta_{\text{pre},i} \mathbf{1}(t \bmod 52 = 0) \\ & + \delta_{\text{mid},i} \mathbf{1}(t \bmod 52 = 1) \\ & + \delta_{\text{post},i} \mathbf{1}(t \bmod 52 = 2) \end{aligned} \quad (6)$$

The Mid West reset captures a three week window around August 8th 2024, where numerous cases were deferred.

$$\begin{aligned}\mathcal{W}_{i,t} = & \mathbf{1}(i = \text{HSE Mid West}) \cdot (\sigma_{\text{pre}} \mathbf{1}(t = 86) \\ & + \sigma_{\text{mid}} \mathbf{1}(t = 87) \\ & + \sigma_{\text{post}} \mathbf{1}(t = 88))\end{aligned}\quad (7)$$

Extending the base model described above, two models were added. One with an added AR(2) component.

$$\begin{aligned}y_{i,t} & \sim \mathcal{N}(\mu_{i,t} + \mathcal{E}_{i,t}^1 + \mathcal{E}_{i,t}^2, \tau_i^{-1}) \\ \mu_{i,t} & = \alpha_i + \mathcal{C}_{i,t}^{52} + \mathcal{R}_{i,t} + \mathcal{W}_{i,t}\end{aligned}\quad (8)$$

A second with an added 6-month cycle.

$$\begin{aligned}y_{i,t} & \sim \mathcal{N}(\mu_{i,t} + \mathcal{E}_{i,t}^1, \tau_i^{-1}) \\ \mu_{i,t} & = \alpha_i + \mathcal{C}_{i,t}^{52} + \mathcal{C}_{i,t}^{26} + \mathcal{R}_{i,t} + \mathcal{W}_{i,t}\end{aligned}\quad (9)$$

3.4 Posterior Sampling and Bayesian Ranking

Posterior sampling was performed using *Just another Gibbs sampler (JAGs)* (Plummer, 2003), implemented by pyJAGS (Nowotny, 2024), a Python interface to JAGS. Health region baselines, α_j^s , were used for ranking. Where j is one of the 6 HSE health regions, $j \in \{1, \dots, 6\}$, and s is the draw index, for $s \in \{1, \dots, n\}$ and total samples is defined as $n = 20,000$.

$$A = \begin{bmatrix} \alpha_1^1 & \dots & \alpha_j^1 & \dots & \alpha_6^1 \\ \vdots & & \vdots & & \vdots \\ \alpha_1^s & \dots & \alpha_j^s & \dots & \alpha_6^s \\ \vdots & & \vdots & & \vdots \\ \alpha_1^n & \dots & \alpha_j^n & \dots & \alpha_6^n \end{bmatrix}$$

Ranks follow the posterior ranking approach of Laird & Louis (1989). Where each sample is posterior sample is ranked to form rank distribution, R .

$$R = \begin{bmatrix} r_1^1 & \dots & r_j^1 & \dots & r_6^1 \\ \vdots & & \vdots & & \vdots \\ r_1^s & \dots & r_j^s & \dots & r_6^s \\ \vdots & & \vdots & & \vdots \\ r_1^n & \dots & r_j^n & \dots & r_6^n \end{bmatrix}$$

Where $r_j^s = \text{rank}(\alpha_j^s) = \sum_{i=1}^6 I(\alpha_j^s > \alpha_i^s)$. Given a tie, ranks were increased by $\frac{1}{2}$ steps for each tied value:

$$r_j^s = \sum_{i=1}^6 [\mathbf{1}(\alpha_j^s > \alpha_i^s) + \frac{1}{2} \mathbf{1}(\alpha_j^s = \alpha_i^s)]$$

Then posterior rank averages, $\hat{r}_i$, were calculated taking the mean across samples.

$$\hat{r}_j = \frac{1}{n} \sum_{s=1}^n r_j^s$$

$$\hat{R} = [\hat{r}_1, \hat{r}_2, \hat{r}_3, \hat{r}_4, \hat{r}_5, \hat{r}_6]$$

Then posterior ranked averages were ranked once again to achieve ranked posterior ranked averages were calculated.

$$r_j^* = \text{rank}(\alpha_j^*) = \sum_{i=1}^6 I(\alpha_j^* > \alpha_i^*)$$

$$R^* = [r_1^*, r_2^*, r_3^*, r_4^*, r_5^*, r_6^*]$$

Where $r_j^* = \text{rank}(\hat{r}_j) = \sum_{i=1}^6 I(\hat{r}_j > \hat{r}_i)$.

3.5 Code and Data Availability

[Placeholder: link to GitHub repository, languages and tools used (Python, R, JAGS), and note that source data is publicly available from the HSE Urgent and Emergency Care portal.]

4 Results

4.1 Model Comparison

4.1.1 Regional Baseline

4.1.2 Week-to-Week Autocorrelation

4.1.3 Annual Cycle

4.1.4 Regional Rankings

5 Discussion

6 References

- Gelman, A. (2004). Exploratory Data Analysis for Complex Models. *Journal of Computational and Graphical Statistics*, 13(4), 755–779. <https://doi.org/10.1198/106186004X11435>
- Henry, L., & Bergin, E. (2026). *HSE Urgent Care Report Bayesian Analysis*. <https://github.com/hotsoupisgood/scraping-hse-trolley-data>
- Laird, N. M., & Louis, T. A. (1989). Empirical Bayes ranking methods. *Journal of Educational Statistics*, 14(1), 29–46. <https://doi.org/10.3102/10769986014001029>
- Nowotny, M. (2024). *pyJAGS: A Python interface to JAGS* (Version 2.4.0) [Computer software]. <https://github.com/michaelnowotny/pyjags>
- Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. *Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC 2003)*. <https://www.r-project.org/conferences/DSC-2003/Proceedings/Plummer.pdf>

7 Appendix

7.1 Model Inheritance




Figure 2: Model inheritance diagram. Spine nodes (blue) form the selected model sequence; grey nodes were tested variants. Thick arrows indicate the selected path; thin arrows indicate branches.

7.2 Full Model Specifications

Model v0.1

- AR(1) baseline; no cycle, no event effects

Likelihood

$$y_{i,1} \sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1})$$

$$y_{i,t} \sim \mathcal{N}(\mu_{i,t} + \mathcal{E}_{i,t}^1, \tau_i^{-1}), \quad t > 1$$

Where $\mu_{i,t}$ is the baseline mean for region i at time t .

$$\mu_{i,t} = \alpha_i$$

$\mathcal{E}_{i,t}^1$ is the AR(1) error term.

$$\mathcal{E}_{i,t}^1 = \phi(y_{i,t-1} - \mu_{i,t-1})$$

Priors

$$\begin{aligned}\alpha_i &\sim \mathcal{N}(0, 0.001) \\ \tau_i &\sim \Gamma(0.001, 0.001) \\ \phi &\sim \text{Uniform}(-1, 1)\end{aligned}$$

Model v1.1

- Extends v0.1; adds annual cycle

Likelihood

$$\begin{aligned}\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\ y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\ y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1\end{aligned}$$

Priors

$$\begin{aligned}\alpha_i, \beta_i, \gamma_i &\sim \mathcal{N}(0, 0.001) \\ \tau_i &\sim \Gamma(0.001, 0.001) \\ \phi &\sim \text{Uniform}(-1, 1)\end{aligned}$$

Model v2.1

- Extends v1.1; adds regional New Year effect and Mid West funding reset

Likelihood

$$\begin{aligned}\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\ &\quad + \delta_{\text{pre},i} \mathbf{1}(t \bmod 52 = 0) + \delta_{\text{mid},i} \mathbf{1}(t \bmod 52 = 1) + \delta_{\text{post},i} \mathbf{1}(t \bmod 52 = 2) \\ &\quad + (\sigma_{\text{pre}} \mathbf{1}(t = 86) + \sigma_{\text{mid}} \mathbf{1}(t = 87) + \sigma_{\text{post}} \mathbf{1}(t = 88)) \cdot \text{mw}_i \\ y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\ y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1\end{aligned}$$

Priors

$$\begin{aligned}\alpha_i, \beta_i, \gamma_i &\sim \mathcal{N}(0, 0.001) \\ \delta_{\text{pre},i}, \delta_{\text{mid},i}, \delta_{\text{post},i} &\sim \mathcal{N}(0, 0.001) \\ \sigma_{\text{pre}}, \sigma_{\text{mid}}, \sigma_{\text{post}} &\sim \mathcal{N}(0, 0.001) \\ \tau_i &\sim \Gamma(0.001, 0.001) \\ \phi &\sim \text{Uniform}(-1, 1)\end{aligned}$$

Model v2.2

- Extends v1.1; adds global New Year effect (shared across regions)

Likelihood

$$\begin{aligned}\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\ &\quad + \delta_{\text{pre}} \mathbf{1}(t \bmod 52 = 0) + \delta_{\text{mid}} \mathbf{1}(t \bmod 52 = 1) + \delta_{\text{post}} \mathbf{1}(t \bmod 52 = 2) \\ y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\ y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1\end{aligned}$$

Priors

$$\begin{aligned}\alpha_i, \beta_i, \gamma_i &\sim \mathcal{N}(0, 0.001) \\ \delta_{\text{pre}}, \delta_{\text{mid}}, \delta_{\text{post}} &\sim \mathcal{N}(0, 0.001) \\ \tau_i &\sim \Gamma(0.001, 0.001) \\ \phi &\sim \text{Uniform}(-1, 1)\end{aligned}$$

Model v2.3

- Extends v1.1; adds regional New Year effect (no MW reset)

Likelihood

$$\begin{aligned}\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\ &\quad + \delta_{\text{pre},i} \mathbf{1}(t \bmod 52 = 0) + \delta_{\text{mid},i} \mathbf{1}(t \bmod 52 = 1) + \delta_{\text{post},i} \mathbf{1}(t \bmod 52 = 2) \\ y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\ y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1\end{aligned}$$

Priors

$$\begin{aligned}\alpha_i, \beta_i, \gamma_i &\sim \mathcal{N}(0, 0.001) \\ \delta_{\text{pre},i}, \delta_{\text{mid},i}, \delta_{\text{post},i} &\sim \mathcal{N}(0, 0.001) \\ \tau_i &\sim \Gamma(0.001, 0.001) \\ \phi &\sim \text{Uniform}(-1, 1)\end{aligned}$$

Model v2.4

- Extends v1.1; adds Mid West funding reset (no New Year effect)

Likelihood

$$\begin{aligned}\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\ &\quad + (\sigma_{\text{pre}} \mathbf{1}(t = 86) + \sigma_{\text{mid}} \mathbf{1}(t = 87) + \sigma_{\text{post}} \mathbf{1}(t = 88)) \cdot \text{mw}_i \\ y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\ y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1\end{aligned}$$

Priors

$$\begin{aligned}
\alpha_i, \beta_i, \gamma_i &\sim \mathcal{N}(0, 0.001) \\
\sigma_{\text{pre}}, \sigma_{\text{mid}}, \sigma_{\text{post}} &\sim \mathcal{N}(0, 0.001) \\
\tau_i &\sim \Gamma(0.001, 0.001) \\
\phi &\sim \text{Uniform}(-1, 1)
\end{aligned}$$

Model v3.1

- Extends v2.1; adds 8-week sub-annual cycle

Likelihood

$$\begin{aligned}
\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\
&\quad + \beta_{2,i} \cos\left(\frac{2\pi t}{8}\right) + \gamma_{2,i} \sin\left(\frac{2\pi t}{8}\right) \\
&\quad + \delta_{\text{pre},i} \mathbf{1}(t \bmod 52 = 0) + \delta_{\text{mid},i} \mathbf{1}(t \bmod 52 = 1) + \delta_{\text{post},i} \mathbf{1}(t \bmod 52 = 2) \\
&\quad + (\sigma_{\text{pre}} \mathbf{1}(t = 86) + \sigma_{\text{mid}} \mathbf{1}(t = 87) + \sigma_{\text{post}} \mathbf{1}(t = 88)) \cdot \text{mw}_i \\
y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\
y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1
\end{aligned}$$

Priors

$$\begin{aligned}
\alpha_i, \beta_i, \gamma_i, \beta_{2,i}, \gamma_{2,i} &\sim \mathcal{N}(0, 0.001) \\
\delta_{\text{pre},i}, \delta_{\text{mid},i}, \delta_{\text{post},i} &\sim \mathcal{N}(0, 0.001) \\
\sigma_{\text{pre}}, \sigma_{\text{mid}}, \sigma_{\text{post}} &\sim \mathcal{N}(0, 0.001) \\
\tau_i &\sim \Gamma(0.001, 0.001) \\
\phi &\sim \text{Uniform}(-1, 1)
\end{aligned}$$

Model v3.2

- Extends v2.1; adds 26-week sub-annual cycle

Likelihood

$$\begin{aligned}
\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\
&\quad + \beta_{2,i} \cos\left(\frac{2\pi t}{26}\right) + \gamma_{2,i} \sin\left(\frac{2\pi t}{26}\right) \\
&\quad + \delta_{\text{pre},i} \mathbf{1}(t \bmod 52 = 0) + \delta_{\text{mid},i} \mathbf{1}(t \bmod 52 = 1) + \delta_{\text{post},i} \mathbf{1}(t \bmod 52 = 2) \\
&\quad + (\sigma_{\text{pre}} \mathbf{1}(t = 86) + \sigma_{\text{mid}} \mathbf{1}(t = 87) + \sigma_{\text{post}} \mathbf{1}(t = 88)) \cdot \text{mw}_i \\
y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\
y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi(y_{i,t-1} - \mu_{i,t-1}), \tau_i^{-1}), \quad t > 1
\end{aligned}$$

Priors

$$\begin{aligned}
\alpha_i, \beta_i, \gamma_i, \beta_{2,i}, \gamma_{2,i} &\sim \mathcal{N}(0, 0.001) \\
\delta_{\text{pre},i}, \delta_{\text{mid},i}, \delta_{\text{post},i} &\sim \mathcal{N}(0, 0.001) \\
\sigma_{\text{pre}}, \sigma_{\text{mid}}, \sigma_{\text{post}} &\sim \mathcal{N}(0, 0.001) \\
\tau_i &\sim \Gamma(0.001, 0.001) \\
\phi &\sim \text{Uniform}(-1, 1)
\end{aligned}$$

Model v3.3

- Extends v2.1; replaces AR(1) with AR(2)

Likelihood

$$\begin{aligned}
\mu_{i,t} &= \alpha_i + \beta_i \cos\left(\frac{2\pi t}{52}\right) + \gamma_i \sin\left(\frac{2\pi t}{52}\right) \\
&\quad + \delta_{\text{pre},i} \mathbf{1}(t \bmod 52 = 0) + \delta_{\text{mid},i} \mathbf{1}(t \bmod 52 = 1) + \delta_{\text{post},i} \mathbf{1}(t \bmod 52 = 2) \\
&\quad + (\sigma_{\text{pre}} \mathbf{1}(t = 86) + \sigma_{\text{mid}} \mathbf{1}(t = 87) + \sigma_{\text{post}} \mathbf{1}(t = 88)) \cdot \text{mw}_i \\
y_{i,1} &\sim \mathcal{N}(\mu_{i,1}, \tau_i^{-1}) \\
y_{i,2} &\sim \mathcal{N}(\mu_{i,2} + \phi_1 (y_{i,1} - \mu_{i,1}), \tau_i^{-1}) \\
y_{i,t} &\sim \mathcal{N}(\mu_{i,t} + \phi_1 (y_{i,t-1} - \mu_{i,t-1}) + \phi_2 (y_{i,t-2} - \mu_{i,t-2}), \tau_i^{-1}), \quad t > 2
\end{aligned}$$

Priors

$$\begin{aligned}
\alpha_i, \beta_i, \gamma_i &\sim \mathcal{N}(0, 0.001) \\
\delta_{\text{pre},i}, \delta_{\text{mid},i}, \delta_{\text{post},i} &\sim \mathcal{N}(0, 0.001) \\
\sigma_{\text{pre}}, \sigma_{\text{mid}}, \sigma_{\text{post}} &\sim \mathcal{N}(0, 0.001) \\
\tau_i &\sim \Gamma(0.001, 0.001) \\
\phi_1, \phi_2 &\sim \mathcal{N}(0, 0.01)
\end{aligned}$$

7.3 Alternative Response Scales

The primary analysis uses population-scaled trolley counts (per 10,000 catchment population). Two sensitivity scalings were also fitted to assess robustness of model selection and regional rankings to the choice of denominator.

7.3.1 Budget scaling

Weekly trolley counts scaled by the 2026 HSE regional budget allocation, expressing the outcome as trolleys per €1 billion allocated to the region.

$$y_{i,t} = \frac{\text{weekly trolley count}_{i,t}}{\text{health region budget}_i / 10^6}$$

where health region budget_{*i*} is the 2026 budget allocation for region *i* in €thousands. This controls for funding differentials: a region receiving a larger allocation is expected to handle a proportionally larger patient volume.

Table 1: DIC comparison across the models (trolley count scaled by 2026 regional budget).

Model	Description	Deviance	pD	DIC
v4.1	V3.1 + Partial Pooling	19858.4	79.8	19938.1
v2.1	V1.1 + Global New Year	19920.3	54.9	19975.3
v3.1	V2.2 + Mid West Reset	19899.9	79.4	19979.3
v2.2	V1.1 + Regional New Year	19908.3	77.4	19985.7
v2.3	V1.1 + Mid West Reset	20169.5	50.4	20219.9
v1.1	V0.1 + Annual Cycle	20176.0	48.6	20224.6
v1.5	V1.1 + 8-Week Cycle	20171.4	72.1	20243.5
v1.4	V1.1 + 6-Month Cycle	20195.4	70.6	20266.0
v0.1	AR(1) Only	20253.9	27.0	20280.9
v0.2	V0.1 + AR(2)	20253.9	29.3	20283.2
v1.3	V0.1 + 8-Week Cycle	20248.9	50.6	20299.5
v1.2	V0.1 + 26-Week Cycle	20274.2	48.8	20323.0

7.3.2 Bed scaling

Weekly trolley counts scaled by the number of inpatient beds in each region, expressing the outcome as an overcapacity rate: for every 100 inpatient beds, how many patients were waiting on trolleys that week.

$$y_{i,t} = \frac{\text{weekly trolley count}_{i,t}}{\text{health region inpatient beds}_i} \times 100$$

where health region inpatient beds_{*i*} is the total inpatient bed count across hospitals in region *i*, summed from the June 2025 bed census. Hospitals with missing bed counts were excluded from the regional sum.

Table 2: DIC comparison across models (response scaled by June 2025 health region inpatient bed count).

Model	Description	Deviance	pD	DIC
v3.1	V2.2 + Mid West Reset	11448.9	93.0	11541.9
v4.1	V3.1 + Partial Pooling	11449.1	93.3	11542.4
v2.2	V1.1 + Regional New Year	11470.0	87.3	11557.3
v2.1	V1.1 + Global New Year	11510.1	57.2	11567.3
v2.3	V1.1 + Mid West Reset	11738.6	56.7	11795.3
v1.1	V0.1 + Annual Cycle	11756.7	51.0	11807.7
v1.5	V1.1 + 8-Week Cycle	11751.8	75.1	11826.9
v1.4	V1.1 + 6-Month Cycle	11773.3	75.3	11848.6
v0.1	AR(1) Only	11828.7	26.6	11855.3
v0.2	V0.1 + AR(2)	11828.3	28.6	11856.8
v1.3	V0.1 + 8-Week Cycle	11823.4	50.7	11874.1
v1.2	V0.1 + 26-Week Cycle	11845.5	50.9	11896.4

7.4 Residual Diagnostics

[Residual diagnostic plots for each model.]

7.5 Posterior Mean Fit

[Posterior mean $\mu_{i,t}$ overlaid on observed $y_{i,t}$ for each spine model.]

7.6 Gelman-Rubin Convergence Summary

[Gelman-Rubin $\hat{R}$ summary table for each spine model.]

7.7 Posterior Parameter Estimates

[Posterior means and 95% credible intervals for each spine model.]