

$$\mathbb{S}_{\mathrm{MU}}^{\wedge}$$

Imagine maps:

$$\mathrm{Map}_*^G(A,B)$$

$$\mathrm{Map}_*^G(A,B)$$

where $A,B \in \mathrm{Top}_*^G$ (where Top is a suitable category).

If $X \in \mathrm{Top}_*$ and $Y \in \mathrm{Top}$, we can form the spectra:

$$\Sigma^\infty X, \Sigma_+^\infty Y$$

and for a spectrum $Z \in \mathrm{Sp}$, we can form the pointed space:

$$\Omega^\infty Z$$

THEOREM 1 (Pythagoras). $a^2 + b^2 = c^2$.

PROOF. Trivial □

◀

LEMMA 2 (Zorn's Lemma). *An poset $P \in \mathrm{Poset}$ where all chains are bounded above has a maximal element.*

PROOF. Axiom of choice. □

◀

PROPOSITION 3. $\sqrt[3]{2}$ is not a rational number.

PROOF. Suppose it was. Then, there will be two integers $a, b \in \mathbb{Z}$ such that $a^3 + a^3 = b^3$. We will need the following:

THEOREM 4 (Fermat's Last Theorem, due to Andrew Wiles). *The equation $a^n + b^n = c^n$ has no integer solutions when $n \geq 3$.*

PROOF. Trivial [Theorem 4] □

◀

We see that $\sqrt[3]{2}$ being rational produces a contradiction to Fermat's Last Theorem. Therefore, it must be irrational. [Proposition 3] □

◀

REMARK 5 (Jordan curve theorem). It happens

IDEA. Duh □

◀

$$\mathbb{S}, \mathrm{MU}, \mathrm{tmf}, \mathrm{BP}, T(n), K(n), H\mathbb{Z}, E_n$$

$$S, MU, \mathrm{tmf}, \mathrm{BP}, T(n), K(n), H\mathbb{Z}, E_n$$

CLAIM 6. When $z \in \mathbb{C}$, we have that:

$$e^z = e^{\Re(z)}(\cos \Im(z) + i \sin \Im(z))$$

◀

$$K(\mathbb{Z}) \rightarrow K \rightarrow K(n - \mathrm{cork}_p(\mathcal{F})) \rightarrow K(n) \rightarrow K(\mathbb{Z}) \rightarrow K(\infty)$$

$$\Delta$$

AXIOM (Axiom of Choice). FOR ANY COLLECTION OF NON-EMPTY SETS \mathcal{S} , THERE EXISTS A CHOICE FUNCTION $\phi : \mathcal{S} \rightarrow \bigcup_{s \in \mathcal{S}} s$, SUCH THAT $\phi(s) \in s$. ◀

abcdefghijklmnopqrstuvwxyz
ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz
ABCDEFGHIJKLMNOPQRSTUVWXYZ
αβγδεζηθικλμνξοπρστυφχψω
ΑΒΓΔΕΖΗΘΙΚΑΜΝΞΟΠΡΣΤΥΦΧΨΩ
εδρφ

ΣΠΛΙΟ⊕⊗