

LINEAR ALGEBRA to Digital Image Processing

January 14, 2018

Singular Value Decomposition

This is similar to factorization of integers. For any matrix $A_{m \times n}$, there exist orthogonal matrices U, V and a diagonal matrix Σ such that $A = U\Sigma V^T$

Singular Value Decomposition

This is similar to factorization of integers. For any matrix $A_{m \times n}$, there exist orthogonal matrices U, V and a diagonal matrix Σ such that $A = U\Sigma V^T$

A matrix $A_{m \times n} = [q_1 \ q_2 \dots \ q_n]$

is said to be orthogonal if the columns of A that is the vectors

q_1, q_2, \dots, q_n are orthonormal vectors. This means they should satisfy the condition $q_i \cdot q_j = \delta_{ij}$

Singular Value Decomposition

This is similar to factorization of integers. For any matrix $A_{m \times n}$, there exist orthogonal matrices U, V and a diagonal matrix Σ such that $A = U\Sigma V^T$

A matrix $A_{m \times n} = [q_1 \ q_2 \dots \ q_n]$

is said to be orthogonal if the columns of A that is the vectors q_1, q_2, \dots, q_n are orthonormal vectors. This means they should satisfy the condition $q_i \cdot q_j = \delta_{ij}$

When the matrix $A_{m \times n}$ is orthogonal, it can be easily proved that $A^T A = I_{n \times n}$. However, $AA^T = I$, if and only if the matrix is square.

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Let the rank of $A = r$

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Let the rank of $A = r$

Consider $B = A^T A$.

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Let the rank of $A = r$

Consider $B = A^T A$.

Clearly B is $n \times n$ matrix and is symmetric.

Hence it will have n eigen values and n eigenvectors

- this follows from the fact that a real square symmetric matrix can always be diagonalized.

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Let the rank of $A = r$

Consider $B = A^T A$.

Clearly B is $n \times n$ matrix and is symmetric.

Hence it will have n eigen values and n eigenvectors

- this follows from the fact that a real square symmetric matrix can always be diagonalized.

These eigenvectors are orthogonal to each other.

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Let the rank of $A = r$

Consider $B = A^T A$.

Clearly B is $n \times n$ matrix and is symmetric.

Hence it will have n eigen values and n eigenvectors

- this follows from the fact that a real square symmetric matrix can always be diagonalized.

These eigenvectors are orthogonal to each other.

Let $\{v_1, v_2, \dots, v_n\}$ be an orthonormal basis of R^n consisting of the eigenvectors of $A^T A$. Let the associated eigenvalues be $\lambda_1, \lambda_2, \dots, \lambda_n$

Singular Value Decomposition-Construction

Let us assume that A is an $m \times n$ matrix.

This means that A is a transformation from $R^n \rightarrow R^m$.

Let the rank of $A = r$

Consider $B = A^T A$.

Clearly B is $n \times n$ matrix and is symmetric.

Hence it will have n eigen values and n eigenvectors

- this follows from the fact that a real square symmetric matrix can always be diagonalized.

These eigenvectors are orthogonal to each other.

Let $\{v_1, v_2, \dots, v_n\}$ be an orthonormal basis of R^n consisting of the eigenvectors of $A^T A$. Let the associated eigenvalues be $\lambda_1, \lambda_2, \dots, \lambda_n$

Then, $\|Av_i\|^2 = (Av_i)^T (Av_i) = v_i^T A^T A v_i = v_i^T (A^T A v_i) = v_i^T \lambda_i v_i = \lambda_i \|v_i\|^2 = \lambda_i$

Hence, eigenvalues of $A^T A$ are all non-negative.

Singular Value Decomposition-construction

Assume that they are all arranged as

$$\lambda_1 \geq \lambda_2 \geq \dots \lambda_n \geq 0$$

Singular Value Decomposition-construction

Assume that they are all arranged as

$$\lambda_1 \geq \lambda_2 \geq \dots \lambda_n \geq 0$$

DEFINITION-The singular values of A are the square roots of the eigenvalues of $A^T A$ denoted by $\sigma_1, \sigma_2, \dots, \sigma_n$

Singular Value Decomposition-construction

Assume that they are all arranged as
 $\lambda_1 \geq \lambda_2 \geq \dots \lambda_n \geq 0$

DEFINITION-The singular values of A are the square roots of the eigenvalues of $A^T A$ denoted by $\sigma_1, \sigma_2, \dots, \sigma_n$

Hence, $\sigma_i = \sqrt{\lambda_i} = \|Av_i\|, i = 1, 2, \dots, n$ and they are the lengths of the vectors Av_i

Ranks of $A^T A$ and A are same

We shall now prove that $\text{rank}(A^T A) = \text{rank}(A)$

Ranks of $A^T A$ and A are same

We shall now prove that $\text{rank}(A^T A) = \text{rank}(A)$

Towards this, we shall prove that $\text{Null}(A^T A) = \text{Null}(A)$

Ranks of $A^T A$ and A are same

We shall now prove that $\text{rank}(A^T A) = \text{rank}(A)$

Towards this, we shall prove that $\text{Null}(A^T A) = \text{Null}(A)$

Let $x \in N(A) \Rightarrow Ax = 0$.

Ranks of $A^T A$ and A are same

We shall now prove that $\text{rank}(A^T A) = \text{rank}(A)$

Towards this, we shall prove that $\text{Null}(A^T A) = \text{Null}(A)$

Let $x \in N(A) \Rightarrow Ax = 0$.

Hence, $A^T Ax = 0 \Rightarrow x \in N(A^T A)$

Ranks of $A^T A$ and A are same

We shall now prove that $\text{rank}(A^T A) = \text{rank}(A)$

Towards this, we shall prove that $\text{Null}(A^T A) = \text{Null}(A)$

Let $x \in N(A) \Rightarrow Ax = 0$.

Hence, $A^T Ax = 0 \Rightarrow x \in N(A^T A)$

This implies that $N(A) \subset N(A^T A) \dots (1)$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Hence, from (2) and (3), we get $Ax = 0 \Rightarrow x \in N(A)$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Hence, from (2) and (3), we get $Ax = 0 \Rightarrow x \in N(A)$

This proves that $N(A^T A) \subset N(A) \dots (4)$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Hence, from (2) and (3), we get $Ax = 0 \Rightarrow x \in N(A)$

This proves that $N(A^T A) \subset N(A) \dots (4)$

From (1) and (4), we get $N(A^T A) = N(A)$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Hence, from (2) and (3), we get $Ax = 0 \Rightarrow x \in N(A)$

This proves that $N(A^T A) \subset N(A) \dots (4)$

From (1) and (4), we get $N(A^T A) = N(A)$

Hence, $\dim(N(A^T A)) = \dim(N(A))$

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Hence, from (2) and (3), we get $Ax = 0 \Rightarrow x \in N(A)$

This proves that $N(A^T A) \subset N(A) \dots (4)$

From (1) and (4), we get $N(A^T A) = N(A)$

Hence, $\dim(N(A^T A)) = \dim(N(A))$

We also know that for any matrix, $\text{rank}(A) + \dim(N(A)) = \text{number of columns of } A$.

Singular Value Decomposition

Let now, $x \in N(A^T A) \Rightarrow A^T A x = 0$

This means $A^T(Ax) = 0 \Rightarrow Ax \in N(A^T) \dots (2)$

But $Ax \in C(A) = R(A^T) \dots (3)$

But we know that the Null space and Row space of any matrix are orthogonal to each other.

Hence, from (2) and (3), we get $Ax = 0 \Rightarrow x \in N(A)$

This proves that $N(A^T A) \subset N(A) \dots (4)$

From (1) and (4), we get $N(A^T A) = N(A)$

Hence, $\dim(N(A^T A)) = \dim(N(A))$

We also know that for any matrix, $\text{rank}(A) + \dim(N(A)) = \text{number of columns of } A$.

Since, both $A^T A$ and A have the same number of columns n , this means $r(A^T A) = r(A)$

Proved.

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

However, $\text{rank}(\text{Col}(A^T A)) = r$

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

However, $\text{rank}(\text{Col}(A^T A)) = r$

This implies only r of these vectors $A^T A v_i$ can be linearly independent.

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

However, $\text{rank}(\text{Col}(A^T A)) = r$

This implies only r of these vectors $A^T A v_i$ can be linearly independent.

This is same as saying that only r of the vectors $\lambda_i v_i$ are linearly independent.

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

However, $\text{rank}(\text{Col}(A^T A)) = r$

This implies only r of these vectors $A^T A v_i$ can be linearly independent.

This is same as saying that only r of the vectors $\lambda_i v_i$ are linearly independent.

Since all the $v_i, i = 1, 2, \dots, n$ are linearly independent, this is possible if and only if only r of the λ_i 's are non-zeroes.

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

However, $\text{rank}(\text{Col}(A^T A)) = r$

This implies only r of these vectors $A^T A v_i$ can be linearly independent.

This is same as saying that only r of the vectors $\lambda_i v_i$ are linearly independent.

Since all the $v_i, i = 1, 2, \dots, n$ are linearly independent, this is possible if and only if only r of the λ_i 's are non-zeroes.

We arrange the λ_i 's such that

$\lambda_1 \geq \lambda_2 \geq \dots \lambda_r > 0$ and $\lambda_i = 0, i = r + 1, \dots, n$

Singular Value Decomposition-construction

Since v_i are eigenvectors of $A^T A$, therefore, $A^T A v_i = \lambda_i v_i$

Now, $v_i, i = 1, 2, \dots, n$ are orthogonal to each other and hence they form a linearly independent set.

Hence, $A^T A v_i = \lambda_i v_i, i = 1, 2, \dots, n$ also form a linearly independent set.

However, $\text{rank}(\text{Col}(A^T A)) = r$

This implies only r of these vectors $A^T A v_i$ can be linearly independent.

This is same as saying that only r of the vectors $\lambda_i v_i$ are linearly independent.

Since all the $v_i, i = 1, 2, \dots, n$ are linearly independent, this is possible if and only if only r of the λ_i 's are non-zeroes.

We arrange the λ_i 's such that

$\lambda_1 \geq \lambda_2 \geq \dots \lambda_r > 0$ and $\lambda_i = 0, i = r + 1, \dots, n$

Consequently, $\sigma_i > 0, i = 1, 2, \dots, r$ and $\sigma_i = 0, i = r + 1, \dots, n$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i} \quad i = 1, 2, \dots, r$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Form the matrices $V = [v_1 \ v_2 \ \dots v_r]$

$U = [u_1 \ u_2 \ \dots u_r]$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}$ $i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Form the matrices $V = [v_1 \ v_2 \ \dots v_r]$

$U = [u_1 \ u_2 \ \dots u_r]$

Now $AV = A[v_1 \ v_2 \ \dots v_r]$

$= [Av_1 \ Av_2 \ \dots Av_r] = [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r]$

$$= [u_1 \ u_2 \ \dots u_r] \begin{bmatrix} \sigma_1 & 0 & 0 \dots & 0 \\ 0 & \sigma_2 & 0 \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sigma_r \end{bmatrix}$$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Form the matrices $V = [v_1 \ v_2 \ \dots v_r]$

$U = [u_1 \ u_2 \ \dots u_r]$

Now $AV = A[v_1 \ v_2 \ \dots v_r]$

$= [Av_1 \ Av_2 \ \dots Av_r] = [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r]$

$= [u_1 \ u_2 \ \dots u_r] \begin{bmatrix} \sigma_1 & 0 & 0 \dots & 0 \\ 0 & \sigma_2 & 0 \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sigma_r \end{bmatrix}$

Check: $A_{m \times n} V_{n \times r} = U_{m \times r} \Sigma_{r \times r}$

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Form the matrices $V = [v_1 \ v_2 \ \dots v_r]$

$U = [u_1 \ u_2 \ \dots u_r]$

Now $AV = A[v_1 \ v_2 \ \dots v_r]$

$= [Av_1 \ Av_2 \ \dots Av_r] = [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r]$

$$= [u_1 \ u_2 \ \dots u_r] \begin{bmatrix} \sigma_1 & 0 & 0 \dots & 0 \\ 0 & \sigma_2 & 0 \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sigma_r \end{bmatrix}$$

Check: $A_{m \times n} V_{n \times r} = U_{m \times r} \Sigma_{r \times r}$

Clearly V is orthogonal and so is U .

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Form the matrices $V = [v_1 \ v_2 \ \dots v_r]$

$U = [u_1 \ u_2 \ \dots u_r]$

Now $AV = A[v_1 \ v_2 \ \dots v_r]$

$= [Av_1 \ Av_2 \ \dots Av_r] = [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r]$

$$= [u_1 \ u_2 \ \dots u_r] \begin{bmatrix} \sigma_1 & 0 & 0 \dots & 0 \\ 0 & \sigma_2 & 0 \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sigma_r \end{bmatrix}$$

Check: $A_{m \times n} V_{n \times r} = U_{m \times r} \Sigma_{r \times r}$

Clearly V is orthogonal and so is U .

Σ is a square diagonal matrix, with the diagonal elements being the singular values of A .

Singular Value Decomposition-construction

Define $u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, i = 1, 2, \dots, r$

Clearly $v_i \in R^n, i = 1, 2, \dots, n$ and $u_i \in R^m, i = 1, 2, \dots, r$

Also $\|u_i\| = 1, i = 1, 2, \dots, r$ and $\|v_i\| = 1, i = 1, 2, \dots, n$

Form the matrices $V = [v_1 \ v_2 \ \dots v_r]$

$U = [u_1 \ u_2 \ \dots u_r]$

Now $AV = A[v_1 \ v_2 \ \dots v_r]$

$= [Av_1 \ Av_2 \ \dots Av_r] = [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r]$

$$= [u_1 \ u_2 \ \dots u_r] \begin{bmatrix} \sigma_1 & 0 & 0 \dots & 0 \\ 0 & \sigma_2 & 0 \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sigma_r \end{bmatrix}$$

Check: $A_{m \times n} V_{n \times r} = U_{m \times r} \Sigma_{r \times r}$

Clearly V is orthogonal and so is U .

Σ is a square diagonal matrix, with the diagonal elements being the singular values of A .

However, V and U are not square!!

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

For U matrix, we have to add $m - r$ orthonormal vectors.

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

For U matrix, we have to add $m - r$ orthonormal vectors.

This is done by extending the r orthonormal vectors $u_i, i = 1, \dots, r$ to an orthonormal basis of R^m . This is always possible by using Gram-Schmidt Orthogonalisation process.

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

For U matrix, we have to add $m - r$ orthonormal vectors.

This is done by extending the r orthonormal vectors $u_i, i = 1, \dots, r$ to an orthonormal basis of R^m . This is always possible by using Gram-Schmidt Orthogonalisation process.

Hence, we now define

$$V = [v_1 \ v_2 \ \dots v_r \ v_{r+1} \ \dots v_n]$$

$$U = [u_1 \ u_2 \ \dots u_r \ u_{r+1} \ \dots u_m]$$

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

For U matrix, we have to add $m - r$ orthonormal vectors.

This is done by extending the r orthonormal vectors $u_i, i = 1, \dots, r$ to an orthonormal basis of R^m . This is always possible by using Gram-Schmidt Orthogonalisation process.

Hence, we now define

$$V = [v_1 \ v_2 \ \dots v_r \ v_{r+1} \ \dots v_n]$$

$$U = [u_1 \ u_2 \ \dots u_r \ u_{r+1} \ \dots u_m]$$

Now both the matrices V and U are square orthogonal matrices of orders $n \times n$ and $m \times m$ respectively

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

For U matrix, we have to add $m - r$ orthonormal vectors.

This is done by extending the r orthonormal vectors $u_i, i = 1, \dots, r$ to an orthonormal basis of R^m . This is always possible by using Gram-Schmidt Orthogonalisation process.

Hence, we now define

$$V = [v_1 \ v_2 \ \dots v_r \ v_{r+1} \ \dots v_n]$$

$$U = [u_1 \ u_2 \ \dots u_r \ u_{r+1} \ \dots u_m]$$

Now both the matrices V and U are square orthogonal matrices of orders $n \times n$ and $m \times m$ respectively

$$\text{Now, } AV = A[v_1 \ v_2 \ \dots v_r \ v_{r+1} \ \dots v_n]$$

Singular Value Decomposition-construction

We add the remaining $n - r$ v vectors as columns to V .

Clearly for each of these new vectors $Av_i = 0, i = r + 1, \dots, n$

For U matrix, we have to add $m - r$ orthonormal vectors.

This is done by extending the r orthonormal vectors $u_i, i = 1, \dots, r$ to an orthonormal basis of R^m . This is always possible by using Gram-Schmidt Orthogonalisation process.

Hence, we now define

$$V = [v_1 \ v_2 \ \dots v_r \ v_{r+1} \ \dots v_n]$$

$$U = [u_1 \ u_2 \ \dots u_r \ u_{r+1} \ \dots u_m]$$

Now both the matrices V and U are square orthogonal matrices of orders $n \times n$ and $m \times m$ respectively

$$\begin{aligned} \text{Now, } AV &= A[v_1 \ v_2 \ \dots v_r \ v_{r+1} \ \dots v_n] \\ &= [Av_1 \ Av_2 \ \dots Av_r \ Av_{r+1} \ \dots Av_n] \end{aligned}$$

Singular Value Decomposition-construction

$$= [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r \ 0 \ \dots 0]$$

Singular Value Decomposition-construction

$$= [\sigma_1 u_1 \ \sigma_2 u_2 \ \dots \sigma_r u_r \ 0 \ \dots 0]$$

$$= [u_1 \ u_2 \ \dots u_r \ u_{r+1} \ \dots u_m]$$

Singular Value Decomposition-construction

$$\begin{aligned}
 &= [\sigma_1 u_1 \quad \sigma_2 u_2 \quad \dots \sigma_r u_r \quad 0 \quad \dots 0] \\
 &= [u_1 \quad u_2 \quad \dots u_r \quad u_{r+1} \quad \dots u_m] \begin{bmatrix} \sigma_1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & \sigma_2 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \sigma_r & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \end{bmatrix}
 \end{aligned}$$

Singular Value Decomposition-construction

$$\begin{aligned}
 &= [\sigma_1 u_1 \quad \sigma_2 u_2 \quad \dots \sigma_r u_r \quad 0 \quad \dots 0] \\
 &= [u_1 \quad u_2 \quad \dots u_r \quad u_{r+1} \quad \dots u_m] \begin{bmatrix} \sigma_1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & \sigma_2 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \sigma_r & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \end{bmatrix}
 \end{aligned}$$

i.e., $A_{m \times n} V_{n \times n} = U_{m \times m} \Sigma_{m \times n}$

The matrix Σ has m rows and n columns, in the first r rows and r columns, the diagonal elements contain the singular values of A . The remaining rows and columns are zeroes.

Singular Value Decomposition-construction

Observations: $v_i, i = r + 1, \dots, n$ form a basis for the Null Space of $A^T A$

Singular Value Decomposition-construction

Observations: $v_i, i = r + 1, \dots, n$ form a basis for the Null Space of $A^T A$
Since V is now orthogonal and square it can be inverted and in fact
 $V^{-1} = V^T$

Singular Value Decomposition-construction

Observations: $v_i, i = r + 1, \dots, n$ form a basis for the Null Space of $A^T A$
Since V is now orthogonal and square it can be inverted and in fact $V^{-1} = V^T$

Hence, multiplying both sides of the above equation by V^{-1} from the right, we get

$$A = U\Sigma V^{-1} = U\Sigma V^T$$

Singular Value Decomposition-construction

Observations: $v_i, i = r + 1, \dots, n$ form a basis for the Null Space of $A^T A$.
Since V is now orthogonal and square it can be inverted and in fact $V^{-1} = V^T$

Hence, multiplying both sides of the above equation by V^{-1} from the right, we get

$$A = U\Sigma V^{-1} = U\Sigma V^T$$

The last result is by virtue of V being square orthogonal.

Singular Value Decomposition

Let us expand the result $A = U\Sigma V^T$

$$U = [u_{col_1} \quad u_{col_2} \quad \dots \quad u_{col_n}], \quad \Sigma = \begin{bmatrix} \sigma_1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & \sigma_2 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \sigma_r & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \end{bmatrix}$$

$$V^T = \begin{bmatrix} v_{row_1}^T \\ v_{row_2}^T \\ \dots \\ v_{row_n}^T \end{bmatrix}$$

Singular Value Decomposition

Let us expand the result $A = U\Sigma V^T$

$$U = \begin{bmatrix} u_{col_1} & u_{col_2} & \dots & u_{col_n} \end{bmatrix}, \Sigma = \begin{bmatrix} \sigma_1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & \sigma_2 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \sigma_r & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \end{bmatrix}$$

$$V^T = \begin{bmatrix} v_{row_1}^T \\ v_{row_2}^T \\ \dots \\ v_{row_n}^T \end{bmatrix}$$

$$\text{Then, } U\Sigma V^T = \sigma_1 u_{col_1} v_{row_1}^T + \sigma_2 u_{col_2} v_{row_2}^T + \dots + \sigma_r u_{col_r} v_{row_r}^T$$

Singular Value Decomposition

Let us expand the result $A = U\Sigma V^T$

$$U = \begin{bmatrix} u_{col_1} & u_{col_2} & \dots & u_{col_n} \end{bmatrix}, \Sigma = \begin{bmatrix} \sigma_1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & \sigma_2 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \sigma_r & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \end{bmatrix}$$
$$V^T = \begin{bmatrix} v_{row_1}^T \\ v_{row_2}^T \\ \dots \\ v_{row_n}^T \end{bmatrix}$$

Then, $U\Sigma V^T = \sigma_1 u_{col_1} v_{row_1}^T + \sigma_2 u_{col_2} v_{row_2}^T + \dots + \sigma_r u_{col_r} v_{row_r}^T$

The remaining terms in the summation are zeroes, by virtue of

$\sigma_i = 0, i = r + 1, \dots, n$

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$,

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$, and since a basis of a vector space is not unique, these $n - r$ vectors are not unique,

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$, and since a basis of a vector space is not unique. these $n - r$ vectors are not unique, and hence V is not unique.

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$, and since a basis of a vector space is not unique. these $n - r$ vectors are not unique, and hence V is not unique.

In case of U , the last $m - r$ vectors are obtained by extending the set $\{u_1, u_2, \dots, u_r\}$ to an orthonormal basis of R^m

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$, and since a basis of a vector space is not unique. these $n - r$ vectors are not unique, and hence V is not unique.

In case of U , the last $m - r$ vectors are obtained by extending the set $\{u_1, u_2, \dots, u_r\}$ to an orthonormal basis of R^m and since this process is not unique,

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$, and since a basis of a vector space is not unique. these $n - r$ vectors are not unique, and hence V is not unique.

In case of U , the last $m - r$ vectors are obtained by extending the set $\{u_1, u_2, \dots, u_r\}$ to an orthonormal basis of R^m and since this process is not unique, therefore U is not unique.

Singular Value Decomposition -

Each term on the RHS is an $m \times n$ matrix of rank 1. Hence A is expressed as a sum of rank 1 matrices.

The importance of each term is given by the magnitude of the corresponding singular value. This fact can be used for data compression.

- NOTE U and V are not unique -the last $n - r$ vectors of V form a basis of the null space of $A^T A$, and since a basis of a vector space is not unique. these $n - r$ vectors are not unique, and hence V is not unique.

In case of U , the last $m - r$ vectors are obtained by extending the set $\{u_1, u_2, \dots, u_r\}$ to an orthonormal basis of R^m and since this process is not unique, therefore U is not unique.

However, Σ is unique.

SVD to Image Processing

One of the applications of Singular Value Decomposition to Image Processing is in Image Compression.

SVD to Image Processing

One of the applications of Singular Value Decomposition to Image Processing is in Image Compression.

Other areas of Image Processing where SVD can be used are noise filtering, image matching etc.

SVD to Image Processing

One of the applications of Singular Value Decomposition to Image Processing is in Image Compression.

Other areas of Image Processing where SVD can be used are noise filtering, image matching etc.

We shall talk only of Image Compression.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

The spatial variables are discretized to give different picture elements, called pixels, and the intensity function is discretized in some suitable range. In monochrome images, discretized intensity is called gray level

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

The spatial variables are discretized to give different picture elements, called pixels, and the intensity function is discretized in some suitable range. In monochrome images, discretized intensity is called gray level. Digital images are stored in computers and occupy some disk space.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

The spatial variables are discretized to give different picture elements, called pixels, and the intensity function is discretized in some suitable range. In monochrome images, discretized intensity is called gray level. Digital images are stored in computers and occupy some disk space. which depends upon the 'fineness' of spatial and intensity information.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

The spatial variables are discretized to give different picture elements, called pixels, and the intensity function is discretized in some suitable range. In monochrome images, discretized intensity is called gray level

Digital images are stored in computers and occupy some disk space. which depends upon the 'fineness' of spatial and intensity information.

The gray level at any pixel represents the average intensity in the region covered by that particular pixel.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

The spatial variables are discretized to give different picture elements, called pixels, and the intensity function is discretized in some suitable range. In monochrome images, discretized intensity is called gray level

Digital images are stored in computers and occupy some disk space. which depends upon the 'fineness' of spatial and intensity information.

The gray level at any pixel represents the average intensity in the region covered by that particular pixel.

Finer the pixel size, which is called Resolution, better will be the image clarity.

What is an image?

An image may be defined as a two-dimensional function $f(x, y)$ where x, y are spatial coordinates and the amplitude f is called the intensity of the image at that point.

A digital image is the discrete version of an image-all the three x, y, f are discretized.

The spatial variables are discretized to give different picture elements, called pixels, and the intensity function is discretized in some suitable range. In monochrome images, discretized intensity is called gray level

Digital images are stored in computers and occupy some disk space. which depends upon the 'fineness' of spatial and intensity information.

The gray level at any pixel represents the average intensity in the region covered by that particular pixel.

Finer the pixel size, which is called Resolution, better will be the image clarity. Similarly, more the gray level range of an image, better will be the discriminability of the image.




Figure: 1. Images at Resolutions 16Km and 8Km






Figure: 2. Image at Resolution 4 Km


Result Satellite Image



1 SingularValue




1 SValue CR=202



18 SingularValues



18 SValues CR=11




35 Singular Values




35 SValues CR=5.8

Reconstructed and Error Images



Result Satellite Image




52 SValue CR=3.9



69 SValues CR=2.9




86 SValues CR=2.3




Reconstructed and Error Images

CompressionRatio/RMSError vs No of Singular Values



Results-Flower



Reconstructed and Error Images with Singular Values and Comp. Ratio

