

Homework 03

Helena Parsons

Part 1: Setting up tasks

Reading in packages

```
library(tidyverse)
library(here)
library(janitor)
library(readxl)
```

Creating data object

```
# creating new object for temp-kelp data
kelp <-
  # reading csv
  read_csv(
    # directing function to data file
    here("data", "temp-kelp.csv"))

# creating new object for personal data
job_search <-
  # reading csv
  read_csv(
    # directing function to data file
    here("data", "personal-data.csv"))
```

Part 2: Problems

Problem 1: Giant kelp fronds

*Climate change threatens all ecosystems, including giant kelp (*Macrocystis pyrifera*) forests.*

You are interested in the potential relationship between ocean temperature (measured in ° C) and giant kelp frond elongation rate (also known as growth rate, measured in cm day⁻¹).

You conduct a field study in which you track giant kelp individuals growing in different temperatures (on average) and measure their elongation rates. Admittedly, this is not a perfect study, but you do the best with what you have!

1a. An appropriate test

In 1-3 sentences, name the appropriate test(s) to determine the strength of the relationship between temperature and giant kelp frond elongation rate (hint: there are two). Describe the differences between the two tests.

Be specific in your response to demonstrate your understanding of the variables in this question.

Because I solely want to understand the strength of the relationship between temperature and giant kelp frond elongation rate, I want to use a correlation test.

A Pearson's r is used when there is a linear relationship between two variables. A Spearman's ρ is used when the relationship between two variables is monotonic.

1b. Create a visualization

Create a visualization that would be appropriate for showing the relationship between temperature and giant kelp frond elongation rate.

In addition to using the correct geometries, be sure to:


- relabel the x- and y-axes and include units*
- use different colors from the ggplot() defaults*
- use a different theme from the ggplot() default*

```

# creating a new object
temp_kelp_visual <-
  # base layer: ggplot
  ggplot(
    # df: kelp
    data = kelp,
    # x-axis: temp_c
    # y-axis: kelp_elong
    mapping = aes(x = temp_c,
                  y = kelp_elong)) +
  # first layer: scatterplot
  geom_point(
    # changing color of points
    color = "magenta4") +
  # changing theme
  theme_classic() +
  # changing title and axes titles
  labs(title = "Kelp frond elongation rates decrease as temp increases",
        x = expression("Temperature (*~degree*C*")"),
        y = expression("Giant kelp frond elongation rates (cm/day~-1)"))

# displaying object
temp_kelp_visual

```




1c. Check your assumptions and run your test

Assumptions


To run a Pearson's r , you want to ensure that there is a linear relationship between the two variables. From the scatter plot I created in part 1b, I can conclude that the relationship mostly appears to be linear.

```
# displaying object from 1b  
temp_kelp_visual
```




I also want to make sure that the data is normally distributed. For this, I can create a QQ plot.

```
qqplot_temp <- ggplot(data = kelp,  
  aes(sample = temp_c)) +  
  stat_qq(distribution=qnorm) +  
  stat_qq_line( col = "black")  
  
qqplot_temp
```



```
qqplot_kelp <- ggplot(data = kelp,  
  aes(sample = kelp_elong)) +  
  stat_qq(distribution=qnorm) +  
  stat_qq_line( col = "black")  
  
qqplot_kelp
```



It seems that the variables are mostly normally distributed.

Additionally, the variables of kelp frond elongation rates and temperature need to be continuous, which they are since they are not integers, categorical, or ordinal data.

Finally, the observations also need to be independent, which is true because the observations of kelp growth rates do not all come from the same individual.

All of our assumptions have been checked so we can now run the Pearson's r .

Pearson's r test