

CSCI567 Machine Learning (Fall 2016)

Dr. Yan Liu

yanliu.cs@usc.edu

September 26, 2016

Outline

- 1 Linear regression
 - Motivation
 - Algorithm
 - Univariate solution
 - Probabilistic interpretation
 - Solution
 - Multivariate solution in matrix form

- 2 Perceptron

Regression

Predicting a continuous outcome variable

- Predicting a company's future stock price using its past and existing financial info
- Predicting the amount of rain fall
- Predicting ...

Regression

Predicting a continuous outcome variable

- Predicting a company's future stock price using its past and existing financial info
- Predicting the amount of rain fall
- Predicting ...


Key difference from classification

- We measure prediction errors differently.
- This will lead to quite different learning models and algorithms.

Ex: predicting the sale price of a house

Retrieve historical sales records

(This will be our training data)



Features used to predict

[Overview](#)
[Property Details](#)
[Tour Insights](#)
[Property History](#)
[Public Records](#)
[Activity](#)
[Schools](#)

Five unit apartment complex within 2 blocks of USC campus, Gate #6. Great for students (most student leases have parents as guarantors). Most USC students live off campus, so housing units like this are always fully leased. Situated on a gated, corner lot, and across from an elementary school, this complex was recently renovated, and has in-unit laundry hook ups, wall-unit AC, and 12 parking spaces. It is within a DPS (Department of Public Safety) and Campus Cruiser patrolled area. This is a great income generating property, not to be missed!

Property Type: Multi-Family
Style: Two Level, Low Rise
Community: Downtown Los Angeles
County: [Los Angeles](#)
MLS#: 22176741

Property Details for 3620 South BUDLONG, Los Angeles, CA 90007

Details provided by i-Tech MLS and may not match the public record. [Learn More](#)

Interior Features

Kitchen Information

- Remodeled
- Oven, Range

Multi-Unit Information

Community Features

- Units in Complex (Total): 5

Multi-Family Information

- # Leased: 5
- # of Buildings: 1
- Owner Pays Water
- Tenant Pays Electricity, Tenant Pays Gas

Unit 1 Information

- # of Beds: 2
- # of Baths: 1
- Unfurnished
- Monthly Rent: \$1,700

Laundry Information

- Inside Laundry

Heating & Cooling

- Wall Cooling Unit(s)

Unit 2 Information

- # of Beds: 3
- # of Baths: 1
- Unfurnished
- Monthly Rent: \$2,250

Unit 3 Information

- Unfurnished

Unit 4 Information

- # of Beds: 3
- # of Baths: 1
- Unfurnished

- Monthly Rent: \$2,350

Unit 5 Information

- # of Beds: 3
- # of Baths: 2
- Unfurnished
- Monthly Rent: \$2,325

Unit 6 Information

- # of Beds: 3
- # of Baths: 1
- Monthly Rent: \$2,250

Property / Lot Details

Property Features

- Automatic Gate, Card/Code Access

Lot Information

- Lot Size (Sq. Ft.): 9,649
- Lot Size (Acres): 0.2215
- Lot Size Source: Public Records

- Automatic Gate, Lawn, Sidewalks
- Corner Lot, Near Public Transit

Property Information

- Updated/Renovated
- Square Footage Source: Public Records

- Tax Parcel Number: 5040017019

Parking / Garage, Exterior Features, Utilities & Financing

Parking Information

- # of Parking Spaces (Total): 12
- Parking Space
- Garage

Building Information

- Total Floors: 2

Utility Information

- Green Certification Rating: 0.00
- Green Location: Transportation, Walkability
- Green Walk Score: 0
- Green Year Certified: 0

Financial Information

- Capitalization Rate (%): 6.25
- Actual Annual Gross Rent: \$126,331
- Gross Rent Multiplier: 11.29

Location Details, Misc. Information & Listing Information

Location Information

- Cross Streets: W 38th Pl


Expense Information

- Operating: \$37,664

Listing Information

- Listing Terms: Cash, Cash To Existing Loan
- Buyer Financing: Cash


Correlation between square footage and sale price



(Unlike classification, the colors of the dots in this scatterplot do not mean anything.)

Possibly linear relationship

Sale price \approx price_per_sqft \times square_footage + fixed_expense



How to learn the unknown parameters?

training data (past sales record)

sqft	sale price
2000	800K
2100	907K
1100	312K
5500	2,600K
...	...

Reduce prediction error

How to measure errors?

- The classification error(*hit* or *miss*) is not appropriate for continuous outcomes.
- We can look at the *absolute* difference: $|\text{prediction} - \text{sale price}|$

However, for simplicity, we look at the *squared* errors:
 $(\text{prediction} - \text{sale price})^2$

sqft	sale price	prediction	error	squared error
2000	800K	710K	80K	$80K^2$
2100	907K	800K	107K	$107K^2$
1100	312K	350K	38K	$38K^2$
5500	2,600K	2,600K	0	0
...	...			

Minimize squared errors

Our model

Sale price = price_per_sqft \times square_footage + fixed_expense + unexplainable_stuff

Training data

sqft	sale price	prediction	error	squared error
2000	800K	720K	80K	$80K^2$
2100	907K	800K	107K	$107K^2$
1100	312K	350K	38K	$38K^2$
5500	2,600K	2,600K	0	0
...	...			
Total				$80K^2 + 107K^2 + 38K^2 + 0 + \dots$

Aim

Adjust price_per_sqft and fixed_expense such that the sum of the squared error is minimized — i.e., the residual/unexplainable_stuff is

Linear regression

Setup

- Input: $\mathbf{x} \in \mathbb{R}^D$ (covariates, predictors, features, etc)
- Output: $y \in \mathbb{R}$ (responses, targets, outcomes, outputs, etc)
- Training data: $\mathcal{D} = \{(\mathbf{x}_n, y_n), n = 1, 2, \dots, N\}$
- Model: $f : \mathbf{x} \rightarrow y$, with $f(\mathbf{x}) = w_0 + \sum_d w_d x_d = w_0 + \mathbf{w}^T \mathbf{x}$
 $\mathbf{w} = [w_1 \ w_2 \ \cdots \ w_D]^T$ is called *weights*, *parameters*, or *parameter vector*

w_0 is called *bias*.

We also sometimes call $\tilde{\mathbf{w}} = [w_0 \ w_1 \ w_2 \ \cdots \ w_D]^T$ parameters too!
So please pay attention to contexts when you read papers, textbooks, or assigned reading material.

Goal

Minimize prediction error as much as possible

- Residual sum of squares

$$RSS(\tilde{\mathbf{w}}) = \sum_n [y_n - f(\mathbf{x}_n)]^2 = \sum_n [y_n - (w_0 + \sum_d w_d x_{nd})]^2$$

- Other definitions of errors are also possible
We will see an example very soon.

A simple case: x is just one-dimensional

Identify stationary points, by taking derivative with respect to parameters, and setting to zeroes

$$\begin{cases} \frac{\partial RSS(\tilde{\mathbf{w}})}{\partial w_0} = 0 \\ \frac{\partial RSS(\tilde{\mathbf{w}})}{\partial w_1} = 0 \end{cases} \Rightarrow \begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix} \begin{pmatrix} w_0 \\ w_1 \end{pmatrix} = \begin{pmatrix} \sum_n y_n \\ \sum_n x_n y_n \end{pmatrix}$$

Why minimizing RSS is a sensible thing?

Probabilistic interpretation

- Noisy observation model

$$Y = w_0 + w_1 X + \eta$$

where $\eta \sim N(0, \sigma^2)$ is a Gaussian random variable

- Likelihood of one training sample (x_n, y_n)

$$p(y_n|x_n) = N(w_0 + w_1 x_n, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{[y_n - (w_0 + w_1 x_n)]^2}{2\sigma^2}}$$

Probabilistic interpretation (cont'd)

Log-likelihood of the training data \mathcal{D} (assuming i.i.d)

$$\log P(\mathcal{D}) = \log \prod_{n=1}^N p(y_n|x_n) = \sum_n \log p(y_n|x_n)$$

Probabilistic interpretation (cont'd)

Log-likelihood of the training data \mathcal{D} (assuming i.i.d)

$$\begin{aligned}\log P(\mathcal{D}) &= \log \prod_{n=1}^N p(y_n|x_n) = \sum_n \log p(y_n|x_n) \\ &= \sum_n \left\{ -\frac{[y_n - (w_0 + w_1 x_n)]^2}{2\sigma^2} - \log \sqrt{2\pi}\sigma \right\}\end{aligned}$$

Probabilistic interpretation (cont'd)

Log-likelihood of the training data \mathcal{D} (assuming i.i.d)

$$\begin{aligned}\log P(\mathcal{D}) &= \log \prod_{n=1}^N p(y_n|x_n) = \sum_n \log p(y_n|x_n) \\ &= \sum_n \left\{ -\frac{[y_n - (w_0 + w_1 x_n)]^2}{2\sigma^2} - \log \sqrt{2\pi}\sigma \right\} \\ &= -\frac{1}{2\sigma^2} \sum_n [y_n - (w_0 + w_1 x_n)]^2 - \frac{N}{2} \log \sigma^2 - N \log \sqrt{2\pi}\end{aligned}$$

Probabilistic interpretation (cont'd)

Log-likelihood of the training data \mathcal{D} (assuming i.i.d)

$$\begin{aligned}\log P(\mathcal{D}) &= \log \prod_{n=1}^N p(y_n|x_n) = \sum_n \log p(y_n|x_n) \\&= \sum_n \left\{ -\frac{[y_n - (w_0 + w_1 x_n)]^2}{2\sigma^2} - \log \sqrt{2\pi}\sigma \right\} \\&= -\frac{1}{2\sigma^2} \sum_n [y_n - (w_0 + w_1 x_n)]^2 - \frac{N}{2} \log \sigma^2 - N \log \sqrt{2\pi} \\&= -\frac{1}{2} \left\{ \frac{1}{\sigma^2} \sum_n [y_n - (w_0 + w_1 x_n)]^2 + N \log \sigma^2 \right\} + \text{const}\end{aligned}$$

i.i.d stands for independently and identically distributed.

Maximum likelihood estimation

Estimating σ , w_0 and w_1 are decoupled

- Maximize over w_0 and w_1

$$\max \log P(\mathcal{D}) \Leftrightarrow \min \sum_n [y_n - (w_0 + w_1 x_n)]^2 \leftarrow \text{That is RSS}(\tilde{w})!$$

Maximum likelihood estimation

Estimating σ , w_0 and w_1 are decoupled

- Maximize over w_0 and w_1

$$\max \log P(\mathcal{D}) \Leftrightarrow \min \sum_n [y_n - (w_0 + w_1 x_n)]^2 \leftarrow \text{That is RSS}(\tilde{\mathbf{w}})!$$

- Maximize over $s = \sigma^2$ (we could estimate σ directly)

$$\frac{\partial \log P(\mathcal{D})}{\partial s} = -\frac{1}{2} \left\{ -\frac{1}{s^2} \sum_n [y_n - (w_0 + w_1 x_n)]^2 + \mathbf{N} \frac{1}{s} \right\} = 0$$

Maximum likelihood estimation

Estimating σ , w_0 and w_1 are decoupled

- Maximize over w_0 and w_1

$$\max \log P(\mathcal{D}) \Leftrightarrow \min \sum_n [y_n - (w_0 + w_1 x_n)]^2 \leftarrow \text{That is RSS}(\tilde{\mathbf{w}})!$$

- Maximize over $s = \sigma^2$ (we could estimate σ directly)

$$\begin{aligned} \frac{\partial \log P(\mathcal{D})}{\partial s} &= -\frac{1}{2} \left\{ -\frac{1}{s^2} \sum_n [y_n - (w_0 + w_1 x_n)]^2 + \mathbf{N} \frac{1}{s} \right\} = 0 \\ \rightarrow \sigma^2 = s &= \frac{1}{\mathbf{N}} \sum_n [y_n - (w_0 + w_1 x_n)]^2 \end{aligned}$$

Interpretation

Least mean square (LMS) solution (minimizing residual sum of errors)

$$\begin{pmatrix} w_0^{LMS} \\ w_1^{LMS} \end{pmatrix} = \begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix}^{-1} \begin{pmatrix} \sum_n y_n \\ \sum_n x_n y_n \end{pmatrix}$$

Interpretation

Least mean square (LMS) solution (minimizing residual sum of errors)

$$\begin{pmatrix} w_0^{LMS} \\ w_1^{LMS} \end{pmatrix} = \begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix}^{-1} \begin{pmatrix} \sum_n y_n \\ \sum_n x_n y_n \end{pmatrix}$$

Additionally

$$\sigma^2 = \frac{1}{N} \sum_n [y_n - (w_0^{LMS} + w_1^{LMS} x_n)]^2$$

Interpretation

Least mean square (LMS) solution (minimizing residual sum of errors)

$$\begin{pmatrix} w_0^{LMS} \\ w_1^{LMS} \end{pmatrix} = \begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix}^{-1} \begin{pmatrix} \sum_n y_n \\ \sum_n x_n y_n \end{pmatrix}$$

Additionally

$$\sigma^2 = \frac{1}{N} \sum_n [y_n - (w_0^{LMS} + w_1^{LMS} x_n)]^2$$

Remarks

- LMS is the same as the maximum likelihood estimation when the noise is assumed to be *Gaussian*.
- The remaining residuals provide a maximum likelihood estimate of the noise's *variance*.

NB. We sometimes call it least square solutions too.

Solution when x is one-dimensional

Least mean square (LMS) solution (minimizing residual sum of errors)

$$\begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix} \begin{pmatrix} w_0 \\ w_1 \end{pmatrix} = \begin{pmatrix} \sum_n y_n \\ \sum_n x_n y_n \end{pmatrix}$$
$$\rightarrow \begin{pmatrix} w_0^{LMS} \\ w_1^{LMS} \end{pmatrix} = \begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix}^{-1} \begin{pmatrix} \sum_n y_n \\ \sum_n x_n y_n \end{pmatrix}$$

NB. We sometimes call it least square solutions too.

LMS when \mathbf{x} is D-dimensional

$RSS(\tilde{\mathbf{w}})$ in matrix form

$$RSS(\tilde{\mathbf{w}}) = \sum_n [y_n - (w_0 + \sum_d w_d x_{nd})]^2 = \sum_n [y_n - \tilde{\mathbf{w}}^T \tilde{\mathbf{x}}_n]^2$$

where we have redefined some variables (by augmenting)

$$\tilde{\mathbf{x}} \leftarrow [1 \ x_1 \ x_2 \ \dots \ x_D]^T, \quad \tilde{\mathbf{w}} \leftarrow [w_0 \ w_1 \ w_2 \ \dots \ w_D]^T$$

LMS when x is D-dimensional

$RSS(\tilde{w})$ in matrix form

$$RSS(\tilde{w}) = \sum_n [y_n - (w_0 + \sum_d w_d x_{nd})]^2 = \sum_n [y_n - \tilde{w}^T \tilde{x}_n]^2$$

where we have redefined some variables (by augmenting)

$$\tilde{x} \leftarrow [1 \ x_1 \ x_2 \ \dots \ x_D]^T, \quad \tilde{w} \leftarrow [w_0 \ w_1 \ w_2 \ \dots \ w_D]^T$$

which leads to

$$RSS(\tilde{w}) = \sum_n (y_n - \tilde{w}^T \tilde{x}_n)(y_n - \tilde{x}_n^T \tilde{w})$$

LMS when x is D-dimensional

$RSS(\tilde{w})$ in matrix form

$$RSS(\tilde{w}) = \sum_n [y_n - (w_0 + \sum_d w_d x_{nd})]^2 = \sum_n [y_n - \tilde{w}^T \tilde{x}_n]^2$$

where we have redefined some variables (by augmenting)

$$\tilde{x} \leftarrow [1 \ x_1 \ x_2 \ \dots \ x_D]^T, \quad \tilde{w} \leftarrow [w_0 \ w_1 \ w_2 \ \dots \ w_D]^T$$

which leads to

$$\begin{aligned} RSS(\tilde{w}) &= \sum_n (y_n - \tilde{w}^T \tilde{x}_n)(y_n - \tilde{x}_n^T \tilde{w}) \\ &= \sum_n \tilde{w}^T \tilde{x}_n \tilde{x}_n^T \tilde{w} - 2y_n \tilde{x}_n^T \tilde{w} + \text{const.} \end{aligned}$$

LMS when x is D-dimensional

$RSS(\tilde{w})$ in matrix form

$$RSS(\tilde{w}) = \sum_n [y_n - (w_0 + \sum_d w_d x_{nd})]^2 = \sum_n [y_n - \tilde{w}^T \tilde{x}_n]^2$$

where we have redefined some variables (by augmenting)

$$\tilde{x} \leftarrow [1 \ x_1 \ x_2 \ \dots \ x_D]^T, \quad \tilde{w} \leftarrow [w_0 \ w_1 \ w_2 \ \dots \ w_D]^T$$

which leads to

$$\begin{aligned} RSS(\tilde{w}) &= \sum_n (y_n - \tilde{w}^T \tilde{x}_n)(y_n - \tilde{x}_n^T \tilde{w}) \\ &= \sum_n \tilde{w}^T \tilde{x}_n \tilde{x}_n^T \tilde{w} - 2y_n \tilde{x}_n^T \tilde{w} + \text{const.} \\ &= \left\{ \tilde{w}^T \left(\sum_n \tilde{x}_n \tilde{x}_n^T \right) \tilde{w} - 2 \left(\sum_n y_n \tilde{x}_n^T \right) \tilde{w} \right\} + \text{const.} \end{aligned}$$

$RSS(\tilde{\mathbf{w}})$ in new notations

Design matrix and target vector

$$\mathbf{X} = \begin{pmatrix} \mathbf{x}_1^T \\ \mathbf{x}_2^T \\ \vdots \\ \mathbf{x}_N^T \end{pmatrix} \in \mathbb{R}^{N \times D}, \quad \tilde{\mathbf{X}} = (\mathbf{1} \quad \mathbf{X}) \in \mathbb{R}^{N \times (D+1)}, \quad \mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{pmatrix}$$

$RSS(\tilde{\mathbf{w}})$ in new notations

Design matrix and target vector

$$\mathbf{X} = \begin{pmatrix} \mathbf{x}_1^T \\ \mathbf{x}_2^T \\ \vdots \\ \mathbf{x}_N^T \end{pmatrix} \in \mathbb{R}^{N \times D}, \quad \tilde{\mathbf{X}} = (\mathbf{1} \quad \mathbf{X}) \in \mathbb{R}^{N \times (D+1)}, \quad \mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{pmatrix}$$

Compact expression

$$RSS(\tilde{\mathbf{w}}) = \left\{ \tilde{\mathbf{w}}^T \tilde{\mathbf{X}}^T \tilde{\mathbf{X}} \tilde{\mathbf{w}} - 2 \left(\tilde{\mathbf{X}}^T \mathbf{y} \right)^T \tilde{\mathbf{w}} \right\} + \text{const}$$

Solution in matrix form

Normal equation

Take derivative with respect to $\tilde{\mathbf{w}}$

$$\frac{\partial RSS(\tilde{\mathbf{w}})}{\partial \tilde{\mathbf{w}}} \propto \tilde{\mathbf{X}}^T \tilde{\mathbf{X}} \mathbf{w} - \tilde{\mathbf{X}}^T \mathbf{y} = 0$$

This leads to the least-mean-square (LMS) solution

$$\tilde{\mathbf{w}}^{LMS} = \left(\tilde{\mathbf{X}}^T \tilde{\mathbf{X}} \right)^{-1} \tilde{\mathbf{X}}^T \mathbf{y}$$

Solution in matrix form

Normal equation

Take derivative with respect to $\tilde{\mathbf{w}}$

$$\frac{\partial RSS(\tilde{\mathbf{w}})}{\partial \tilde{\mathbf{w}}} \propto \tilde{\mathbf{X}}^T \tilde{\mathbf{X}} \mathbf{w} - \tilde{\mathbf{X}}^T \mathbf{y} = 0$$

This leads to the least-mean-square (LMS) solution

$$\tilde{\mathbf{w}}^{LMS} = \left(\tilde{\mathbf{X}}^T \tilde{\mathbf{X}} \right)^{-1} \tilde{\mathbf{X}}^T \mathbf{y}$$

Verify the solution when $D = 1$

$$\tilde{\mathbf{X}}^T \tilde{\mathbf{X}} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_1 & x_2 & \cdots & x_N \end{pmatrix} \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \cdots & \cdots \\ 1 & x_N \end{pmatrix} = \begin{pmatrix} \sum_n 1 & \sum_n x_n \\ \sum_n x_n & \sum_n x_n^2 \end{pmatrix}$$

Mini-Summary

- Linear regression is the linear combination of features.
 $f : \mathbf{x} \rightarrow y$, with $f(\mathbf{x}) = w_0 + \sum_d w_d x_d = w_0 + \mathbf{w}^T \mathbf{x}$
- If we minimize residual sum squares as our learning objective, we get a closed-form solution of parameters.
- Probabilistic interpretation: maximum likelihood if assuming residual is Gaussian distributed
- Other interpretations exist: if interested, please consult the slides from last year's lectures.

Outline

1 Linear regression

2 Perceptron

- Intuition
- Algorithm

Main idea

Consider a linear model for binary classification

$$\mathbf{w}^T \mathbf{x}$$

is used to distinguish two classes $\{-1, +1\}$.

Our goal

$$\varepsilon = \sum_n \mathbb{I}[y_n \neq \text{sign}(\mathbf{w}^T \mathbf{x}_n)]$$

i.e., at least the errors on the training dataset are reduced.

Hard, but easy if we have only one training example

How to change \mathbf{w} such that

$$y_n = \text{sign}(\mathbf{w}^T \mathbf{x}_n)$$

Two cases

- If $y_n = \text{sign}(\mathbf{w}^T \mathbf{x}_n)$, do nothing.
- If $y_n \neq \text{sign}(\mathbf{w}^T \mathbf{x}_n)$,

$$\mathbf{w}^{\text{NEW}} \leftarrow \mathbf{w}^{\text{OLD}} + y_n \mathbf{x}_n$$

Why would it work?

If $y_n \neq \text{sign}(\mathbf{w}^T \mathbf{x}_n)$, then

$$y_n(\mathbf{w}^T \mathbf{x}_n) < 0$$

Why would it work?

If $y_n \neq \text{sign}(\mathbf{w}^T \mathbf{x}_n)$, then

$$y_n(\mathbf{w}^T \mathbf{x}_n) < 0$$

What would happen if we change to new $\mathbf{w}^{\text{NEW}} = \mathbf{w} + y_n \mathbf{x}_n$?

$$y_n[(\mathbf{w} + y_n \mathbf{x}_n)^T \mathbf{x}_n] = y_n \mathbf{w}^T \mathbf{x}_n + y_n^2 \mathbf{x}_n^T \mathbf{x}_n$$

We are adding a positive number, so it would be possible for the new \mathbf{w}^{NEW}

$$y_n(\mathbf{w}^{\text{NEW}T} \mathbf{x}_n) > 0$$

i.e., classify correctly!

Perceptron

Iteratively solving one case at a time

- REPEAT
- Pick a data point \mathbf{x}_n (can be a fixed order of the training instances)
- Make a prediction $y = \text{sign}(\mathbf{w}^T \mathbf{x}_n)$ using the *current* \mathbf{w}
- If $y = y_n$, do nothing. Else,

$$\mathbf{w} \leftarrow \mathbf{w} + y_n \mathbf{x}_n$$

- UNTIL converged.

Properties

- If the training data is linearly separable, the algorithm stops in a finite number of steps.
- The parameter vector is always a linear combination of training instances.