

# Using NDVI to improve LiDAR-based Canopy Height Models

Hugo Hunt, Eden Project University Centre



## MSc Dissertation References

Acebes, P.; Lillo, P.; Jaime-González, C. (2021) Disentangling LiDAR Contribution in Modelling Species–Habitat Structure Relationships in Terrestrial Ecosystems Worldwide. A Systematic Review and Future Directions. *Remote Sens.* 2021, 13, 3447. <https://doi.org/10.3390/rs13173447>

Aerial Photography Great Britain (APGB) (2024). APGB 50cm Colour Infrared (CIR) imagery. Tiles: ny30 clipped. Downloaded: 2024-04-12

Allin, Z (2024). Conversation with Hugo Hunt, 21 August.

Bakx, T.R., Koma, Z., Seijmonsbergen, A.C. and Kissling, W.D., 2019. Use and categorization of light detection and ranging vegetation metrics in avian diversity and species distribution research. *Diversity and Distributions*, 25(7), pp.1045-1059.

Blackburn, G.A., Abd Latif, Z. and Boyd, D.S., 2014. Forest disturbance and regeneration: a mosaic of discrete gap dynamics and open matrix regimes?. *Journal of Vegetation Science*, 25(6), pp.1341-1354.

Brownnett, J. (2024). Email to Hugo Hunt, 24 January. Natural England

Carter, J. V., Pan, J., Rai, S. N., & Galandiuk, S. (2016). ROC-ing along: Evaluation and interpretation of receiver operating characteristic curves. *Surgery*, 159(6), 1638-1645.

Chávez, R.O., Clevers, J.G.P.W., Decuyper, M., De Bruin, S. and Herold, M., 2016. 50 years of water extraction in the Pampa del Tamarugal basin: Can Prosopis tamarugo trees survive in the hyper-arid Atacama Desert (Northern Chile)?. *Journal of Arid Environments*, 124, pp.292-303.

Choudhury, B. J., & Tucker, C. J. (1987). Satellite observed seasonal and inter-annual variation of vegetation over the Kalahari, the Great Victoria Desert, and the Great Sandy Desert: 1979–1984. *Remote Sensing of Environment*, 23(2), 233-241.

De'ath, G., & Fabricius, K. E. (2000). Classification and regression trees: a powerful yet simple technique for ecological data analysis. *Ecology*, 81(11), 3178-3192.

DEFRA (2023), “Environmental Improvement Plan 2023”. First revision of the 25 Year Environment Plan. *Environmental Improvement Plan 2023*, DEFRA, 31 Jan. 2023. Available at: [assets.publishing.service.gov.uk/media/64a6d9c1c531eb000c64fffa/environmental-improvement-plan-2023.pdf](https://assets.publishing.service.gov.uk/media/64a6d9c1c531eb000c64fffa/environmental-improvement-plan-2023.pdf). (Accessed: 17 August 2024).

DEFRA (2024), *National Lidar Programme* - [data.gov.uk](https://www.data.gov.uk/dataset/f0db0249-f17b-4036-9e65-309148c97ce4/national-lidar-programme). Available at: <https://www.data.gov.uk/dataset/f0db0249-f17b-4036-9e65-309148c97ce4/national-lidar-programme> (Accessed: 26 July 2024).

Environmental Systems Research Institute (Esri). (2022). ArcGIS Pro 3.0.1. Redlands, CA: Environmental Systems Research Institute (Esri)

Frampton, W. J., Dash, J., Watmough, G., & Milton, E. J. (2013). Evaluating the capabilities of Sentinel-2 for quantitative estimation of biophysical variables in vegetation. *ISPRS journal of photogrammetry and remote sensing*, 82, 83-92.

Garzon, M. B., Blazek, R., Neteler, M., De Dios, R. S., Ollero, H. S., & Furlanello, C. (2006). Predicting habitat suitability with machine learning models: the potential area of *Pinus sylvestris* L. in the Iberian Peninsula. *Ecological modelling*, 197(3-4), 383-393.

Gaulton, R., Olayaa, G., Wallington, E. D., Malthus, T. J., Woodhouse, I. H., & Suarez, J. C. (2005). CONTINUOUS COVER FORESTRY IN THE UK? QUANTIFYING FOREST STRUCTURE USING REMOTE SENSING.

Gitelson, A. A., Kaufman, Y. J., & Merzlyak, M. N. (1996). Use of a green channel in remote sensing of global vegetation from EOS-MODIS. *Remote sensing of Environment*, 58(3), 289-298.

Gwenzi, D. and Lefsky, M.A. (2014) 'Modeling canopy height in a savanna ecosystem using spaceborne lidar waveforms', *Remote Sensing of Environment*, 154, pp. 338–344. doi:10.1016/j.rse.2013.11.024.

Hamer, A. (2024). Conversation with Hugo Hunt, 29 August.

Hamer, A., Dixon, S., Kratz, C., Dornan, C., Miller, C., Prince, M., Hart, C., Hunt, T., and Webb, A. (2023). *England Peat Map: The challenges of using Earth observation data and machine learning approaches at the national scale*. EGU General Assembly 2023, Vienna, Austria, 24–28 Apr, EGU23-3514, <https://doi.org/10.5194/egusphere-egu23-3514>, 2023.

Hanley, J. A. (1989). Receiver operating characteristic (ROC) methodology: the state of the art. *Crit Rev Diagn Imaging*, 29(3), 307-335.

Hijmans R (2024). *\_terra: Spatial Data Analysis\_*. R package version 1.7-71, <<https://CRAN.R-project.org/package=terra>>

Huang, S., Tang, L., Hupy, J.P., Wang, Y. and Shao, G., 2021. A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *Journal of Forestry Research*, 32(1), pp.1-6.

Hyde, P., Dubayah, R., Walker, W., Blair, J.B., Hofton, M. and Hunsaker, C., 2006. Mapping forest structure for wildlife habitat analysis using multi-sensor (LiDAR, SAR/InSAR, ETM+, Quickbird) synergy. *Remote Sensing of Environment*, 102(1-2), pp.63-73.

Jiang, Z., Huete, A. R., Didan, K., & Miura, T. (2008). Development of a two-band

enhanced vegetation index without a blue band. *Remote sensing of Environment*, 112(10), 3833-3845.

Justice, C. O., Townshend, J. R. G., Holben, B. N., & Tucker, E. C. (1985). Analysis of the phenology of global vegetation using meteorological satellite data. *International Journal of Remote Sensing*, 6(8), 1271-1318.

Kampichler, C., Wieland, R., Calmé, S., Weissenberger, H., & Arriaga-Weiss, S. (2010). Classification in conservation biology: a comparison of five machine-learning methods. *Ecological Informatics*, 5(6), 441-450.

Kane, V.R., Bakker, J.D., McGaughey, R.J., Lutz, J.A., Gersonde, R.F. and Franklin, J.F., 2010. Examining conifer canopy structural complexity across forest ages and elevations with LiDAR data. *Canadian Journal of Forest Research*, 40(4), pp.774-787.

Karnieli, A., Agam, N., Pinker, R. T., Anderson, M., Imhoff, M. L., Gutman, G. G., ... & Goldberg, A. (2010). Use of NDVI and land surface temperature for drought assessment: Merits and limitations. *Journal of climate*, 23(3), 618-633.

Kashani, A. G., Olsen, M. J., Parrish, C. E., & Wilson, N. (2015). A review of LiDAR radiometric processing: From ad hoc intensity correction to rigorous radiometric calibration. *Sensors*, 15(11), 28099-28128.

Khosravipour, A., Skidmore, A.K., Wang, T., Isenburg, M. and Khoshelham, K., 2015. Effect of slope on treetop detection using a LiDAR Canopy Height Model. *ISPRS journal of photogrammetry and remote sensing*, 104, pp.44-52.

Lee, W.J. and Lee, C.W., 2018. Forest canopy height estimation using multiplatform remote sensing dataset. *Journal of Sensors*, 2018(1), p.1593129.

Lefsky, M.A., Cohen, W.B., Parker, G.G. and Harding, D.J., 2002. Lidar remote sensing for ecosystem studies: Lidar, an emerging remote sensing technology that directly measures the three-dimensional distribution of plant canopies, can accurately estimate vegetation structural attributes and should be of particular interest to forest, landscape, and global ecologists. *BioScience*, 52(1), pp.19-30.

Lidar Composite Digital Terrain Model England 1m resolution [TIFF geospatial data], Scale 1:5000, Tiles: ny30ne, Updated: 5 May 2022, Open Government Licence, Using: EDINA LIDAR Digimap Service, <<https://digimap.edina.ac.uk>>, Downloaded: 2024-04-12 11:21:01.692

Lidar Composite First Return Digital Surface Model England 1m resolution [TIFF geospatial data], Scale 1:5000, Tiles: ny30ne, Updated: 5 May 2022, Open Government Licence, Using: EDINA LIDAR Digimap Service, <<https://digimap.edina.ac.uk>>, Downloaded: 2024-04-12 11:21:01.692

Luscombe, D. J., Gatis, N., Anderson, K., Carless, D., & Brazier, R. E. (2023). Rapid, repeatable landscape-scale mapping of tree, hedgerow, and woodland habitats (THaW), using airborne LiDAR and spaceborne SAR data. *Ecology and Evolution*, 13(5), e10103.

Mangewa, L. J., Ndakidemi, P. A., Alward, R. D., Kija, H. K., Bukombe, J. K., Nasolwa,

E. R., & Munishi, L. K. (2022). Comparative assessment of UAV and sentinel-2 NDVI and GNDVI for preliminary diagnosis of habitat conditions in Burunge wildlife management area, Tanzania. *Earth*, 3(3), 769-787.

Manning, A. D., Kesteven, J., Stein, J., Lunn, A., Xu, T., & Rayner, B. (2010). Could native Scots pines (*Pinus sylvestris*) still persist in northern England and southern Scotland?. *Plant Ecology & Diversity*, 3(2), 187-201.

Matsushita, B., Yang, W., Chen, J., Onda, Y., & Qiu, G. (2007). Sensitivity of the enhanced vegetation index (EVI) and normalized difference vegetation index (NDVI) to topographic effects: a case study in high-density cypress forest. *Sensors*, 7(11), 2636-2651.

Meneses-Tovar, C. L. (2011). NDVI as indicator of degradation. *Unasylva*, 62(238), 39-46.

Müller, J., Brandl, R., Brändle, M., Förster, B., de Araujo, B.C., Gossner, M.M., Ladas, A., Wagner, M., Maraun, M., Schall, P. and Schmidt, S. (2018). LiDAR-derived canopy structure supports the more-individuals hypothesis for arthropod diversity in temperate forests. *Oikos*, 127(6), pp.814-824.

Murdoch D, Adler D (2024). `_rgl: 3D Visualization Using OpenGL_`. R package version 1.3.1, <<https://CRAN.R-project.org/package=rgl>>.

Ordnance Survey (2024), 1:25 000 Scale Colour Raster [TIFF geospatial data], Scale 1:25000, Tiles: ny30\_clipped, Updated: 15 February 2024, Using: EDINA Digimap Ordnance Survey Service, <<https://digimap.edina.ac.uk>>, Downloaded: 2024-04-02 10:57:53.514

Peñuelas, J., Filella, I., Biel, C., Serrano, L., & Save, R. (1993). The reflectance at the 950–970 nm region as an indicator of plant water status. *International journal of remote sensing*, 14(10), 1887-1905.

Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J. M., Tucker, C. J., & Stenseth, N. C. (2005). Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in ecology & evolution*, 20(9), 503-510.

Powell, C. (2024). Conversation with Hugo Hunt, 21 August. Centre for Information Resilience.

QGIS.org (2024). QGIS Geographic Information System. Open Source Geospatial Foundation Project

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Roca, D., Martínez-Sánchez, J., Lagüela, S., & Arias, P. (2016). Novel aerial 3D mapping system based on UAV platforms and 2D laser scanners. *Journal of Sensors*, 2016(1), 4158370.

Roussel, J.R., and Auty, D. (2024). Airborne LiDAR Data Manipulation and Visualization for Forestry Applications. R package version 4.1.2. <https://cran.r->

[project.org/package=lidR](https://project.org/package=lidR)

Roussel, J.R., Auty, D., Coops, N. C., Tompalski, P., Goodbody, T. R. H., Sánchez Meador, A., Bourdon, J.F., De Boissieu, F., Achim, A. (2020). lidR : An R package for analysis of Airborne Laser Scanning (ALS) data. *Remote Sensing of Environment*, 251 (August), 112061. <[doi:10.1016/j.rse.2020.112061](https://doi.org/10.1016/j.rse.2020.112061)>.

Schultz, A., Wieland, R., & Lutze, G. (2000). Neural networks in agroecological modelling—stylish application or helpful tool?. *Computers and Electronics in Agriculture*, 29(1-2), 73-97.

Sibona, E., Vitali, A., Meloni, F., Caffo, L., Dotta, A., Lingua, E., Motta, R. and Garbarino, M. (2016). Direct measurement of tree height provides different results on the assessment of LiDAR accuracy. *Forests*, 8(1), p.7.

Solano, F., Modica, G., Pratico, S., Box, O.F. and Piovesan, G., 2022. Unveiling the complex canopy spatial structure of a Mediterranean old-growth beech (*Fagus sylvatica* L.) forest from UAV observations. *Ecological Indicators*, 138, p.108807.

Steele, B. M. (2000). Combining multiple classifiers: An application using spatial and remotely sensed information for land cover type mapping. *Remote sensing of environment*, 74(3), 545-556.

Sun, J., Wang, P., Gao, Z., Liu, Z., Li, Y., Gan, X., & Liu, Z. (2021). Wood–leaf classification of tree point cloud based on intensity and geometric information. *Remote Sensing*, 13(20), 4050.

Suzuki, R., Kobayashi, H., Delbart, N., Asanuma, J. and Hiyama, T., 2011. NDVI responses to the forest canopy and floor from spring to summer observed by airborne spectrometer in eastern Siberia. *Remote sensing of environment*, 115(12), pp.3615-3624.

Törmä, M., Lumme, J., Pyysalo, U., Patrikainen, N. and Luojus, K. (2004). Tree species classification using ers sar and modis ndvi images. *International Archives of Photogrammetry Remote Sensing and Spatial Information Sciences*, 35(7), pp.927-932.

United Nations. (2019). *Resolution adopted by the General Assembly on 1 March 2019: United Nations Decade on Ecosystem Restoration (2021–2030)* (A/RES/73/284).

<https://documents.un.org/doc/undoc/gen/n19/060/16/pdf/n1906016.pdf>

Vallet, P., Dhôte, J.F., Le Moguédec, G., Ravart, M. and Pignard, G. (2006). Development of total aboveground volume equations for seven important forest tree species in France. *Forest Ecology and Management*, 229(1-3), pp.98-110.

Wang, J., Rich, P.M., Price, K.P. and Kettle, W.D., 2004. Relations between NDVI and tree productivity in the central Great Plains. *International Journal of Remote Sensing*, 25(16), pp.3127-3138.

Wasser, L.H. (2024) *What is a CHM, DSM and DTM? about gridded, raster lidar data / Nsf neon | open data to understand our ecosystems*, National Ecological Observatory

*Network (NEON) Learning Hub.* Available at:  
<https://www.neonscience.org/resources/learning-hub/tutorials/chm-dsm-dtm>  
(Accessed: 20 July 2024).

Wickham, H., Henry, L. (2023). *\_purrr: Functional Programming Tools\_*. R package version 1.0.2, <<https://CRAN.R-project.org/package=purrr>>.

Woodwell, G. M., Houghton, R. A., Stone, T. A., & Park, A. B. (1986). Changes in the area of forests in Rondônia, Amazon Basin, measured by satellite imagery. In *The changing carbon cycle: a global analysis* (pp. 242-257). New York, NY: Springer New York.

