

## Analysis of DNA methylation in invertebrates requires consideration of genome characteristics and methylation landscape

Shelly A. Trigg<sup>\*1</sup>, Yaamini R. Venkataraman<sup>\*1</sup>, Mackenzie R. Gavery<sup>2</sup>, Steven B. Roberts<sup>1</sup>, Debashish Bhattacharya<sup>3</sup>, Alan Downey-Wall<sup>4</sup>, Jose M. Eirin-Lopez<sup>5</sup>, Kevin M. Johnson<sup>6</sup>, Katie E/ Lotterhos<sup>4</sup>, Jonathan B. Puritz<sup>7</sup> and Hollie M/ Putnam<sup>7\*</sup>

<sup>1</sup> University of Washington, School of Aquatic and Fishery Sciences 1122 NE Boat St. Seattle, WA, 98195, USA

<sup>2</sup> NOAA Northwest Fisheries Science Center Montlake 2725 Montlake Blvd E, Seattle, WA, 98112, USA

<sup>3</sup> Department of Biochemistry and Microbiology, Rutgers University, New Brunswick, NJ 08901 USA

<sup>4</sup> Department of Marine and Environmental Sciences, Northeastern University, 430 Nahant Road, Nahant, MA 01908

<sup>5</sup> Florida International University, Environmental Epigenetics Laboratory, Institute of Environment 3000 NE 151 St. North Miami, FL, 33181, USA

<sup>6</sup>Department of Biological Sciences, University of Rhode Island, Kingston, RI 02881, USA

<sup>+</sup>corresponding author: [hputnam@uri.edu](mailto:hputnam@uri.edu)

<sup>\*</sup> equal contribution

### Table of Contents:

**Figure SF1:** Summary of sequencing depth and alignments for all libraries. (page 3)

**Figure SF2:** Bisulfite conversion efficiency. (page 4)

**Figure SF3:** Methylation status of CpGs covered by each method (page 5)

**Figure SF4:** PCA of CpG methylation for loci covered at 5x read depth by all *M. capitata* and *P. acuta* samples. (page 6)

**Figure SF5:** Estimated fraction of genome-wide CpG loci covered by at least 5 reads with different sequencing depths for each species. (page 7)

**Figure SF6:** Coverage of orthologs across methods. (page 8)

**Table ST1:** [Trimming statistics](#)

**Table ST2:** [Alignment statistics](#)


**Table ST3:** [C1 alignment statistics](#)

**Table ST4:** [Lambda conversion efficiency and alignment statistics](#)


**Table ST5:** [Bisulfite conversion efficiency comparisons](#)

**Table ST6:** [Results from PERMANOVA and beta-dispersion tests for genomic location.](#)


**Table ST7:** [Contingency test results for \*M. capitata\* and \*P. acuta\*](#)



**Figure SF1.** Summary of sequencing depth and alignments for all libraries. Bars show average number of reads for each method and species and error bars show standard deviation.




**Figure SF2.** Bisulfite conversion efficiency assessment. Bisulfite conversion efficiency is different across library preparation methods whether calculated from lambda alignments or estimated from non-CpG methylation from coral alignments for *M. capitata* libraries and *P. acuta* libraries. Generally, bisulfite conversion efficiency calculation methods are not different.




**Figure SF3.** Percent of highly methylated ( $\geq 50\%$ ; darkest shade), moderately methylated ( $10\text{--}50\%$ ; medium shade), and lowly methylated CpGs ( $< 10\%$ ; lightest shade) detected by each method **A)** for *M. capitata* and **B)** *P. acuta*, based on the number of CpGs captured by each method separately. Principal Coordinate Analyses associated with perMANOVA and beta-dispersion tests related to **Table ST6** that show


differences in proportion of CpGs that are highly ( $\geq 50\%$ ), moderately (10-50%), or lowly ( $\leq 10\%$ ) methylated in **C)** *M. capitata* and **D)** *P. acuta*. WGBS is represented by green circles, RRBS by purple triangles, and MBDBS by orange diamonds. Percent variation explained by each PCoA axis is included in the axis label. Ellipses depict 95% confidence intervals for each sequencing method. All eigenvectors are significant at the  $\alpha = 0.05$  level.



**Figure SF4.** PCA of CpG methylation for loci covered at 5x read depth in all samples for **(A)** *M. capitata* and **(B)** *P. acuta*.



**Figure SF5.** Estimated fraction of CpG sites in the genome covered by at least 5 reads at different sequencing depths (number of M read pairs) for **(A)** *M. capitata* and **(B)** *P. acuta* samples for each bisulfite sequencing method. ‘Observed’ (opaque line and dots) denotes the fraction of genome-wide CpG loci covered by at least 5 reads determined from pooled data that was subsampled at 50M, 100M, 150M, and 200M reads. ‘Estimated’ (translucent line and dots) denotes the fraction of genome-wide CpG loci covered by at least 5 reads estimated by michaelis-menten modelling of the ‘observed’ data with standard error shown by shaded areas. All samples within a bisulfite sequencing method were pooled for the downsampling analyses.



**Figure SF6.** Mean proportion (n=3 samples per method) of CpGs per gene that have at least 5x coverage in all of the one-to-one orthologous genes, as identified by OrthoFinder (Putnam et al., 2020) for **A**) *M. capitata* and **B**) *P. acuta*.