

	perm	ncal	mobj	plant	lrn	comp	inp	int	spc
validity tests reported by the row's records, as k/n over the records whose design can carry the class (control task: none reported, applicability unresolved)									
F1a clinical findings and anatomy (linear)	1/18	1/18	0/18	1/18	1-2/18	–	–	–	–
F1a' the same, non-linear readouts	0/15	0/15	0/15	0/15	0/15	–	–	–	–
F1b held-out concept recovery, text scorers	–	0/6	0/6	1/6	0/6	–	–	–	–
F1b' concept-vector gradient association	0/2	0/2	0/2	0/2	0/2	–	–	–	–
F1b'' labelled representational correspondence	–	0-1/5	0/5	0-1/5	0/5	–	–	–	–
F1c concept layers on foundation features	0/13	0/13	0/13	0/13	1/13	–	–	1/13	–
F2 sex, age or race decodable	2/9	0/9	0/9	1/9	1/9	–	–	–	–
F2b acquisition site and device decodable	1/12	0/12	1/12	0/12	0/12	–	–	–	–
F3a centre and scanner structure, pathology	0/9	0/18	0/10	0/10	0/18	2-3/8	–	–	–
F3b post-hoc harmonisation raises invariance	–	–	–	–	–	–	–	–	–
F3c site and artefact structure in brain MRI	0/1	0/3	1/1	0/1	0/3	0-1/2	–	–	–
F4 sex redundant for a refitted probe	–	–	–	–	–	–	–	–	–
F5 anisotropy and the effect of whitening	–	0/1	–	–	0/1	1/1	–	–	–
F5b medical encoders share structure	–	1/5	–	–	0/5	4/5	–	–	–
F6 modality gap; separation task-dependent	–	0/1	–	–	0/1	1/1	–	–	–
F8a attribution sets fail an instrumented test	–	–	–	–	0-1/6	–	5-6/6	0/6	–
F8a' attribution methods disagree	–	–	–	–	0-1/3	–	1-2/3	0/3	–
F8a'' maps implausible on inspection	–	–	–	–	0/2	–	0/2	0/2	–
F8b labels less decodable after the connector	–	–	–	–	0/1	–	0/1	0/1	–
F9a targeted interventions report changed outputs	–	–	–	–	–	–	–	–	2-3/8
F7 reported validation of dictionary features	row selected on semantic validation; no share shown								
F9b inventory: editing and unlearning	–	–	–	–	–	–	–	–	0/1
F10 inventory: medical vs general decodability	0/12	0-1/20	0/20	0/20	1-2/20	–	–	0/3	–
F11 inventory: properties shared or differing	2/5	2/19	0/8	2-3/8	0/21	11/11	2/2	0/2	–