

# Probability & Statistics Project

Hana Ali Rashid, hr05940  
Tasmiya Malik, Student ID  
Ifrah Ilyas, Student ID

April 27, 2021

## Q1: Random Walk



### 1.1

Function implementation in Python:

```
1 def get_updated_position(n,p):
2     pos = 0 #position
3     for _ in range(n):
4         rand = random.randint(1,100) #generating a random number in the range 1 to 100
5         if rand < p*100:
6             pos += 1 #move one step right
7         else:
8             pos -= 1 #move one step left
9     return pos #return final position
```


Calling the function for several iterations to get multiple expected values:

```
1 def main_11():
2     '''Calculating and plotting expected outcomes for various combinations of n and p'''
3     p = 0.7
4     n = 10
5     expected = []
6     for j in range(50): #expected values for each (n,p) for 50 iterations
7         outcomes = []
8         for i in range(25):
9             outcomes.append(get_updated_position(n,p))
10            expected.append(sum(outcomes)/25) #appending the expected (average) value for each(n,p)
11        #plotting and showing a histogram of calculated expected values
12        fig, ax = plt.subplots(figsize=(10, 7))
13        ax.hist(expected, bins = range(-5,10))
14        plt.title('n = '+str(n)+', p = '+str(p)+'\n Mean = '+str(round(statistics.mean(expected),3))+'\n Variance = '+str(round(statistics.variance(expected),3)))
15        plt.savefig("Q1_histograms/q1"+'_n = '+str(n)+'_ p = '+str(p)+'.png')
16        plt.show()
```



(a) No. of steps taken = 10 with probability of moving a step right = 0.5.

(b) No. of steps taken = 18 with probability of moving a step right = 0.5.



(c) No. of steps taken = 10 with probability of moving a step right = 0.7.

Histograms produced by the above code for expected final positions of objects against frequency, where  $n$  is the number of steps taken and  $p$  is the probability of the object moving one step to the right.

Histograms (a) and (b) have the same value of  $p$  and varying number of steps  $n$ .

Both histograms appear to follow a normal distribution, with (a) having a mean of  $-0.235$  and a variance of  $0.35$  and (b) having a mean of  $-0.205$  and a variance of  $0.61$ . Increasing the number of steps has not had a significant effect on the mean but did increase the variance.

Histograms (a) and (c) have the same number of steps  $n$  and varying values of  $p$ .

Histogram (c) also appears to follow a normal distribution, having a mean of  $3.859$  and a variance of  $0.273$ . Increasing  $p$  appears to have affected the mean of the distribution but not the variance as such.



## 1.2

Function implementation in Python:

```
1 def get_updated_position_restricted(n,p):
2     pos = 0 #position
3     for _ in range(n):
4         rand = random.randint(1,100) #generating a random number in the range 1 to 100
5         if rand < p*100 or pos <= 0: #move one step right if pos == 0
6             pos += 1
7         else:
8             pos -= 1 #move one step left
9     return pos #return final position
```



Calling the function for several iterations to get multiple expected values:

```
1 def main_12():
2     '''Calculating and plotting expected outcomes for various combinations of n and p'''
3     p = 0.7
4     n = 24
5     expected = []
6     for j in range(50): #expected value for each (n,p) for 25 iterations
7         outcomes = []
8         for i in range(25):
9             outcomes.append(get_updated_position_restricted(n,p))
10        expected.append(sum(outcomes)/25) #appending the expected (average) value for each(n,p)
11    #plotting and showing a histogram of calculated expected values
12    fig, ax = plt.subplots(figsize =(10, 7))
13    ax.hist(expected, bins =
14    [4,4.5,5,5.5,6,6.5,7,7.5,8,8.5,9,9.5,10,10.5,11,11.5,12,12.5,13,13.5,14])
15    plt.title('n = '+str(n)+', p = '+str(p)+'\n Mean = '+str(round(statistics.mean(expected)
16    ,3))+'\n Variance = '+str(round(statistics.variance(expected),3)))
17    plt.savefig("Q1_histograms/Q1.22 "+"_n = '+str(n)+'_p = '+str(p)+'.png')
18    plt.show()
```



(d) No. of steps taken = 20 with probability of moving a step right = 0.5.

(e) No. of steps taken = 24 with probability of moving a step right = 0.5.



(f) No. of steps taken = 20 with probability of moving a step right = 0.7.

(g) No. of steps taken = 24 with probability of moving a step right = 0.7.

Histograms produced by the above code for expected final positions of objects against frequency, where  $n$  is the number of steps taken and  $p$  is the probability of the object moving one step to the right.

Histograms (d) and (e) have the same value of  $p$  and varying number of steps  $n$ . Both histograms appear to follow a normal distribution, with (d) having a mean of 3.267 and a variance of 0.233 and (e) having a mean of 3.466 and a variance of 0.367. Again, increasing the number of steps has not had a significant effect on the mean but did increase the variance.

Histograms (f) and (g) have the same number of steps  $n$  respectively as (d) and (e) and varying values of  $p$ . They also appear to follow a normal distribution, with (f) having a mean of 8.462 and a variance of 0.559 and (g) having a mean of 10.187 and a variance of 0.668. Increasing  $p$  appears to have increased the mean and variance, but the mean has increased more significantly.

Additionally, due to the added constraint in this part, we see that no expected value for the final position is negative.

## 1.3


Function implementation in Python:

```
1 def stepsToMeet(pos1,pos2,p1,p2):
2     count = 0 #keeps count of number of steps taken for objects to meet
3     while pos1 != pos2:
4         rand = random.randint(1,100) #generating a random number in the range 1 to 100 to
         determine outcome
5         if rand < p1*100:
6             pos1 += 1 #move one step right
7         else:
8             pos1 -= 1 #move one step left
9         rand = random.randint(1,100) #generating a random number in the range 1 to 100
10        if rand < p2*100:
11            pos2 += 1 #move one step right
12        else:
13            pos2 -= 1 #move one step left
14        count += 1
15    return count
```


Calling the function for several iterations to get multiple expected values:

```
1 def main_13():
2     '''Calculating and plotting expected outcomes for various combinations of n & p'''
3     expected = []
4     p1 = 0.7
5     p2 = 0.3
6     pos1 = -4
7     pos2 = 4
8     for i in range(25): #calculating the expected value for each (n,p) for 25 iterations
9         outcomes = []
10        for j in range(25):
11            outcomes.append(stepsToMeet(pos1,pos2,p1,p2))
12        #calculating the average expected value for each(n,p)
13        expected.append(sum(outcomes)/25) #appending the expected (average) value for each(n,p)
14    #plotting a histogram of calculated expected values
15    fig, ax = plt.subplots(figsize =(10, 7))
16    ax.set_xlabel('Expected no. of steps taken to meet')
17    ax.set_ylabel('Frequency')
18    ax.hist(expected, bins = range(5,18))
19    plt.title('p1 = '+str(p1)+' , p2 = '+str(p2)+'\npos1 = '+str(pos1)+' , pos2 = '+str(pos2)+'\n
n Mean = '+str(round(statistics.mean(expected),3))+' , Variance = '+str(round(statistics.
variance(expected),3))')
20    plt.savefig("Q1_histograms/Q1.3 "+'_p1 = '+str(p1)+'_p2 = '+str(p2)+'_pos1 = '+str(pos1)+'
_pos2 = '+str(pos2)+'(6).png')
21    plt.show()
```

Histograms produced by the above code for expected number of steps taken for two objects walking randomly to meet, against frequency.  $p_1$  and  $p_2$  are the respective probabilities of object 1 and object 2 moving one step to the right, and  $pos_1$  and  $pos_2$  denote the starting positions of object 1 and object 2 respectively.



The above histogram shows that the expected number of steps to meet for two objects follows a normal distribution of mean 10.09 and variance 1.94 when they begin 6 steps apart with the given probabilities.



The above histogram shows that the expected number of steps to meet for two objects follows a normal distribution of mean 9.90 and variance 1.16 when they begin 8 steps apart with the given probabilities.

## Q5: Hypothesis Testing

### 5.1


Function that implements the simulation of a fair coin:

```
1 def cointoss():
2     #assuming 0 = T & 1 = H
3     return(random.randint(0,1))
```

Function that uses the above function to simulate 10 coin tosses multiple times and finds the expected number of times the null hypothesis is rejected:

```
1 def simulate_tosses():
2     expected = []
3     for n in range(100): #to get 100 expected values
4         rejected = []
5         for j in range(100): #to get 1 expected value
6             outcomes = []
7             for i in range(10): #to get 10 outcomes/1 rejected value
8                 outcomes.append(cointoss())
9             #check if hypothesis is accepted or rejected
10            if sum(outcomes) == 0 or sum(outcomes) == 1 or sum(outcomes) == 10 or sum(outcomes) == 9:
11                rejected.append(1) #hypothesis rejected
12            else:
13                rejected.append(0) #hypothesis accepted
14            expected.append(sum(rejected)/100)
15        #plotting histogram
16        fig, ax = plt.subplots(figsize=(10, 7))
17        ax.hist(expected,width = 0.01)
18        ax.set_xlabel('Times the null hypothesis is rejected')
19        ax.set_ylabel('Frequency')
20        plt.title('Mean: '+str(round(statistics.mean(expected),3))+', Variance: '+str(round(
21        statistics.variance(expected),3)))
22        plt.savefig("Q5_histograms/Q5.1.png")
23        plt.show()
```

Histogram of expected values:



Probability we will reject the null hypothesis even though it is true:

*Simulation-wise:* According to the mean of the expected values, the probability is 2.3%.

*Mathematically:* We reject the null hypothesis if the probability of the outcomes is below the threshold. Since the probability of the outcomes being accepted is 0.95 and the probability of them being rejected is 0.05, the probability that we reject the null hypothesis even though it is true is also 0.05 which is equal to the threshold.

### 5.2.1

The following function conducts a single hypothesis test by taking a sample of 30 fish and calculating the mean and standard deviation of their lengths to check the following condition to accept or reject the null hypothesis:

$$P(|S - u_0| \geq a) < 0.05$$


The function returns 1 if the hypothesis is rejected, and 0 if it is accepted.

```
1 def hypothesis_test(u_0, n):
2     s = []
3     for i in range(n): #catching a sample of 30 fish
4         s.append(f.fish())
5     s = np.array(s)
6     std = s.std() #sample standard deviation
7     mean = s.mean() #sample mean
8     a = abs(mean - u_0)
9     prob = 2*(stats.norm(u_0, std/math.sqrt(n)).cdf(u_0 - a))
10    if prob < 0.05:
11        return 1 #rejected
12    else:
13        return 0 #accepted
```

The following function performs 50 experiments and uses the average of their outcome to calculate a single expected value. It repeats this for 100 iterations and plots a histogram of the expected values.

```
1 def experiments():
2     u_0 = 23 #population mean
3     n = 30 #sample size
4     expected = []
5     for j in range(100): #getting 100 expected values
6         outcomes = []
7         for i in range(50): #conducting 50 hypothesis tests
8             outcomes.append(hypothesis_test(u_0,n))
9         expected.append(sum(outcomes)/50) #calculating one expected value from the 50 tests
10    #plotting histogram of expected values
11    fig, ax = plt.subplots(figsize=(10, 7))
12    ax.hist(expected, width = 0.04)
13    ax.set_xlabel('Times the null hypothesis is rejected')
14    ax.set_ylabel('Frequency')
15    plt.title('Population Mean: '+str(u_0)+' , Sample Size: '+str(n)+'\n Mean: '+str(round(
16    statistics.mean(expected),3))+', Variance: '+str(round(statistics.variance(expected),3)))
17    plt.savefig("Q5_histograms/Q5.2.1.png")
18    plt.show()
```

Histogram of expected values of the null hypothesis being rejected:



The histogram follows a normal distribution with mean 0.951 and variance 0.001. According to this, the null hypothesis is rejected 95% of the times so it is false. A single hypothesis test may have been sufficient to accept or reject the null hypothesis as the variance is very low.




## 5.2.2

The following function calls the `hypothesis_test()` function defined in 5.2.1 using the same value for the population mean  $u_0$  (23) but passing a value of 70 for the sample size  $n$ . It then performs 50 experiments and uses the average of their outcome to calculate a single expected value. It repeats this for 100 iterations and plots a histogram of the expected values.

```
1 def experiments_updated():
2     u_0 = 23 #population mean
3     n = 70 #sample size
4     expected = []
5     for j in range(100):
6         outcomes = []
7         for i in range(50):
8             outcomes.append(hypothesis_test(u_0,n))
9         expected.append(sum(outcomes)/50)
10    #plotting histogram
11    fig, ax = plt.subplots(figsize =(10, 7))
12    ax.hist(expected, width = 0.1)
13    ax.set_xlabel('Times the null hypothesis is rejected')
14    ax.set_ylabel('Frequency')
15    plt.title('Population Mean: '+str(u_0)+' , Sample Size: '+str(n)+'\n Mean: '+str(round(
16    statistics.mean(expected),3))+', Variance: '+str(round(statistics.variance(expected),3)))
17    plt.savefig("Q5_histograms/Q5.2.2.png")
18    plt.show()
```

Histogram of expected values of the null hypothesis being rejected:



Increasing the value of  $n$  made the variance become 0 where it was previously 0.001. In other words, the null hypothesis is rejected 100% of the times. A single hypothesis test may have been sufficient to reject the null hypothesis in this case as there is no variance in the outcome.