

An explicit weighted digit construction for small sumsets and large difference sets

Benjamin Guo
Hraness (hraness.com)*

28 September 2026

Abstract

The small-sumset, large-difference-set problem asks how large $|E - F|$ can be when $|E + F| \leq K|E|$, for fixed $K > 1$ and arbitrarily large finite integer sets E, F . We give a weighted digit construction for its exponent θ . Equal counts of opposite differences give a lower bound for the difference set, while a product-weight estimate bounds the sumset from above. The resulting formula applies to any finite digit alphabet. For the alphabet $\{0, 3, 4, 6, 7, \dots, 16\}$ and geometric weights $(16/19)^a$, it proves $\theta > 1.1855$. The numerical conclusion reduces to one explicit integer inequality. This exceeds Zheng's published lower bound of 1.173077. The proof uses the finite-set transfer inequality of Gyarmati, Hennecart and Ruzsa and does not require optimality of the chosen parameters.

Keywords. sumset; difference set; weighted digits; entropy; exact certificate. **MSC 2020.** 11B75; 05A16.

1 The exponent and the finite-set reduction

Let θ be the supremum of the real numbers t with the following property: for every fixed $K > 1$, there is a constant $c(K) > 0$ and arbitrarily large finite sets $E, F \subset \mathbb{Z}$ satisfying

$$|E + F| \leq K|E|, \quad |E - F| \geq c(K)|E + F|^t. \quad (1)$$

Here $E \pm F = \{e \pm f : e \in E, f \in F\}$, and all logarithms are natural. The constant may depend on the fixed exponent t as well as on K .

The lemma in Section 2 of Gyarmati, Hennecart and Ruzsa [1] gives a finite-set transfer inequality: for every finite $U \subset \mathbb{Z}_{\geq 0}$ with $0 \in U$, $\max U > 0$ and $|U - U| < 2 \max U + 1$,

$$\theta \geq 1 + \frac{\log(|U - U|/|U + U|)}{\log(2 \max U + 1)}. \quad (2)$$

For each such U , its exponent is attained for every fixed $K > 1$ and arbitrarily large sets. The strict difference-count hypothesis matters; we will ensure it by dilating each encoded set. A family whose difference-to-sum ratio grows exponentially relative to its diameter then gives a lower bound for the supremum θ .

Zheng uses integer vectors whose coordinates lie in a full interval and whose coordinate sum is bounded. His large-deviation estimates give the lower bound 1.173077 [2]. We allow a smaller coordinate alphabet and general nonnegative digit costs. Matching the counts of a difference

*Developed with AI research agents. The proof is accompanied by a separate verifier using exact integer arithmetic.

and its negative makes the costs of its two representatives exactly equal. This symmetry gives a direct counting argument without optimizing a large-deviation rate function. The constructions below improve the bound in Zheng's stated theorem.

A separate problem concerns the one-set quantity

$$C(A) = \frac{\log(|A + A|/|A|)}{\log(|A - A|/|A|)}, \quad A \subset \mathbb{Z}, \quad |A| > 1,$$

whose supremum is 2 by Lin and Li [3]. Numerical comparisons must distinguish $C(A)$ from θ . The "Sets" comparison in Table 1 of Beck, Ogren and Kobren [4] reports a value of 1.21 from Hyra, while its Appendix A, equation (7), and Appendix D use the objective considered here. Lin and Li [3, Section 3] identify that 1.21 search value with $C(A)$. The comparison therefore does not establish a 1.21 lower bound for θ .

2 A weighted digit construction

Fix an integer $B \geq 1$ and a digit alphabet $\mathcal{A} \subseteq \{0, \dots, B\}$ containing at least two elements. Attach a positive weight w_a to each $a \in \mathcal{A}$. Multiplying every weight by a common factor changes neither of the ratios below, so assume $\max_{a \in \mathcal{A}} w_a = 1$. Put

$$\mathcal{S} = \mathcal{A} + \mathcal{A}, \quad S_s = \max_{a+b=s} w_a w_b, \quad P = \sum_{s \in \mathcal{S}} S_s, \quad (3)$$

$$\mathcal{D} = \mathcal{A} - \mathcal{A}, \quad D_d = \max_{a-b=d} w_a w_b, \quad Q = \sum_{d \in \mathcal{D}} D_d. \quad (4)$$

All pairs in these maxima belong to $\mathcal{A}^2$; only represented sums and differences are included.

Theorem 1. *If $Q > P$, then*

$$\theta \geq 1 + \frac{\log(Q/P)}{\log(2B+1)}. \quad (5)$$

Proof. Write $c_a = -\log w_a \geq 0$ and $\tau_d = -\log D_d$. Choose a pair (a_d, b_d) attaining D_d for each positive $d \in \mathcal{D}$, and set $(a_{-d}, b_{-d}) = (b_d, a_d)$. For $d = 0$, use (a_*, a_*) , where $w_{a_*} = 1$. Then $\tau_0 = 0$ and $\tau_{-d} = \tau_d$. Define

$$p_d = \frac{D_d}{Q}, \quad r = \frac{1}{2} \sum_{d \in \mathcal{D}} p_d \tau_d, \quad V_m = \left\{ x \in \mathcal{A}^m : \sum_{i=1}^m c_{x_i} \leq rm \right\}.$$

The probabilities p_d are positive and satisfy $p_{-d} = p_d$.

Difference count. For $d \neq 0$, take $n_d = \lfloor mp_d \rfloor$ and put $n_0 = m - \sum_{d \neq 0} n_d$. Hence $n_d = n_{-d}$ and $n_d/m \rightarrow p_d$ for every d . Consider any ordering $z \in \mathcal{D}^m$ of this multiset of differences. Replace each coordinate d by the selected pair (a_d, b_d) , obtaining vectors x, y with $x - y = z$. Their two costs are equal, since opposite differences occur equally often. Each cost is

$$\sum_{\substack{d \in \mathcal{D} \\ d > 0}} n_d (c_{a_d} + c_{b_d}) \leq m \sum_{\substack{d \in \mathcal{D} \\ d > 0}} p_d \tau_d = mr.$$

The zero coordinates contribute no cost. Thus every such z belongs to $V_m - V_m$. The number of distinct orderings is $m! / \prod_{d \in \mathcal{D}} n_d!$. Stirling's formula gives

$$\log |V_m - V_m| \geq mH(p) - o(m), \quad (6)$$

$$H(p) = - \sum_{d \in \mathcal{D}} p_d \log p_d = \log Q + \sum_{d \in \mathcal{D}} p_d \tau_d = \log Q + 2r. \quad (7)$$

Sum count. If $z = x + y$ for $x, y \in V_m$, the defining maximum in (3) implies

$$\prod_{i=1}^m S_{z_i} \geq \prod_{i=1}^m w_{x_i} w_{y_i} \geq e^{-2rm}.$$

Summing these nonnegative products over all of $\mathcal{S}^m$ yields P^m . Consequently

$$|V_m + V_m| \leq e^{2rm} P^m. \quad (8)$$

Combining (6)–(8),

$$\log \frac{|V_m - V_m|}{|V_m + V_m|} \geq m \log(Q/P) - o(m). \quad (9)$$

Encoding as integers. Set $L = 2B + 1$ and $\phi_m(x) = \sum_{i=1}^m x_i L^{i-1}$. This map is injective on both $\{0, \dots, 2B\}^m$ and $\{-B, \dots, B\}^m$. Indeed, if two different strings first differ at their highest index j , that term has absolute value at least L^{j-1} , whereas all lower terms together have absolute value at most $2B \sum_{i < j} L^{i-1} = L^{j-1} - 1$.

Let $U_m = \phi_m(V_m) - \min \phi_m(V_m)$. It contains zero, its sum and difference cardinalities equal those of V_m , and

$$2 \max U_m + 1 \leq 2B \sum_{i=1}^m L^{i-1} + 1 = L^m. \quad (10)$$

For all sufficiently large m , the logarithm in (9) is positive, so $M_m = \max U_m > 0$. Set $W_m = \{2u : u \in U_m\}$. This dilation preserves both sum and difference cardinalities and ensures the strict hypothesis of (2), because

$$|W_m - W_m| = |U_m - U_m| \leq 2M_m + 1 < 4M_m + 1 = 2 \max W_m + 1.$$

Moreover, (10) gives $2 \max W_m + 1 \leq 2L^m - 1$. Applying (2) to W_m yields

$$\theta \geq 1 + \frac{m \log(Q/P) - o(m)}{m \log L + \log 2}.$$

Taking $m \rightarrow \infty$ proves (5) for the supremum θ . In particular, every exponent strictly below its right-hand side is attainable in (1); we do not assert attainment of the limit. $\square$

3 Sparse geometric weights

For an alphabet containing zero and geometric weights $w_a = q^a$, with $0 < q < 1$, the two totals have a particularly simple form:

$$P(q) = \sum_{s \in \mathcal{A} + \mathcal{A}} q^s, \quad Q(q) = \sum_{d \in \mathcal{A} - \mathcal{A}} q^{e(d)}, \quad e(d) = \min_{a-b=d} (a+b). \quad (11)$$

Thus both totals are polynomials with nonnegative integer coefficients. The formula follows directly from $w_a w_b = q^{a+b}$ and the strict decrease of q^j with j .

The finite integer family is also explicit. Define

$$\rho(q) = \frac{qQ'(q)}{2Q(q)}, \quad V_m = \left\{ x \in \mathcal{A}^m : \sum_{i=1}^m x_i \leq \lfloor m\rho(q) \rfloor \right\}. \quad (12)$$

The radius r in Theorem 1 is $(-\log q)\rho(q)$, so (12) gives precisely its sets V_m . Multiply their base- $(2B+1)$ encodings by two, as in that proof, to obtain the sets used in the transfer inequality. Zero already belongs to this integer family, and $\rho(q)$ is rational whenever q is rational.

3.1 A one-digit omission

Take $B \geq 3$ and $\mathcal{A}_B = \{0, 2, 3, \dots, B\}$. Its sumset is $\{0, 2, 3, \dots, 2B\}$ and its difference set is $\{-B, \dots, B\}$. For $|d| \geq 2$, the pair $(|d|, 0)$ or its reversal minimizes $a + b$, so $e(d) = |d|$. The difference 1 first occurs as $3 - 2$, giving $e(1) = e(-1) = 5$, and $e(0) = 0$. Therefore

$$P_B(q) = \sum_{j=0}^{2B} q^j - q, \quad Q_B(q) = 1 + 2q^5 + 2 \sum_{j=2}^B q^j. \quad (13)$$

The omitted digit removes one sum, while every difference remains represented. This changes the weighted ratio used by Theorem 1.

Corollary 2. *The exponent in (1) satisfies $\theta > 1.1813$.*

Proof. Set $B = 11$ and $q = 377/500$ in (13). Define the positive integers

$$N_S = \sum_{\substack{0 \leq j \leq 22 \\ j \neq 1}} 377^j 500^{22-j}, \quad (14)$$

$$N_D = 500^{22} + 2 \cdot 377^5 500^{17} + 2 \sum_{j=2}^{11} 377^j 500^{22-j}. \quad (15)$$

Then $Q_{11}(q)/P_{11}(q) = N_D/N_S$. Exact integer exponentiation verifies

$$N_D^{10000} > 23^{1813} N_S^{10000}. \quad (16)$$

The accompanying verifier checks this inequality with integer arithmetic. Taking logarithms in (16) and applying Theorem 1 gives

$$\theta \geq 1 + \frac{\log(N_D/N_S)}{\log 23} > 1 + \frac{1813}{10000} = 1.1813.$$

□

3.2 Omitting the digits 1, 2 and 5

Let $B \geq 8$ and $\mathcal{C}_B = \{0, 3, 4, 6, 7, \dots, B\}$. Its sumset is $\{0, \dots, 2B\} \setminus \{1, 2, 5\}$ and its difference set is $\{-B, \dots, B\}$. The three missing positive digits still occur as differences: $1 = 4 - 3$, $2 = 6 - 4$ and $5 = 8 - 3$. These pairs minimize the sum of their endpoints, giving $e(\pm 1) = 7$, $e(\pm 2) = 10$ and $e(\pm 5) = 11$. Every other nonzero difference uses a digit and zero. Thus

$$P(q) = 1 + q^3 + q^4 + \sum_{j=6}^{2B} q^j, \quad (17)$$

$$Q(q) = 1 + 2 \left(q^3 + q^4 + \sum_{j=6}^B q^j + q^7 + q^{10} + q^{11} \right). \quad (18)$$

Corollary 3. *The exponent in (1) satisfies $\theta > 1.1855$.*

Proof. Take $B = 16$, $q = 16/19$ and $T_j = 16^j 19^{32-j}$. By (17)–(18), $Q(q)/P(q) = M_D/M_S$, where

$$M_S = T_0 + T_3 + T_4 + \sum_{j=6}^{32} T_j, \quad (19)$$

$$M_D = T_0 + 2 \left(T_3 + T_4 + \sum_{j=6}^{16} T_j + T_7 + T_{10} + T_{11} \right). \quad (20)$$

The exact integer check below verifies

$$M_D^{2000} > 33^{371} M_S^{2000}. \quad (21)$$

Consequently Theorem 1 gives

$$\theta \geq 1 + \frac{\log(M_D/M_S)}{\log 33} > 1 + \frac{371}{2000} = 1.1855.$$

□

4 Exact reproduction and scope

The certificate can be checked with the following complete Python program. It uses integer arithmetic and the standard library only.

```
from math import gcd

p, q, B = 16, 19, 16
term = lambda j: p**j * q**(2*B-j)
S = term(0) + term(3) + term(4) + sum(term(j) for j in range(6, 33))
D = term(0) + 2*(term(3) + term(4) + sum(term(j) for j in range(6, 17))
    + term(7) + term(10) + term(11))
g = gcd(S, D)
if (D//g)**2000 <= 33**371 * (S//g)**2000:
    raise ArithmeticError("The certificate failed")
print("Verified: theta > 1.1855")
```

The accompanying verifier also constructs the maxima in (3)–(4) directly from all ordered digit pairs and compares them with the displayed polynomial formulas. The integer inequality, rather than a rounded logarithm or a numerical optimizer, certifies the stated decimal threshold.

The theorem holds for every fixed alphabet and every set of positive weights satisfying $Q > P$. The selected parameters need not maximize its right-hand side. Nor does a finite search over alphabets determine the optimal exponent in (1). This note records the construction, its proof and an exact certificate. It does not assert that the bound or method is new, or that the parameters attain a current record.

References

- [1] K. Gyarmati, F. Hennecart and I. Z. Ruzsa, *Sums and differences of finite sets*, Functiones et Approximatio Commentarii Mathematici **37**(1) (2007), 175–186. <https://gyarmatikati.web.elte.hu/publ/sumdiffv.pdf>.
- [2] F. Zheng, *Sums and differences of sets: a further improvement over AlphaEvolve*, arXiv:2506.01896 (2025). <https://arxiv.org/abs/2506.01896>.
- [3] H. Lin and S. Li, *Settling the optimal exponent relating sumsets and difference sets*, arXiv:2607.27199v1 (2026). <https://arxiv.org/abs/2607.27199v1>.
- [4] J. Beck, P. V. Ogren and A. Kobren, *Hill Sampling for Test-Time Scaling: A Simple and Better Alternative to Repeated Sampling, Evolution, and Training*, arXiv:2609.25510v2 (2026). <https://arxiv.org/abs/2609.25510v2>.