STUDENT NUMBER:

APSC 171 - Term Test 1
Tuesday October 3, 2023

A. Ableson S. Pradhan K. Zhang

Last Name (BLOCK LETTERS): _____

First Name: _____

Student #: _____

- This is a 2 hour test.
- No textbooks, notes, or aids other than your calculator (Casio 991 series) are allowed.
- For full marks, you must show all your work and explain how you arrived at your answers, unless explicitly told to do otherwise.
- Wherever appropriate, **include units in your answers**.
- When drawing graphs, **add labels and scales on all axes**.
- Correct final answers with incorrect support may be given no credit.
- While waiting to start, fill in your name and student number above.
- If you need more room, there are blank pages at the end of the test. **If you use these pages, you must provide clear directions on the original page**, e.g. "Continued on page 10".
- Only answers written in this booklet will be graded; all scrap paper from the test will be discarded.

Q 1	Q 2	Q 3	Q 4	Q 5	Q 6	Q 7	Q 8	Total
8	8	9	8	8	5	9	5	60

STUDENT NUMBER:

1. Find the derivative of the following functions. You may assume that the input variables will always be within the domain of their respective functions.
   You do **not** need to simplify after you have taken the derivative.

   (a) $f(x) = 3x^2 \cos(4x)$. [/2]

   (b) $g(t) = \dfrac{e^{at}}{1 - e^{bt}}$. Both a and b are constants. [/3]

   (c) $h(x) = \ln\left(3\sqrt{x^2 + 1}\right)$. [/3]

STUDENT NUMBER:

2. The following three parts are independent of each other.

(a) Find the domain of the function $\vec{r}(t) = \left\langle 2\ln(t^2+1),\ \dfrac{2t}{t^2-4},\ \sqrt{1-t} \right\rangle$.
Express the domain as an interval, or a union of intervals. E.g. "$t \in [-2, 2)$", or "$t \in (-\infty, 2) \cup [3, \infty)$". [/2]

(b) Compute the angle between the vectors $\vec{a} = \langle -2, 5, 1 \rangle$ and $\vec{b} = \langle 2, 1, 2 \rangle$.
Your answer should be in radians, and in the range $[0, \pi]$; report your answer to 4 significant figures. [/2]

(c) Find the exact value of $x = \arcsin\left(\sin\left(\dfrac{11\pi}{4}\right)\right)$. [/2]

(d) Express the relationship $y = \tan\left(\arccos\left(\dfrac{5}{x}\right)\right)$ without trigonometric functions. [/2]

STUDENT NUMBER:

3. Consider a particle whose trajectory is defined by $\vec{s}(t) = \langle 3\sin(2t) + 2, \ -3\cos(2t) + 4 \rangle$.

   (a) The particle's trajectory lies on one of the circles defined by the Cartesian equations below:

   - Circle A: $(x+2)^2 + (y+4)^2 = 9$
   - Circle B: $(x+2)^2 + (y-4)^2 = 9$
   - Circle C: $(x-2)^2 + (y-4)^2 = 9$

   Circle the correct formula and then support your answer with an appropriate calculation. [/2]


   (b) Find the velocity vector for the particle's trajectory. [/2]

   (c) Find the velocity vector for the particle specifically at $t = \frac{\pi}{3}$, then compute the **unit** vector pointing in the direction of the velocity at that time. [/2]

   unit vector = _____ [/3]

   (d) On the grid to the right, draw x and y axes then sketch the graph of the trajectory.
   Once you have drawn the graph:

   - Draw arrows on the trajectory to indicate the direction of motion;
   - Add your answer to part (c) at the appropriate location on the trajectory.




STUDENT NUMBER:

4. The following three parts are independent of each other.

   (a) Find a value s for which the vectors $\vec{a} = \langle s,\ s-1 \rangle$ and $\vec{b} = \langle s^2,\ s \rangle$ will be parallel. [/3]

   (b) On the axes to the right, sketch the graph of the function $y = \arcsin(2x)$. [/3]
   - Use filled circles for endpoints that are reached on the graph (end of a closed interval), and
   - open circles for endpoints that are not reached (end of an open interval).

   Clearly indicate the scale on your axes



   (c) Compute the derivative of $f(x) = \arctan(4x + 5)$. [/2]


STUDENT NUMBER:

5. The position of a particle is defined by the vector-valued function $\vec{r}(t) = \langle t^2 - 1,\ 2 - t \rangle$.

(a) Eliminate parameters to find a Cartesian equation for the function $\vec{r}(t)$. [/3]

(b) On the axes to the right, sketch the graph of the Cartesian equation you found in part (a). Once you have drawn the graph: [/3]

- Clearly indicate what part of this graph the function $\vec{r}(t)$ actually passes over, and
- draw arrows to indicate the direction of motion.



(c) Find a formula for the acceleration of the particle at time t. [/2]

STUDENT NUMBER:

6. This question involves the same vector-valued function as the previous question:


$$\vec{r}(t) = \langle t^2 - 1,\ 2 - t \rangle.$$

For what value(s) of t does a line tangent to the trajectory pass through the point (-4 ,2)? [/5]

STUDENT NUMBER:

7. You throw a ball down the side of a hill. Relative to the xy coordinate axes shown in the diagram, the velocity of the ball at time t (in seconds) is given by

$$\vec{v}(t) = \langle \underbrace{6t+3}_{v_x(t)}, \underbrace{-12t+5}_{v_y(t)} \rangle \text{ m/s.}$$



(a) Give a function $\vec{r}(t) = \langle x(t), y(t) \rangle$ which describes the position of the ball at time t. Assume that the ball is thrown from $\langle x, y \rangle = \langle 0, 0 \rangle$ at $t = 0$. [/3]


(b) At what time t will the ball hit the hillside? [/2]

(c) Since there is no wind, the only acceleration on the ball is due to gravity. Give a vector in xy coordinates which points in the direction of gravity. [/2]

(d) The gravity on Earth, Mars and Neptune are 9.8m/s^2, 3.8m/s^2 and 13.3m/s^2 respectively. Which planet are we on? Be sure to justify your answer. [/2]

STUDENT NUMBER:

8. Assume time t runs from zero to 2π and that the unit circle has been marked with the hours as a clock face (see diagram at right). Match each of the parametric equations with the best description of the curve from the following list. For this question, no explanation or calculations are necessary.



A. Starts at 12 o'clock and moves clockwise one time around.

B. Starts at 12 o'clock and moves counterclockwise one time around.

C. Starts at 6 o'clock and moves counterclockwise one time around.

D. Starts at 3 o'clock and moves counterclockwise one time around.

E. Starts at 9 o'clock and moves counterclockwise one time around.

F. Starts at 9 o'clock and moves clockwise to 3 o'clock.

G. Starts at 9 o'clock and moves counterclockwise two times around.

H. Starts at 9 o'clock and moves clockwise two times around.

[/5]

(a) $\vec{r}(t) = \left\langle -\cos\left(\frac{t}{2}\right), \sin\left(\frac{t}{2}\right)\right\rangle$ matches description _____

(b) $\vec{r}(t) = \langle -\cos(2t), -\sin(2t)\rangle$ matches description _____

(c) $\vec{r}(t) = \langle \cos(t), \sin(t)\rangle$ matches description _____

(d) $\vec{r}(t) = \langle \sin(t), \cos(t)\rangle$ matches description _____

(e) $\vec{r}(t) = \langle -\cos(t), -\sin(t)\rangle$ matches description _____

STUDENT NUMBER:

Space for additional work. **Indicate clearly which question you are continuing if you use this space.**

STUDENT NUMBER:

Space for additional work. **Indicate clearly which question you are continuing if you use this space.**

STUDENT NUMBER:

Space for additional work. **Indicate clearly which question you are continuing if you use this space.**