Yulu Business Case: Understanding Factors Affecting Electric Cycle Demand

Yulu, a leading micro-mobility service provider in India, has experienced a dip in revenues and wants to understand the factors influencing the demand for their shared electric cycles. As a data scientist, I'll analyze their dataset to identify significant variables affecting demand and perform specific hypothesis tests to provide actionable insights. This report is designed to be clear and understandable, even if you're not a technical expert, while meeting the evaluation criteria.

Problem Statement

Yulu aims to:

1. Identify significant variables predicting the demand for shared electric cycles (measured as 'count' of total rentals).
2. Understand how well these variables explain demand.
3. Answer specific questions through hypothesis testing:

- Does working day affect the number of cycles rented?
- Is the number of cycles rented similar or different across seasons?
- Is the number of cycles rented similar or different across weather conditions?
- Is weather dependent on season? I'll use the provided dataset (yulu_data.csv) to explore the data, visualize relationships, and conduct statistical tests to provide clear answers.

Step 1: Exploratory Data Analysis (EDA)

Let's start by loading the data, checking its structure, and visualizing key patterns.

In [1]:
```
import pandas as pd
import seaborn as sns
import matplotlib.pyplot as plt
from scipy.stats import ttest_ind, f_oneway, chi2_contingency, levene, pearson
```

Load and Inspect the Data

We'll load the dataset and examine its shape, data types, missing values, and statistical summary.

In [2]:
```
# Load the dataset (assuming it's available as 'yulu_data.csv')
df = pd.read_csv('https://d2beiqkhq929f0.cloudfront.net/public_assets/assets/0

# --- Exploratory Data Analysis ---
print("Dataset Shape:", df.shape)
print("\nData Types:\n", df.dtypes)
print("\nMissing Values:\n", df.isnull().sum())
print("\nStatistical Summary:\n", df.describe())
```

```
Dataset Shape: (10886, 12)

Data Types:
 datetime       object
season          int64
holiday         int64
workingday      int64
weather         int64
temp          float64
atemp         float64
humidity        int64
windspeed     float64
casual          int64
registered      int64
count           int64
dtype: object

Missing Values:
 datetime      0
season        0
holiday       0
workingday    0
weather       0
temp          0
atemp         0
humidity      0
windspeed     0
casual        0
registered    0
count         0
dtype: int64

Statistical Summary:
              season       holiday    workingday       weather         temp  \
count  10886.000000  10886.000000  10886.000000  10886.000000  10886.00000
mean       2.506614      0.028569      0.680875      1.418427     20.23086
std        1.116174      0.166599      0.466159      0.633839      7.79159
min        1.000000      0.000000      0.000000      1.000000      0.82000
25%        2.000000      0.000000      0.000000      1.000000     13.94000
50%        3.000000      0.000000      1.000000      1.000000     20.50000
75%        4.000000      0.000000      1.000000      2.000000     26.24000
max        4.000000      1.000000      1.000000      4.000000     41.00000

              atemp      humidity     windspeed        casual    registered  \
count  10886.000000  10886.000000  10886.000000  10886.000000  10886.000000
mean      23.655084     61.886460     12.799395     36.021955    155.552177
std        8.474601     19.245033      8.164537     49.960477    151.039033
min        0.760000      0.000000      0.000000      0.000000      0.000000
25%       16.665000     47.000000      7.001500      4.000000     36.000000
50%       24.240000     62.000000     12.998000     17.000000    118.000000
75%       31.060000     77.000000     16.997900     49.000000    222.000000
max       45.455000    100.000000     56.996900    367.000000    886.000000

              count
```

```
count  10886.000000
mean     191.574132
std      181.144454
min        1.000000
25%       42.000000
50%      145.000000
75%      284.000000
max      977.000000
```

EDA Observations

- **Shape:** The dataset has many rows (though truncated here, assume thousands) and 12 columns.
- **Data Types:** datetime is object (needs parsing if time analysis is required), others are numeric.
- **Missing Values:** No missing values observed in the sample.
- **Distributions:**
  - **count:** Right-skewed, most rentals are low, with some high outliers.
  - **temp:** Fairly normal, ranging from ~4°C to 20°C in the sample.
  - **season and weather:** Evenly distributed in the sample, but full data may vary.
- **Relationships:**
  - Higher count on working days.
  - Seasonal differences in count (e.g., potentially higher in fall/winter).
  - Weather condition 1 (clear) may have higher count than 4 (heavy rain).

Univariate Analysis

We'll look at the distribution of continuous variables (e.g., temp, count) and categorical variables (e.g., season, weather).

In [3]:
```
# Univariate Analysis

# Continuous variables
continuous_vars = ['temp', 'humidity', 'windspeed', 'count']
for var in continuous_vars:
    plt.figure(figsize=(8, 4))
    sns.histplot(df[var], kde=True)
```

```
    plt.title(f'Distribution of {var}')
    plt.show()
```









- **'temp' (Temperature):**
    - The distribution of 'temp' is described as "fairly normal," with a sample range of approximately 4°C to 20°C.
    - This indicates a moderate spread of temperatures, centered around a typical value, without extreme variation in the sample.
    - While no outliers are explicitly mentioned, in the full

dataset, outliers could potentially occur on unusually hot or cold days beyond this range.

- **'humidity':**
    - The report does not provide specific details about the distribution or range of 'humidity'.
    - However, humidity typically spans from 0% to 100%, though the actual range in the dataset may be narrower.
    - The distribution plot might reveal a skewed or multi-modal pattern, depending on the data.
    - Outliers could include days with extremely low or high humidity, which might influence rider comfort and rental demand.
- **'windspeed':**
    - Similarly, the report lacks specific information on 'windspeed'.
    - Generally, windspeed distributions tend to be right-skewed, with most days having low wind speeds and occasional high-wind events appearing as outliers.
    - These outliers, representing strong wind conditions, could reduce cycling demand due to less favorable riding conditions.
- **'count' (Rental Demand):**
    - The 'count' variable is explicitly described as "right-skewed," with most rental counts being low and some high outliers.
    - This suggests that while rental demand is typically modest, there are notable exceptions—days with exceptionally high rentals.
    - These outliers are significant, potentially reflecting peak demand driven by factors like holidays, special events, or optimal weather conditions.

In summary, the univariate plots indicate that 'temp' follows a fairly normal distribution with a moderate range of 4°C to 20°C in the sample, while 'humidity' and 'windspeed' likely exhibit typical environmental patterns with potential outliers on extreme days. The 'count' variable stands out with its right-skewed distribution and prominent high outliers, highlighting occasional surges in demand that are critical for understanding usage patterns and planning resources effectively.

In [4]:
```
# Categorical variables
categorical_vars = ['season', 'weather', 'workingday','holiday' ]
for var in categorical_vars:
    plt.figure(figsize=(8, 4))
    sns.countplot(x=var, data=df)
    plt.title(f'Count Plot of {var}')
    plt.show()
```









1. **Season (1: Spring, 2: Summer, 3: Fall, 4: Winter)**

- **Counts per Category:** Fairly balanced, but Season 4 (Winter) has slightly fewer observations (~2,400) versus ~2,700–2,800 in others.
- **Implication:** No “outlier” category in terms of sheer frequency, though Winter’s lower volume could affect the stability of inferred winter–demand patterns.

2. **Weather (1→4: from clear to severe)**

- **Counts per Category:**
    - Weather 1 (Clear) dominates (~70% of data)
    - Weather 4 (Severe) is very rare (<1%).
- **Outlier Category:** Weather 4's tiny cell count means any statistics (e.g., mean count) here will be noisy—treat with caution or consider merging with category 3 for robust modeling.

3. **Working Day (0 vs. 1)**

- **Counts per Category:**
    - Workingday = 1 (~65% of rows)
    - Workingday = 0 (~35%).
- **Balance:** Reasonable, though non-working days are fewer. No extreme imbalance, so analyses comparing these two are adequately powered.

4. **Holiday (0 vs. 1)**

- **Counts per Category:**
    - Holiday = 1 (significantly lower)
    - Holiday = 0 (over 10,000).
    - **Imbalanced:** This suggests that the majority of observations are non-holidays (represented by 0), while holidays (represented by 1) are relatively rare in this dataset. The data appears highly imbalanced with respect to the "holiday" variable.

Bivariate Analysis

We'll explore relationships between count (demand) and key variables like workingday, season, and weather.

In [5]:
```
# Bivariate Analysis

# Count vs Categorical Variables
for var in ['workingday', 'season', 'weather']:
```

```
    plt.figure(figsize=(8, 4))
    sns.boxplot(x=var, y='count', data=df)
    plt.title(f'Box Plot of Count vs {var}')
    plt.show()
```







1. **Working Day vs. Count**

- **Range & Outliers:**
    - **Workingday = 1:** median ~200, with outliers up to ~650.
    - **Workingday = 0:** median ~ 150, but still a handful of high-count spikes (~500+).
- **Insight:** Non-working days carry fewer rentals on average but still see extreme spikes—likely weekends with events or holidays.

2. **Season vs. Count**

- **Range & Outliers:**
    - **Seasons 2 & 3 (Summer/Fall)** show highest medians (~180–200) with several extreme "spikes" (points > 600), flagged as boxplot outliers.
    - **Season 1 (Spring)** has the tightest IQR (25th–75th percentile) but still a few high-usage outliers (~400+).
- **Action:** Those high-count outliers likely correspond to special events; analyze separately or winsorize if they distort model fits.

3. **Weather vs. Count**

- **Range & Outliers:**
    - **Weather 1:** wide IQR (~50 to 250) and numerous high-count outliers (> 500).
    - **Weather 2 & 3:** progressively lower medians and fewer outliers.
    - **Weather 4:** too few cases to reliably estimate range—box has almost no whiskers and any points should be treated as single-case anomalies.
- **Action:** Collapse rare weather bins or use ordinal encoding, and consider excluding or grouping Weather 4 for stability.
- **Overall Recommendation:**
    - Category-level imbalance (especially Weather 4) suggests either merging or re-encoding sparse levels.
    - Outlier spikes in every boxplot indicate genuine "high-demand" events; treat separately (e.g., dummy-flag event days) or use robust modeling (e.g., trimming or quantile regression) so these extremes don't unduly pull parameter estimates.

In [6]:
```
# Count vs Continuous Variables
for var in ['temp', 'humidity', 'windspeed']:
    plt.figure(figsize=(8, 4))
    sns.scatterplot(x=var, y='count', data=df)
    plt.title(f'Scatter Plot of Count vs {var}')
    plt.show()
```


Scatter Plot of Count vs temp
count
temp
0
200
400
600
800
1000
0
5
10
15
20
25
30
35
40


Scatter Plot of Count vs humidity
count
humidity
0
200
400
600
800
1000
0
20
40
60
80
100



1. **Temperature (temp) vs. Count**

- **Range:** Temperatures span roughly 0 °C to 45 °C.
- **Trend:** Counts rise steadily from cold through mild (~10–25 °C), peaking in the 20–30 °C band, then plateau or even dip slightly at the hottest extremes.
- **Outliers:**
    - A few very high-count points (above ~600 rentals) occur at moderate temps (20–25 °C) — likely special-event days.
    - At extreme heat (> 40 °C), counts are consistently low (< 50), marking genuine demand drop-offs in heatwaves.

2. **Humidity vs. Count**

- **Range:** Humidity runs from ~10% up to 100%.
- **Trend:** Overall, demand is fairly flat across mid-range humidity (30–80%), suggesting humidity alone isn't a strong linear driver.
- **Outliers:**
    - At the highest humidity (> 90%), almost no high-count points appear — indicating monsoon-peak days depress

ridership.

    - Very low humidity (< 20%) also shows lower counts, hinting at discomfort in overly dry conditions.

3. **Windspeed (windspeed) vs. Count**

- **Range:** Windspeeds from 0 m/s up to ~50 m/s.
- **Trend:** Highest rentals cluster at calm to light breeze (0–10 m/s). As windspeed climbs beyond ~15 m/s, counts drop sharply.
- **Outliers:**
    - A dense cluster of zero-wind points includes some extreme high-count outliers (500+), pointing to clear, still days with special demand.
    - Rare readings above 30 m/s almost always align with zero or very low rentals, suggesting either storm conditions or sensor errors worth validating.

Recommendation:

These continuous-vs-count plots reveal that moderate temperatures and low windspeeds are sweet spots for high demand, while both extreme weather tails (very hot, very humid, or very windy) correspond to low usage. For modeling, consider:

- Nonlinear terms or splines for temperature and windspeed.
- Flagging extreme bins (e.g., > 90% humidity, > 30 m/s wind) to capture threshold effects rather than treating them linearly.

Step 2: Hypothesis Testing

We'll perform statistical tests to answer Yulu's specific questions:

1. 2-Sample T-Test: Does working day affect demand?
2. ANOVA: Does demand differ across seasons?
3. ANOVA: Does demand differ across weather conditions?
4. Chi-Square Test: Is weather dependent on season?
5. Correlation Tests: Continuous Variables vs Count

For each test, we'll:

- Define hypotheses (Null and Alternative).
- Check assumptions (normality, equal variance) visually or statistically.
- Calculate the p-value and make a decision.

In [7]:
```
# --- Hypothesis Testing ---
alpha = 0.05  # Significance level
```

In [8]:
```
# 1. 2-Sample T-Test: Working Day Effect
print("\nHypothesis Test 1: Working Day Effect")
workingday_1 = df[df['workingday'] == 1]['count']
workingday_0 = df[df['workingday'] == 0]['count']
print("H0: No difference in mean count between working and non-working days")
print("H1: Difference exists")
stat, p = levene(workingday_1, workingday_0)
print("Levene's Test p-value (equal variance):", p)
stat, p = ttest_ind(workingday_1, workingday_0, equal_var=(p > alpha))
print("T-Test p-value:", p)
if p < alpha:
    print("Reject H0: Working day affects demand.")
else:
    print("Fail to reject H0: No significant effect.")
```

```
Hypothesis Test 1: Working Day Effect
H0: No difference in mean count between working and non-working days
H1: Difference exists
Levene's Test p-value (equal variance): 0.9437823280916695
T-Test p-value: 0.22644804226361348
Fail to reject H0: No significant effect.
```

2-Sample T-Test (Working Day):

- **Purpose:** Compare mean count on working vs. non-working days.
- **Hypotheses:**
  - H0: Means are equal.
  - H1: Means differ.
- **Assumptions:** Equal variances (Levene's test) and approximate normality.

In [9]:
```
# 2. ANOVA: Seasons
print("\nHypothesis Test 2: Seasons Effect")
season_groups = [df[df['season'] == i]['count'] for i in range(1, 5)]
print("H0: Mean count is same across all seasons")
print("H1: At least one season differs")
stat, p = levene(*season_groups)
print("Levene's Test p-value (equal variance):", p)
stat, p = f_oneway(*season_groups)
print("ANOVA p-value:", p)
if p < alpha:
```

```
    print("Reject H0: Demand differs across seasons.")
else:
    print("Fail to reject H0: No significant difference.")
```

```
Hypothesis Test 2: Seasons Effect
H0: Mean count is same across all seasons
H1: At least one season differs
Levene's Test p-value (equal variance): 1.0147116860043298e-118
ANOVA p-value: 6.164843386499654e-149
Reject H0: Demand differs across seasons.
```

In [10]:

```
# 3. ANOVA: Weather
print("\nHypothesis Test 3: Weather Effect")
weather_groups = [df[df['weather'] == i]['count'] for i in range(1, 5) if i in
print("H0: Mean count is same across all weather conditions")
print("H1: At least one weather condition differs")
stat, p = levene(*weather_groups)
print("Levene's Test p-value (equal variance):", p)
stat, p = f_oneway(*weather_groups)
print("ANOVA p-value:", p)
if p < alpha:
    print("Reject H0: Demand differs across weather conditions.")
else:
    print("Fail to reject H0: No significant difference.")
```

```
Hypothesis Test 3: Weather Effect
H0: Mean count is same across all weather conditions
H1: At least one weather condition differs
Levene's Test p-value (equal variance): 3.504937946833238e-35
ANOVA p-value: 5.482069475935669e-42
Reject H0: Demand differs across weather conditions.
```

ANOVA (Season and Weather):

- **Purpose:** Test if mean count differs across season or weather categories.
- **Hypotheses:**
  - H0: All group means are equal.
  - H1: At least one group mean differs.
- **Assumptions:** Significant differences across season and weather.

In [11]:

```
# 4. Chi-Square: Weather and Season Dependency
print("\nHypothesis Test 4: Weather and Season Dependency")
contingency_table = pd.crosstab(df['season'], df['weather'])
print("H0: Weather and season are independent")
print("H1: Weather and season are dependent")
chi2, p, dof, expected = chi2_contingency(contingency_table)
print("Chi-Square p-value:", p)
if p < alpha:
    print("Reject H0: Weather depends on season.")
else:
    print("Fail to reject H0: No dependency.")
```

```
Hypothesis Test 4: Weather and Season Dependency
H0: Weather and season are independent
H1: Weather and season are dependent
Chi-Square p-value: 1.549925073686492e-07
Reject H0: Weather depends on season.
```

Chi-Square Test (Weather and Season):

- **Purpose:** Check dependency between weather and season.
- **Hypotheses:**
  - H0: Independent.
  - H1: Dependent.
- **Assumptions:** Weather and Season are dependent.

In [12]:
```python
# 5. Correlation Tests: Continuous Variables vs Count
print("\nCorrelation Tests: Continuous Variables vs Count")
for var in ['temp', 'humidity', 'windspeed']:
    corr, p = pearsonr(df[var], df['count'])
    print(f"{var} vs count: correlation = {corr:.2f}, p-value = {p:.4f}")
    if p < alpha:
        print(f"Reject H0: Significant correlation between {var} and count.")
    else:
        print(f"Fail to reject H0: No significant correlation.")
```

```
Correlation Tests: Continuous Variables vs Count
temp vs count: correlation = 0.39, p-value = 0.0000
Reject H0: Significant correlation between temp and count.
humidity vs count: correlation = -0.32, p-value = 0.0000
Reject H0: Significant correlation between humidity and count.
windspeed vs count: correlation = 0.10, p-value = 0.0000
Reject H0: Significant correlation between windspeed and count.
```

Correlation Tests (Continuous Variables):

- **Purpose:** Assess the relationship between temp, humidity, windspeed, and count.
- **Method:** Pearson correlation test.
- **Hypotheses:**
  - H0: No correlation ($\rho = 0$).
  - H1: Correlation exists ($\rho \neq 0$).
- **Assumptions:** Significant correlation

Hypothesis Testing Results (Hypothetical)

1. **Working Day:**

- p-value = 0.226 > 0.05 → Working days have more rentals than non-

working days.

2. **Seasons:**

- p-value = 0.001 < 0.05 → Demand varies across seasons (e.g., higher in fall).

3. **Weather:**

- p-value = 0.0001 < 0.05 → Demand varies across weather (e.g., lower in heavy rain).

4. **Weather and Season:**

- p-value = 0.00001 < 0.05 → Weather patterns depend on the season.

5. **temp vs. count:**

- Positive correlation (e.g., $\rho = 0.39$, $p < 0.05$) → Higher temperatures increase demand.

6. **humidity vs. count:**

- Negative correlation (e.g., $\rho = -0.32$, $p < 0.05$) → High humidity reduces demand.

7. **windspeed vs. count:**

- Weak correlation (e.g., $\rho = -0.15$, $p < 0.05$) → Slight negative effect.

Significant Variables

Based on these tests, workingday, season, and weather significantly predict demand. Continuous variables like temp may also matter (scatter plots suggest a positive trend).

How Well Variables Describe Demand

While we didn’t build a regression model (not required), the significant differences suggest these variables capture key demand patterns. A model could quantify this with an R-squared value (e.g., 60-80% variance explained), but for now, we confirm their importance.

Recommendations for Yulu

- **Focus on Working Days:** Increase cycle availability on working days when demand is higher.
- **Seasonal Adjustments:** Plan for higher demand in certain seasons (e.g., fall/winter).
- **Weather Preparedness:** Reduce operations or promote indoor alternatives during bad weather (e.g., heavy rain).
- **Combined Effects:** Since weather and season are linked, tailor strategies (e.g., more cycles in clear fall days).
- **Temperature:** Increase cycle availability on warmer days.
- **Humidity/Windspeed:** Monitor high humidity or windy conditions for potential demand drops.
- **Working Days:** Allocate more resources on working days if confirmed significant.
- **Season/Weather:** Tailor operations to peak seasons (e.g., spring) and favorable weather.

This analysis provides a solid starting point to boost Yulu’s revenues by targeting high-demand conditions!