

[10]

5. Prove the following identities.

$$(a) \gamma^\alpha (\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) \gamma_\alpha = 0.$$

$$(b) \gamma^0 S^\dagger \gamma^0 = S^{-1}, \text{ where } S \text{ is the Lorentz transformation on the Dirac field.}$$

[5]

6. Calculate the unpolarized cross section for the process  $\psi(p_1) + \bar{\psi}(p_2) \rightarrow \phi(P)$  at the leading order in the rest frame of  $\phi$  in Yukawa theory,

$$\mathcal{L}_{\text{Yukawa}} = \frac{1}{2} (\partial_\mu \phi \partial^\mu \phi - m_\phi^2 \phi^2) + \bar{\psi} (i \not{\partial} - m) \psi - g \bar{\psi} \psi \phi.$$

[5]

7. Consider the leading order scattering  $e^-(p_1) + \mu^-(p_2) \rightarrow e^-(p_3) + \mu^-(p_4)$  in QED,

$$\mathcal{L}_{\text{QED}} = \sum_i \bar{\psi}_i (i \not{\partial} - m_i) \psi_i - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} - e \bar{\psi}_i \gamma^\alpha \psi_i A_\alpha,$$

where  $i = 1, 2$  refer to electron and muon respectively.(a) Calculate spin-summed and averaged squared amplitude in terms of Mandelstam variables  $s, t$  and  $u$ .(b) Calculate differential cross section  $\frac{d\sigma}{d\Omega}$  in the center-of-mass frame and express your result in terms of  $\alpha = \frac{e^2}{4\pi}$ ,  $s$  and  $\cos\theta$  where  $\theta$  is the angle between scattered electron and the incident electron which can be taken along  $z$ -axis.Take  $m_e = m_\mu = 0$ . Use  $u_e(p_1)$  and  $u_\mu(p_2)$  for incoming electron and muon. Use  $\bar{u}_e(p_3)$  and  $\bar{u}_\mu(p_4)$  for outgoing electron and muon. Photon propagator can be taken as:

$$D_A^{\alpha\beta}(p) = \frac{i(-g^{\alpha\beta})}{p^2 + i\epsilon}$$

[10]

$$im = (-ig) \bar{v}(p_2) u(p_1)$$

$$|Tm|^2 = 4g^2 (p_2 \cdot p_1 - m^2)$$

$$\int \frac{d^3x}{(2\pi)^3} e^{-i(p_1 - p_2) \cdot x} = \delta^{(3)}(p_1 - p_2)$$

$$\gamma_i \gamma_i = (\sigma_i \sigma_i) (\sigma_i \sigma_i)$$