



AI SMART FARMING REPORT

Soybean

PREPARED FOR
Hrushikesh Thombare
Sambhajinagar, Maharashtra

LAND AREA
5

GENERATED ON
13 Aug 2026

REPORT ID
FF-AE80B873

Track live updates and AI recommendations at
<https://farmfleet.ai/itinerary/6a7d54e1bef33f53ae80b873>



Farm Information

Details used by the AI to build your farming plan

STATE Maharashtra	DISTRICT Sambhajinagar
SOIL TYPE Black Soil	WATER SOURCE Borewell
BUDGET 100000	CROP Soybean
SEASON Kharif (June - October)	SEED RECOMMENDATION { variety: 'JS 335 / MAUS 158 / Phule Sangam (KDS 726)', seedQuantity: '25 - 30 kg per Acre (125 - 150 kg for 5 Acres)', estimatedCost: ' ₹3,000 - ₹3,500 per Acre (₹15,000 - ₹17,500 total)' }



Crop Timeline

Week-by-week schedule for your crop cycle

WEEK 1

Summer Ploughing & Land Preparation

Plough land deep, apply FYM, pulverize black soil, and form Broad Bed Furrows (BBF) for drainage.

Scheduled: 13 Aug 2026

Scheduled

WEEK 2

Seed Treatment & Line Sowing

Treat seeds with Thiamethoxam 30 FS, Trichoderma, and Rhizobium; sow at 45 cm row distance after 75-100 mm rainfall.

Scheduled: 20 Aug 2026

Scheduled

WEEK 3

Germination Check & Weed Control

Monitor uniform plant emergence and apply pre-emergence herbicide within 48 hours if not done at sowing.

Scheduled: 27 Aug 2026

Scheduled

WEEK 4

Thinning & Inter-cultivation

Thin excess seedlings to maintain 10-12 cm plant-to-plant spacing and run a hoe/kolpa for root aeration.

Scheduled: 03 Sept 2026

Scheduled

WEEK 5

Post-Emergence Weed Management

Apply post-emergence weedicide for selective control of broadleaf weeds and grassy weeds.

Scheduled: 10 Sept 2026

Scheduled

WEEK 6

Vegetative Growth & Foliar Spray

Spray 19:19:19 NPK soluble fertilizer along with micronutrients to accelerate canopy development.

Scheduled: 17 Sept 2026

Scheduled



Crop Timeline (continued)

WEEK 7

Flowering Stage & Pest Inspection

Inspect crop for girdle beetle and stem fly; administer borewell supplementary irrigation if rain spell halts.

Scheduled: 24 Sept 2026

Scheduled

WEEK 8

Pod Initiation & Crop Protection

Spray recommended insecticide for Spodoptera caterpillar and fungicide for leaf spot prevention.

Scheduled: 01 Oct 2026

Scheduled

WEEK 9

Pod Development & Grain Filling

Maintain optimum soil moisture via borewell if needed; apply 0:52:34 foliar spray to boost grain weight.

Scheduled: 08 Oct 2026

Scheduled

WEEK 10

Crop Maturation Stage

Stop irrigation completely as leaves begin yellowing and drop off naturally.

Scheduled: 15 Oct 2026

Scheduled

WEEK 11

Harvesting

Harvest crop when 85-90% of pods turn golden brown and seeds produce a rattling sound inside pods.

Scheduled: 22 Oct 2026

Scheduled

WEEK 12

Threshing, Sun Drying & Storage

Thresh harvested plants using a thresher, dry seeds to below 10% moisture content, and store in jute bags.

Scheduled: 29 Oct 2026

Scheduled



Equipment

Machinery and tools recommended for this crop

Tractor with Mouldboard Plough & Rotavator
Primary tillage, clod crushing, and fine soil preparation

Estimated Rent: ₦1,200 - ₦1,500 per hour

BBF Seed Drill
Precision line sowing with simultaneous broad bed furrow making

Estimated Rent: ₦800 - ₦1,000 per acre

Knapsack Power Sprayer
Spraying weedicides, insecticides, and foliar soluble fertilizers

Estimated Rent: ₦350 - ₦400 per day

Tractor-Driven Soybean Thresher
Threshing and cleaning harvested crop pods

Estimated Rent: ₦1,400 - ₦1,600 per hour



Labour

Estimated workforce required across activities

Land Preparation, Manure Spreading & Bed Formation	WORKERS 4 workers	EST. DAYS 2 days
Seed Treatment, Seed Drill Sowing & Fertilizer Basal Mix	WORKERS 5 workers	EST. DAYS 1 day
Hand Weeding & Inter-cultivation Hoeing	WORKERS 8 workers	EST. DAYS 2 days
Chemical & Foliar Nutrient Spraying	WORKERS 2 workers	EST. DAYS 3 days
Crop Harvesting & Field Bundling	WORKERS 10 workers	EST. DAYS 2 days
Threshing, Seed Drying, Bagging & Storage	WORKERS 6 workers	EST. DAYS 1 day



Fertilizer Schedule

Recommended fertilizer application plan

Stage	Fertilizer	Quantity	Time
Basal Application	DAP + MOP + Sulphur	35 kg DAP + 20 kg MOP + 10 kg Sulphur Granules per acre	At the time of sowing in seed line
Vegetative Stage (25-30 DAS)	Foliar Spray NPK 19:19:19	1 kg per acre in 150 liters water	25 - 30 days after sowing
Pre-Flowering Stage (40-45 DAS)	Foliar Spray 12:61:0 + Micronutrient Boron	1 kg 12:61:0 + 250 g Boron per acre in 200 liters water	40 - 45 days after sowing
Pod Filling Stage (60-65 DAS)	Foliar Spray NPK 0:0:50 (SOP)	1 kg per acre in 200 liters water	60 - 65 days after sowing



Irrigation Schedule

Recommended watering plan for healthy crop growth

Stage	Frequency	Water Requirement
Germination & Seedling Stage	Rainfed / 1 Light irrigation if monsoon delayed	25 - 30 mm light irrigation
Flowering Stage (35-45 DAS)	1 Critical supplementary irrigation from borewell during dry spell	40 - 50 mm flooding in furrows
Pod Filling Stage (60-70 DAS)	1 Critical supplementary irrigation from borewell if dry spell occurs	40 - 50 mm flooding in furrows



Weather Report

Latest conditions used to plan your farming activities

Condition: Sunny / Clear

TEMPERATURE
28°C

HUMIDITY
65%

WIND SPEED
10 km/h

RAIN PROBABILITY
15%

Weather Recommendation

Favorable weather conditions for current crop stage. Proceed with scheduled farming activities and maintain regular field observation.



Precautions

Important safety and crop-care guidelines

- Ensure good field drainage because black clay soil holds excess water, which causes soybean root rot.
- Do not drill seeds deeper than 3 to 4 cm; deep sowing in heavy soil severely reduces seed germination percentage.
- Always wear personal protective kit (mask, gloves, long sleeves) during weedicide and pesticide application.
- Avoid spraying liquid pesticides during strong midday direct heat or windy environmental conditions.

Tips

Best practices to improve yield and reduce costs

- Sow soybean using Broad Bed Furrow (BBF) plan to increase yield by 15-20% and save water during dry spells.
- Inoculate seeds with *Bradyrhizobium japonicum* and PSB cultures to fix natural atmospheric nitrogen and cut urea costs.
- Reserve borewell water strictly for critical flowering (35-45 DAS) and pod filling (55-65 DAS) moisture stress periods.



Your Journey Continues

This report is just the beginning of your AI-powered farming plan

- Your itinerary will continue to receive updates.
- Weather conditions may automatically change your schedule.
- Login to your FarmFleet account to view the latest recommendations.

Continue on FarmFleet

To receive:

- Live Weather Updates
- AI Recommendations
- Updated Farming Schedule
- Equipment Booking
- Labour Hiring

Login at:

<https://farmfleet.ai/itinerary/6a7d54e1bef33f53ae80b873>

REPORT ID

FF-AE80B873

GENERATED DATE

13 Aug 2026