


Determining Galactic Structure through 21cm Emission Lines

Henry Shackleton


May 11, 2017

Hyperfine Transition of Hydrogen Emits 21cm Wavelength Emission




- Hydrogen electron spin-flip causes electromagnetic radiation at a frequency of 1420.41 MHz.
- Low probability ($2.9 \times 10^{-15} \text{ s}^{-1}$), but the vast amount of hydrogen in the galaxy allows for this detection

Doppler Shift Gives Change in 21cm Line Proportional to Velocity




$$v = c \frac{1420.41 - \nu}{\nu}$$

Doppler Shift Gives Change in 21cm Line Proportional to Velocity




$$v = c \frac{1420.41 - \nu}{\nu} - v_{lsr}$$

Location of Hydrogen Masses Determined through Geometry




- Velocity we observe is the velocity of the mass *projected* onto our line of sight.

Location of Hydrogen Masses Determined through Geometry




- Velocity we observe is the velocity of the mass *projected* onto our line of sight.
- $$v_{obs} = \frac{\Theta}{r} R_0 \sin \ell - \Theta_0 \sin \ell$$

Location of Hydrogen Masses Determined through Geometry




- Velocity we observe is the velocity of the mass *projected* onto our line of sight.
- $v_{obs} = \frac{\Theta}{r} R_0 \sin \ell - \Theta_0 \sin \ell$
- Relation between Θ and r obtained through Galactic Rotation Curve.

Location of Hydrogen Masses Determined through Geometry




- Velocity we observe is the velocity of the mass *projected* onto our line of sight.
- $v_{obs} = \frac{\Theta}{r} R_0 \sin \ell - \Theta_0 \sin \ell$
- Relation between Θ and r obtained through Galactic Rotation Curve.
- Between $90^\circ < \ell < 180^\circ$, Galactic Rotation Curve is approximately constant.

SRT Measures Radio Power Within Given Frequency Domain




- Noise diode allows for calibration of telescope.
- Receiver selects desired bandwidth for data collection.

Peak in Antenna Readings Signal Hydrogen Density Concentration



Peak in Antenna Readings Signal Hydrogen Density Concentration



Sources of Error Largely Due to Unknown Constants

- Velocity of sun lacks certainty, approximately 231 ± 14 km/s.

Sources of Error Largely Due to Unknown Constants

- Velocity of sun lacks certainty, approximately 231 ± 14 km/s.
- Distance from sun to center of galaxy also uncertain, 8.2 ± 0.5 kpc.


Sources of Error Largely Due to Unknown Constants

- Velocity of sun lacks certainty, approximately 231 ± 14 km/s.
- Distance from sun to center of galaxy also uncertain, 8.2 ± 0.5 kpc.
- Beam width approximated as constant across five degree bin width - taking into account beam profile would allow for more accurate data analysis


Sources of Error Largely Due to Unknown Constants

- Velocity of sun lacks certainty, approximately 231 ± 14 km/s.
- Distance from sun to center of galaxy also uncertain, 8.2 ± 0.5 kpc.
- Beam width approximated as constant across five degree bin width - taking into account beam profile would allow for more accurate data analysis
- Poissonian uncertainty in number of counts.


Thermal Broadening Irrelevant at Current Galactic Resolution



Hydrogen Mapping to Polar Histogram Indicates Spiral Arm



Hydrogen Mapping Agrees with Density Data



Results Verify Validity of 21cm Analysis

- Hydrogen hyperfine emissions provide a reliable method of detecting hydrogen.

Results Verify Validity of 21cm Analysis

- Hydrogen hyperfine emissions provide a reliable method of detecting hydrogen.
- Analyzing Doppler shift allows for determining of source location and a mapping of hydrogen density.

Results Verify Validity of 21cm Analysis

- Hydrogen hyperfine emissions provide a reliable method of detecting hydrogen.
- Analyzing Doppler shift allows for determining of source location and a mapping of hydrogen density.
- Resolution limited due to telescope size and thermal broadening, but qualitative features agree with literature.