

# Relativistic Behavior Detection through Electron Acceleration

Henry Shackleton

April 5, 2017

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

- Formalized by Newton in 1687

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

- Formalized by Newton in 1687
- No limit to the speed of a particle

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

- Formalized by Newton in 1687
- No limit to the speed of a particle

## Special Relativity

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

- Formalized by Newton in 1687
- No limit to the speed of a particle

## Special Relativity

- Developed by Einstein in 1905

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

- Formalized by Newton in 1687
- No limit to the speed of a particle

## Special Relativity

- Developed by Einstein in 1905
- The speed of light,  $c$ , is constant in all reference frames

# Classical Mechanics and Special Relativity Differ in Predictions

## Classical Mechanics

- Formalized by Newton in 1687
- No limit to the speed of a particle

## Special Relativity

- Developed by Einstein in 1905
- The speed of light,  $c$ , is constant in all reference frames
- The velocity of any particle is capped at  $c$



# Classical and Relativistic Kinetic Energies are Different

## Classical Kinetic Energy

$$K = \frac{p^2}{2m}$$

# Classical and Relativistic Kinetic Energies are Different

## Classical Kinetic Energy


$$K = \frac{p^2}{2m}$$

## Relativistic Kinetic Energy

$$K = \sqrt{p^2 c^2 + m^2 c^4} - mc^2$$

# Electrons in Magnetic Fields are Accelerated in Circular Orbits

- $\frac{d\mathbf{p}}{dt} = e (\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B})$
- Electrons follow a circular orbit with radii  $\rho$  proportional to their momentum,  $\rho = \frac{pc}{eB}$



# Lorentz Law Yields Kinetic Energy vs. Magnetic Field Relations

Classically,


$$K = \frac{e^2 \rho^2}{2mc^2} B^2$$

Relativistically,


$$K = \sqrt{e^2 \rho^2 B^2 + m^2 c^4} - mc^2$$

Experimentally measuring  $K$  vs.  $B$  will let us pick the more likely model


# Experimental Setup Constrains Radius of Electron Orbit




# Experimental Setup Constrains Radius of Electron Orbit



# Experimental Setup Constrains Radius of Electron Orbit




# Barium-133 Produces MCA Peaks at Known Energies







# MCA Readout for Sr-90/Y-90 Sharply Peaked around Energy Range



# Magnetic Field Affects Peak Energy Range



# Kinetic Energy Determined through Gaussian Fitting



# Gaussian Fits Bring Uncertainty in Kinetic Energy

$B_{approx}$ (G)	K (keV)	$\sigma_K$ (keV)
90	222	1.47
100	265	1.08
110	312	.79
120	355	.66

## Uncertainties in Magnetic Field

## Uncertainties in Magnetic Field

- Variations during individual runs from coil heating

## Uncertainties in Magnetic Field

- Variations during individual runs from coil heating
- Variations between runs

## Uncertainties in Magnetic Field

- Variations during individual runs from coil heating
- Variations between runs
- Inhomogeneous magnetic field during individual runs




## Uncertainties in Magnetic Field


- Variations during individual runs from coil heating
- Variations between runs
- Inhomogeneous magnetic field during individual runs
- Systematic uncertainty in magnetometer

# Inhomogeneity Addressed by Averaging over Multiple Points

- Measured at point C during experimental runs
- Determined correspondance between magnetic field at point C and the average magnetic field over the path of the electron



# Data Follows Relativistic Trend




# Relativistic Fit Predicts Realistic Constants

	Measured Values	Expected Values
$e$ (esu)	$5.02 \pm .03$	4.8
$mc^2$ (keV)	$505 \pm 76$	511

Correlated fit pays off - uncertainty in  $mc^2$  reduces from 200 keV to 76 keV.

# Rest Energy Is Not Dominant In Our Regime

$$K = \sqrt{\rho^2 e^2 B^2 - m^2 c^4} - mc^2$$



# Electromagnetism Allows for Detection of Relativistic Effects

# Electromagnetism Allows for Detection of Relativistic Effects

- Special Relativity and Classical Mechanics make different predictions for  $K(p)$

# Electromagnetism Allows for Detection of Relativistic Effects

- Special Relativity and Classical Mechanics make different predictions for  $K(p)$
- Using Lorentz's Law, we can determine electron momentum by the magnetic field supplied



# Electromagnetism Allows for Detection of Relativistic Effects

- Special Relativity and Classical Mechanics make different predictions for  $K(p)$
- Using Lorentz's Law, we can determine electron momentum by the magnetic field supplied
- Experimentally determined relationship of  $K$  and  $B$  follow relativistic trends and predict electron charge and rest energy within expected values

# Relativistic Energy Reduces to Classical when $v \ll c$

$$K = \sqrt{p^2 c^2 + m^2 c^4} - mc^2$$

# Relativistic Energy Reduces to Classical when $v \ll c$

$$\begin{aligned} K &= \sqrt{p^2 c^2 + m^2 c^4} - mc^2 \\ &= \sqrt{m^2 c^4 \left( 1 + \frac{p^2}{m^2 c^2} \right)} - mc^2 \end{aligned}$$

# Relativistic Energy Reduces to Classical when $v \ll c$

$$\begin{aligned} K &= \sqrt{p^2 c^2 + m^2 c^4} - mc^2 \\ &= \sqrt{m^2 c^4 \left( 1 + \frac{p^2}{m^2 c^2} \right)} - mc^2 \\ &= mc^2 \sqrt{1 + \frac{p^2}{m^2 c^2}} - mc^2 \end{aligned}$$

# Relativistic Energy Reduces to Classical when $v \ll c$

$$\begin{aligned} K &= \sqrt{p^2 c^2 + m^2 c^4} - mc^2 \\ &= \sqrt{m^2 c^4 \left( 1 + \frac{p^2}{m^2 c^2} \right)} - mc^2 \\ &= mc^2 \sqrt{1 + \frac{p^2}{m^2 c^2}} - mc^2 \\ &\approx mc^2 \left( 1 + \frac{p^2}{2m^2 c^2} \right) - mc^2 \end{aligned}$$

# Relativistic Energy Reduces to Classical when $v \ll c$

$$\begin{aligned} K &= \sqrt{p^2 c^2 + m^2 c^4} - mc^2 \\ &= \sqrt{m^2 c^4 \left( 1 + \frac{p^2}{m^2 c^2} \right)} - mc^2 \\ &= mc^2 \sqrt{1 + \frac{p^2}{m^2 c^2}} - mc^2 \\ &\approx mc^2 \left( 1 + \frac{p^2}{2m^2 c^2} \right) - mc^2 \\ &= \frac{p^2}{2m} \end{aligned}$$

# Voltage for Curvature Correction Dependent on Velocity

- For detection, the trajectory of the electron must be adjusted to a straight line

# Voltage for Curvature Correction Dependent on Velocity

- For detection, the trajectory of the electron must be adjusted to a straight line
- From Lorentz's Law, the required electric field is  $E = \frac{vB}{c}$



# Voltage for Curvature Correction Dependent on Velocity

- For detection, the trajectory of the electron must be adjusted to a straight line
- From Lorentz's Law, the required electric field is  $E = \frac{vB}{c}$
- For two plates separated by a distance  $d$ , the voltage required is  $V = \frac{dvB}{c}$

# Calibration Fit for Magnetometer Results in Large Errors

