


Jointcal: Optimized Astrometry & Photometry for Thousands of Exposures with Large Mosaic Cameras

Introduction and Algorithm

- Jointcal is a new software package developed for the LSST Science Pipelines to optimize the astrometric and photometric calibrations of a set of mosaic camera images of an area of sky.
- Jointcal can operate on cameras supported by the LSST software, including Canada-France-Hawaii Telescope (CFHT) Megacam, Dark Energy Camera (DECam), Subaru Hyper SuprimeCam (HSC), and simulated LSST images from phosim and imsim.
- The jointcal algorithm matches both between exposures and to external reference catalogs; this allows many more sources to be fit when the images are significantly deeper than those of the reference catalogs (for example, LSST's 30-second r-band exposure depth will be 24.7 compared with $r < 20$ for Gaia's final survey depth).



$$\chi^2 = \sum_{\gamma,i} [M_\gamma(S_{\gamma,i}) - F_i]^T W_{\gamma,i} [M_\gamma(S_{\gamma,i}) - F_i] \quad (\text{meas. terms})$$

$$+ \sum_j [F_j - R_j]^T W_j [F_j - R_j] \quad (\text{ref. terms})$$



- Jointcal minimizes the χ^2 equation shown above, jointly fitting the sensor-to-sky model and the "true"-to-reference catalog value.
- The 45 exposures in this example include >500,000 stars + >3500 model parameters.
- Jointcal uses the sparseness of the Hessian matrix (2nd derivatives of the model parameters) of χ^2 to rapidly solve the large linear system.
- Running jointcal on 45 visits of HSC takes about 5 hours (single core).
- Plan to parallelize reading data and outlier rejection.



Check out jointcal on GitHub:
github.com/lsst/jointcal
 More info: dmtn-036.lsst.io
 Contact me: parejkoj@uw.edu


Astrometric Calibration

- Here we demonstrate jointcal's astrometric and photometric calibration performance on HSC data (101 CCDs), with 45 exposures of one field (XMM-WIDE HSC-I band).
- We fit a 7th order polynomial per exposure over the whole focal plane, and an affine transform (x/y rotation, translation, and scale) for each sensor (fixed for all observations).
- Residuals are markedly reduced on the edges of CCDs. More sources and a polynomial that continues beyond the edge help the model to not diverge compared with fitting CCDs individually.




Photometric Calibration

- Here is jointcal's photometric calibration of one HSC exposure (one scale factor per CCD, times a 7th order chebyshev polynomial per visit), with the single frame algorithm (`processCcd`) divided out.
- The largest changes are at the edges of the focal plane, where HSC is significantly vigneted.
- The middle of the focal plane has improvements at the >2% level.



- Both panels below show the RMS of the residuals on bright stars that are measured on multiple exposures.
- The dashed lines are the median RMS; jointcal performs markedly better in both astrometry and photometry.



John K. Parejko: University of Washington, DIRAC, LSST
 Pierre Astier: LPNHE/IN2P3/CNRS, Paris, France
 Jim Bosch: Princeton University, LSST
 Dominique Boutigny: LPNHE/IN2P3/CNRS, Paris, France



This material is based upon work supported in part by the National Science Foundation through Cooperative Support Agreement (CSA) Award No. AST-1227061 under Governing Cooperative Agreement 1258333 managed by the Association of Universities for Research in Astronomy (AURA), and the Department of Energy under Contract No. DE-AC02-76SF00515 with the SLAC National Accelerator Laboratory. Additional LSST funding comes from private donations, grants to universities, and in-kind support from LSSTC Institutional Members.

