

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.elsevier.com/locate/jval](http://www.elsevier.com/locate/jval)

## PREFERENCE-BASED ASSESSMENTS

## Associations of Smoking, Physical Inactivity, Heavy Drinking, and Obesity with Quality-Adjusted Life Expectancy among US Adults with Depression

Haomiao Jia, PhD<sup>1,\*</sup>, Matthew M. Zack, MD, MPH<sup>2</sup>, Irving I. Gottesman, PhD<sup>3</sup>, William W. Thompson, PhD<sup>4</sup><sup>1</sup>Department of Biostatistics, Mailman School of Public Health and School of Nursing, Columbia University, New York, NY, USA;<sup>2</sup>National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, GA, USA; <sup>3</sup>Departments of Psychology and Psychiatry, University of Minnesota, Twin Cities, MN, USA; <sup>4</sup>National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention, Atlanta, GA, USA

## A B S T R A C T

**Objectives:** To examine associations between four health behaviors (smoking, physical inactivity, heavy alcohol drinking, and obesity) and three health indices (health-related quality of life, life expectancy, and quality-adjusted life expectancy (QALE)) among US adults with depression. **Methods:** Data were obtained from the 2006, 2008, and 2010 Behavioral Risk Factor Surveillance System data. The EuroQol five-dimensional questionnaire (EQ-5D) health preference scores were estimated on the basis of extrapolations from the Centers for Disease Control and Prevention's healthy days measures. Depression scores were estimated using the eight-item Patient Health Questionnaire. Life expectancy estimates were obtained from US life tables, and QALE was estimated from a weighted combination of the EQ-5D scores and the life expectancy estimates. Outcomes were summarized by depression status for the four health behaviors (smoking, physical inactivity, heavy alcohol drinking, and obesity). **Results:** For depressed adults, current smokers and the physically inactive had significantly lower EQ-5D scores (0.040 and 0.171, respectively), shorter life expectancy

(12.9 and 10.8 years, respectively), and substantially less QALE (8.6 and 10.9 years, respectively). For nondepressed adults, estimated effects were similar but smaller. Heavy alcohol drinking among depressed adults, paradoxically, was associated with higher EQ-5D scores but shorter life expectancy. Obesity was strongly associated with lower EQ-5D scores but only weakly associated with shorter life expectancy. **Conclusions:** Among depressed adults, physical inactivity and smoking were strongly associated with lower EQ-5D scores, life expectancy, and QALE, whereas obesity and heavy drinking were only weakly associated with these indices. These results suggest that reducing physical inactivity and smoking would improve health more among depressed adults.

**Keywords:** burden of illness, depression, quality-adjusted life-years, risk factors.

© 2018 Published by Elsevier Inc. on behalf of International Society for Pharmacoeconomics and Outcomes Research (ISPOR).

## Introduction

Depression is a health outcome associated with genetic predispositions, sex, adverse childhood experiences, exposure to disasters, and poverty [1–5]. It is also associated with several unhealthy behaviors including physical inactivity [6–10], sedentary behavior [11–13], excessive alcohol use [14–17], smoking [18–22], and less social engagement [23]. Finally, depression is associated with most chronic conditions [24]. In a review of 31 studies of associations between depression and other chronic

conditions including diabetes, pulmonary disease, heart disease, and arthritis, those with a chronic condition as well as depression reported significantly more medical symptoms than did those with only a chronic condition [25]. In terms of causal associations, the relationship between depression and other chronic conditions is bidirectional [26,27].

Although depression has an age of onset earlier than that of most other chronic conditions [28,29], this does not necessarily demonstrate that depression causes the new chronic conditions. In a review examining bidirectional effects of depression and chronic

Conflicts of interest: The authors have no conflicts of interest to report.

This analysis used de-identified data produced by federal agencies in the public domain. Data were downloaded from the Centers for Disease Control and Prevention Web site ([ftp://ftp.cdc.gov/pub](http://ftp.cdc.gov/pub)). The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

\* Address correspondence to: Haomiao Jia, Department of Biostatistics, Mailman School of Public Health and School of Nursing, Columbia University, 560 West 168th Street, New York, NY 10032.

E-mail: [hj2198@columbia.edu](mailto:hj2198@columbia.edu)

1098-3015/\$36.00 – see front matter © 2018 Published by Elsevier Inc. on behalf of International Society for Pharmacoeconomics and Outcomes Research (ISPOR).

<http://dx.doi.org/10.1016/j.jval.2017.08.002>

obstructive pulmonary disease (COPD) [30], 16 studies supported a causal effect of depression on COPD, whereas 6 studies supported a causal effect of COPD on depression.

Several different interventions decrease rates of depression, including drug therapy [31], psychotherapy [32,33], physical activity [7–10], and reducing harmful risky behaviors such as smoking and heavy alcohol use [21,22,34]. Counterintuitively, in some studies, smoking cessation has been found to increase rates of depressive symptoms in some individuals, likely because of withdrawal symptoms associated with nicotine dependence.

Depression is a leading cause of disability [24,35–37] and suicide [35,38,39]. It is also associated with poor health-related quality of life (HRQOL) and increased years of living with a disability [35,40,41]. Depression has also been associated with excess mortality and substantially lower life expectancy [42–44]. In a recent study of US adults, the life expectancy for those with depression was 16.4 years less than that for those without depression [44]. The total health loss, for both fatal and nonfatal outcomes associated with depression, was a 28.9-year loss in quality-adjusted life expectancy (QALE) starting at age 18 years, a more than 50% decrease in QALE relative to that in those without depression.

QALE takes into account both the years of life lost and the relative severity of current chronic conditions, making it possible to quantify both population-level and individual-level health losses associated with both fatal and nonfatal outcomes [45–48]. Typically, preference-based measures of HRQOL are used to estimate the effects of these nonfatal health losses. These measures capture respondents' perceived health for different health states using a summary score (called utility value) between 0 (for death) and 1 (for perfect health) [45]. Thus, 1 year of life lived at a utility value of 0.5 equals 0.5 quality-adjusted life-years (QALYs), the same as that of only half a year of life lived at a perfect utility value of 1.0 [24,25]. The QALE at a certain age (such as at age 18 years) is defined as the average QALY throughout the remainder of expected life [44,47].

The primary objectives of this article were to estimate changes in age-adjusted HRQOL scores, life expectancy, and QALE associated with four risk factors—physical inactivity, smoking, heavy drinking, and obesity—previously associated with depression among adults. This study also estimated the potential gains in QALE for adults with high levels of depressive symptoms if they had lowered the frequency and levels of these risk factors.

## Methods

The Behavioral Risk Factor Surveillance System (BRFSS) is an ongoing annual survey of noninstitutionalized civilian adult residents from each of the 50 states, the District of Columbia, and participating territories [49–51]. The BRFSS uses a random-digit-dialed telephone survey method to ascertain a population-based random sample of adults aged 18 years and older [50,51].

During 2006, 2008, and 2010, the eight-item Patient Health Questionnaire (PHQ-8) depression scale was included in the BRFSS state optional modules to classify individuals according to the severity of their self-reported depression symptoms [52,53]. The PHQ-8 is established as a valid and reliable diagnostic measure of depressive symptoms in large clinical studies and has frequently been used to estimate the prevalence of depression in the US adult population [54–56]. Thirty-six states and the District of Columbia asked PHQ-8 questions at least once during these 3 years, yielding a total sample size of 276,442 for this analysis. For the purposes of this study, a PHQ-8 score of 10 or more will be used to categorize adults as depressed [52,53] because such a score has an 88% sensitivity and an 88% specificity for major depressive disorders [54,55].

The BRFSS asks respondents about their physical activity, cigarette smoking, number of drinks of alcohol, and their weight and height without their shoes. Leisure-time physical activity is defined as having performed, during the past 30 days, physical activity or exercise other than that in a regular job. Cigarette smoking is defined as currently smoking any cigarettes every day

**Table 1 – Demographic and risk factor characteristics among US adults with and without depression—BRFSS 2006, 2008, and 2010.**

Demographic and risk factor characteristic	Depression (n = 24,826)		No depression (n = 251,616)	
	% <sup>*</sup>	SE	% <sup>*</sup>	SE
Age (y), mean	43.1	0.2	45.9	0.1
Age groups (y)				
18–44	54.1%	0.7%	50.7%	0.2%
45–64	36.6%	0.7%	32.7%	0.2%
> 65	9.3%	0.3%	16.6%	0.1%
% Women	61.0%	0.3%	49.3%	1.3%
Race/ethnicity				
White non-Hispanic	68.3%	1.8%	72.1%	1.6%
Black non-Hispanic	13.8%	1.0%	10.4%	0.9%
Hispanic	12.7%	1.7%	12.0%	1.5%
Others	5.2%	0.8%	5.5%	0.8%
% Currently smoking	39.9%	1.4%	19.5%	1.2%
% Physically inactive	43.5%	1.5%	20.6%	1.4%
% Heavy alcohol drinking	7.4%	0.8%	5.1%	0.7%
Body mass index (kg/m <sup>2</sup> ) groups				
< 18.5	1.9%	0.6%	0.8%	0.6%
18.5–24.9	23.2%	1.7%	30.0%	1.6%
25.0–29.9	32.1%	1.6%	38.7%	1.5%
≥ 30.0	42.8%	1.3%	30.5%	1.2%

BRFSS, Behavioral Risk Factor Surveillance System; SE, standard error.

<sup>\*</sup> Percentages accounting for sampling weight and age in subgroups (except by age groups).

or some days. Heavy alcohol drinking is defined as drinking alcoholic beverages 15 or more times per week for men and 8 or more times per week for women. Obesity is defined as having a body mass index (BMI; weight [kg] divided by height [m] squared) greater than or equal to 30 kg/m<sup>2</sup>. Only 1.9% of the depressed US adults were underweight (BMI < 18.5 kg/m<sup>2</sup>) (Table 1), and so we did not calculate QALE for those who were underweight because the mortality estimates in such a small group were considered unreliable.

The BRFSS includes four HRQOL questions that ask respondents to rank their general health from 1 (excellent) to 5 (poor) and to report their number of physically and mentally unhealthy days and days with activity limitation during the past 30 days [57]. Because these questions are not preference-based measures of HRQOL, this study applied a previously constructed algorithm to obtain values for the EuroQol five-dimensional questionnaire (EQ-5D) index, a preference-based HRQOL measure, for BRFSS respondents, on the basis of their age and answers to these four questions [58,59]. This algorithm provides valid estimates of the EQ-5D scores of the US population overall and in some demographic subgroups and common health conditions identified from the BRFSS [58,59]. The bias of estimated QALE from these scores has been estimated to be less than 1% of that using the actual EQ-5D questions [60].

Life expectancy at a given age is the average number of years of life remaining starting at that age and is calculated using the life table method [61]. The National Center for Health Statistics

compiles death data for the US population and makes these data available to the public in the Compressed Mortality File at <http://wonder.cdc.gov>. This file includes age-specific mortality rates and can be used to construct life tables for the US population [60,61].

The age-specific death rate ( $m$ ) was obtained by dividing the number of deaths by the population size. Because age-specific death rates stratified by depression and each of the four health behaviors are not available, these rates were estimated. Death rates for those with depression ( $m_1$ ) and those without depression ( $m_0$ ) were estimated using the hazard ratio ( $h$ ) of dying for depressed versus nondepressed and the prevalence of depression ( $p$ ) by  $m_1 = \frac{hm}{hp + (1-p)}$  and  $m_0 = \frac{m}{hp + (1-p)}$ , respectively [47]. The hazard ratio was estimated from the 2004 to 2010 National Health and Nutrition Examination Survey (NHANES) data as linked to the National Death Index through December 31, 2011 (<https://www.cdc.gov/nchs/data-linkage/mortality.htm>), using the Cox proportional hazards model. The prevalence of depression was estimated from the BRFSS.

The QALE was calculated using the life table method [60]. Let  $l_i$  be the number of a hypothetical population surviving to the beginning of age interval  $i$  (18–24, 25–34, ..., > 85 years) and  $L_i$  be the total life-years for the age interval  $i$ . Assuming a constant death rate during an age interval, the probability of dying in an  $n_i$ -year age interval is  $q_i = 1 - e^{-n_i m_i}$  and those who died during the interval for ages  $x < 85$  years lived approximately  $n_i/2$  years, that

**Table 2 – Association between selected risk factors and the EQ-5D index, life expectancy, and QALE by depression status among US adults—BRFSS 2006, 2008, and 2010<sup>\*</sup>.**

Risk factor	EQ-5D <sup>†</sup>		Life expectancy <sup>‡</sup>		QALE <sup>§</sup>	
	depression		depression		depression	
	Yes	No	Yes	No	Yes	No
All	0.594	0.909	47.3	63.7	28.0	56.8
Smoking						
No	0.610	0.914	53.2	65.6	31.9	58.7
Yes	0.570	0.888	40.3	54.1	23.3	47.8
Loss	0.040	0.025	12.9	11.4	8.6	10.8
% Loss	6.6	2.8	24.3	17.4	27.0	8.4
Physical inactivity						
No	0.669	0.921	53.6	64.3	34.0	58.5
Yes	0.498	0.864	42.8	62.3	23.1	52.7
Loss	0.171	0.057	10.8	2.1	10.9	5.8
% Loss	25.6	6.2	20.1	3.2	32.0	9.9
Heavy drinking						
No	0.568	0.909	47.7	63.9	28.0	57.1
Yes	0.671	0.909	41.3	57.2	27.9	52.0
Loss	−0.103	0.000	6.4	6.7	0.1	5.1
% Loss	−18.1	0.0	13.4	10.5	0.4 <sup>*</sup>	8.9
Obesity <sup>  </sup>						
No	0.616	0.918	47.4	64.0	28.7	57.6
Yes	0.563	0.889	47.0	63.5	27.0	55.1
Loss	0.053	0.029	0.4	0.5	1.7	2.5
% Loss	8.6	3.1	0.9	0.8	5.8	4.4

BMI, body mass index; BRFSS, Behavioral Risk Factor Surveillance System; EQ-5D, EuroQol five-dimensional questionnaire; QALE, quality-adjusted life expectancy.

\* All estimates in this table are statistically significant except for the QALE loss for the depressed individuals who reported they engaged in heavy drinking.

<sup>†</sup> Age-adjusted EQ-5D index.

<sup>‡</sup> Life expectancy at age 18 y.

<sup>§</sup> QALE at age 18 y.

<sup>||</sup> Obesity: Yes, BMI ≥ 30.0; No, BMI = 18.5–29.9; excluded, BMI < 18.5.

is,  $L_i = l_i(1 - n_i m_i / 2)$ , and for the last age interval ( $>85$  years), the average years of life at age 85 years is  $L_{85} = l_{85} / m_{85}$  [60,61]. The life expectancy at age  $x$  is  $\sum_{k=x}^{\infty} L_k$ . If  $y_i$  is the mean HRQOL score, the number of QALYs for this age interval is  $L_i y_i$ . The QALE at age  $x$  is calculated by summarizing QALYs throughout the remaining expected life starting at age  $x$  divided by the population surviving to age  $x$  and is calculated as  $\frac{\sum_{k=x}^{\infty} L_k y_k}{l_x}$ .

We estimate life expectancy and QALE by respondents' depression status cross-tabulated by each of the four risk factors. Life expectancy loss associated with a risk factor is the difference in life expectancy between those with and without the risk factor, and QALE loss associated with a risk factor is the difference in QALE between these two groups [47,62]. The nonfatal health loss due to a risk factor is the difference in age-standardized EQ-5D scores (standardized to the year 2010 US population) between those with the factor and those without the factor [44].

## Results


The total sample size was 276,442. Approximately 9.1% (standard error = 0.13%) of adults in this study using our case definition were currently depressed. The depressed adults were about 2.8 years younger than the nondepressed adults (Table 1). After age adjustment, those who were depressed were more likely to be women (61.0% vs. 49.3%) and black non-Hispanics (13.8% vs. 10.4%) and less likely to be white non-Hispanics (68.3% vs. 71.1%) than those who were not depressed. Those who were depressed had about the same proportions of Hispanics (12.7% vs. 12.0%) and other races (5.2% vs. 5.5%) as those who were not depressed. Compared with those who were not depressed, those who were depressed were more than twice as likely to be current

smokers (39.9% vs. 19.5%) and to be physically inactive (43.5% vs. 20.6%) and more than 40% as likely to be heavy drinkers (7.4% vs. 5.1%) and to be obese (42.8% vs. 30.5%).

The age-adjusted EQ-5D index was 0.594 for those who were depressed and 0.909 for those who were not depressed (Table 2). The life expectancy for individuals aged 18 years was 47.3 additional years for those who were depressed and 63.7 additional years for those who were not depressed. The QALE at age 18 years was 28.0 years for those who were depressed and 56.9 years for those who were not depressed. This 28.9-year decrease represents the years of QALE loss associated with depression (or a loss of 50.8% QALE) starting at age 18 years. Because the main aim of this study was to examine the association of each of the four risk factors with HRQOL, life expectancy, and QALE, for those with depression, the rest of this study pertains only to US adults with depression.

## Smoking among Depressed Adults

Nearly 40% of depressed adults were current smokers (Table 1). Generally, smoking had a relatively small adverse relationship with HRQOL but a large adverse relationship with life expectancy among depressed adults (Table 2). The age-adjusted EQ-5D index for depressed current smokers was 0.570, 0.040 points lower (or 6.6% lower) than that for depressed nonsmokers (0.610). The life expectancy for an 18-year-old smoker was 40.3 years, about 12.9 years less (24.3%) than the life expectancy of an 18-year-old nonsmoker (53.2 years). This 12.9-year decrease in life expectancy was the largest among the four risk factors examined, and the 0.040-point decrease in the EQ-5D index was the smallest. Combining the small loss in the EQ-5D index with the large loss in life expectancy led to a relatively large QALE decrease



**Fig. 1 – Decreases in QALE associated with selected risk factors among US adults with depression at different ages—BRFSS 2006, 2008, and 2010.** BRFSS, Behavioral Risk Factor Surveillance System; CI, confidence interval; QALE, quality-adjusted life expectancy.

associated with smoking. Because the QALE for smokers was 23.3 years and for nonsmokers was 31.9 years, smoking was associated with an 8.6-year QALE decrease (or a decrease of 27% QALE), the second largest decrease among the four risk factors.

The decreases in QALE associated with smoking were statistically significant at all ages (Fig. 1). For example, an 85-year-old depressed person had a significantly lower QALE (0.2 years) than an 85-year-old person who was not depressed (0.9 years). The decrease in QALE declined gradually with age, going from 8.6 years at age 18 years to 0.7 years at age 85 years. This occurs because QALE declines with age, and not because smoking has a weaker association with QALE at older ages. Such a large and significant QALE decrease associated with smoking occurred among both men and women and among white non-Hispanics, black non-Hispanics, and Hispanics (Fig. 2). Smoking was associated with approximately the same amount of QALE decrease for both men and women and for the three race/ethnicity subgroups (between 7.9 and 8.6 years).

Depressed adults had 28.9 years of QALE less than nondepressed adults; part of this difference may be related to higher smoking prevalence rates among depressed adults (39.9%) relative to those in adults who were not depressed (19.5%). If depressed adults lowered their smoking prevalence to that of individuals who were not depressed, it could potentially increase their QALE by 1.9 years (6.8%), from the current estimate of 28.0 years to 29.9 years.

### Physical Inactivity among Depressed Adults

Approximately 43.5% of depressed adults were physically inactive (Table 1). Their age-adjusted EQ-5D index was 0.498, 0.171 points (25.6%) lower than that for those who were physically active (0.669) (Table 2). This 0.171-point decrease in the EQ-5D index was


the largest among the four risk factors examined. The life expectancy for an 18-year-old inactive person was 42.8 years, 10.8 years (20.1%) less than the life expectancy of an 18-year-old active person (53.6 years). This 10.8-year decrease in life expectancy was the second largest of the four risk factors. The strong association between physical inactivity and both EQ-5D and life expectancy results in a very large QALE decrease associated with physical inactivity. Being physically inactive was associated with a 10.9-year (32%) decrease in QALE (34.0 years for the active group and 23.1 years for the inactive group). This 10.9-year decrease in QALE was also the largest among the four risk factors.

The decreases in QALE associated with physical inactivity were also significant at all ages (Fig. 1). Like the decrease associated with smoking, the QALE loss to physical inactivity also declined gradually with age, going from 10.9 years at age 18 years to 1.6 years at age 85 years. Such a strong association of physical inactivity with QALE was observed for both men (an 11.8-year decrease) and women (a 9.9-year decrease) and for all three race/ethnicity groups (a 16.6-year decrease among Hispanics, a 11.5-year decrease among white non-Hispanics, and a 10.9-year decrease among black non-Hispanics; Fig. 2). Nevertheless, these decreases among the different demographic subgroups did not differ statistically significantly.

If the prevalence of physical inactivity among depressed adults (43.5%) was reduced to the same prevalence as that among adults who were not depressed (20.6%), the depressed adults could potentially increase their QALE by 2.8 years (10%) to 30.8 years.

### Heavy Alcohol Drinking among Depressed Adults

Among those with depression, about 7.4% were heavy drinkers (Table 1). Among depressed adults, heavy drinkers had a



**Fig. 2 – Decreases in QALE associated with selected risk factors among US adults with depression, overall and for men, women, white non-Hispanics, black non-Hispanics, and Hispanics—BRFSS 2006, 2008, and 2010.** BRFSS, Behavioral Risk Factor Surveillance System; CI, confidence interval; QALE, quality-adjusted life expectancy.



significantly higher EQ-5D index (0.671) compared with non-heavy drinkers (0.568) (Table 2). At age 18 years, heavy drinkers had a significantly lower life expectancy (41.3 years) compared with non-heavy drinkers (47.7 years). On combining a higher EQ-5D index with a lower life expectancy, we found that heavy drinkers had nearly the same QALE as non-heavy drinkers. The QALE was 27.9 years for heavy drinkers and 28.0 years for non-heavy drinkers. This 0.1-year difference was the smallest among the four risk factors examined and was not statistically significant (95% confidence interval [CI] –2.5 to 2.7).

When examined by sex and race/ethnicity, the difference in QALE associated with heavy drinking was not statistically significant for men (95% CI –2.9 to 2.7), women (95% CI –3.6 to 2.2), white non-Hispanics (95% CI –3.4 to 2.0), and black non-Hispanics (95% CI –0.2 to 8.4). Only Hispanics with depression had statistically significantly less QALE associated with heavy drinking (4.5 years [95% CI 0.2 to 8.8]).

### Obesity among Depressed Adults

We did not find a significant and meaningful difference in the EQ-5D index and death rates between normal-weight adults (BMI = 18.5–24.9 kg/m<sup>2</sup>) and overweight adults (BMI = 25.0–29.9 kg/m<sup>2</sup>); therefore, we combined normal-weight and overweight adults into one group (nonobese adults [BMI = 18.5–29.9 kg/m<sup>2</sup>]) and compared them with obese adults (BMI ≥ 30 kg/m<sup>2</sup>). Among those with depression, the age-adjusted EQ-5D index was 0.563 for obese adults and 0.616 for nonobese adults (Table 2). This 0.053-point decrease in EQ-5D associated with obesity exceeded the 0.040-point decrease associated with smoking but was smaller than the 0.171-point decrease associated with physical inactivity and the 0.103-point increase associated with heavy drinking. The life expectancy of an 18-year-old obese person was 47.0 years, about 0.4 years less than the life expectancy of an 18-year-old nonobese person (47.4 years). This 0.4-year decrease in life expectancy was the smallest among the four risk factors but was statistically significant ( $P = 0.04$ ). Overall, obesity was associated with a statistically significant 1.7-year (5.8%) QALE decrease (27.0 years for obese persons and 28.7 years for non-obese persons).

Being obese was associated with a statistically significant 1.8-year (95% CI 0.7 to 2.9) QALE decrease for depressed women (Fig. 2). For depressed men, the obesity-associated QALE decrease was not statistically significant (1.1-year decrease; 95% CI –0.2 to 2.4). The decrease in QALE associated with obesity was significant only among those aged 55 years or younger. In addition, the QALE decrease associated with obesity declined gradually down to 0 between age 18 years and 55 years (Fig. 1). This suggests that obesity is significantly associated with health only among younger depressed adults.

If the prevalence of obesity among depressed adults was reduced to a rate such as that in adults who were not depressed, the depressed adults could potentially increase their QALE by 0.2 years to 28.2 years.

## Discussion

In our study, leisure-time physical inactivity and smoking were strongly associated with two health outcomes and one composite score (i.e., HRQOL, life expectancy, and QALE) among US adults with depression. The other two risk factors, heavy alcohol drinking and obesity, surprisingly, had nonstatistically significant or negligible associations with respondents' QALE (0.1-year decrease for heavy drinkers and 1.7-year decrease for obese persons). These findings were generally consistent at all ages,

for both men and women, and for white non-Hispanics, black non-Hispanics, and Hispanics.

Each of the four risk factors had differences in decreases for fatal and nonfatal outcomes. Smoking was associated with a large reduction in life expectancy (a loss of 12.9 years of life) but a relatively small decrease in HRQOL scores (a decrease of 0.04 points in the EQ-5D index) among depressed adults. This is consistent with findings from previous studies of the impact of smoking on general populations [47,63–66]. For example, smoking is one of the leading causes of death among US adults [64], but in a study of 13,000 adults, current smoking was associated with a statistically significant but only a small decrease of 0.03 points in the EQ-5D index [66].

Physical inactivity also had a large association with both HRQOL and life expectancy—a decrease of 20.1% life expectancy and 25.6% HRQOL. This is likely because physical inactivity is a strong predictor of long-term mortality [41] and because physical inactivity is also associated with physical limitations [67,68]. Therefore, it would be expected that the EQ-5D and physical inactivity would be correlated.

Although excessive alcohol consumption has been associated with deteriorating HRQOL in the general population [69], this study found that among those with depression, heavy drinkers had significantly better age-adjusted HRQOL scores than non-heavy drinkers, with an increase of 18.1% HRQOL. Among depressed people, heavy drinkers rated their general health significantly better than did non-heavy drinkers. For example, 66.4% of heavy drinkers rated their general health as “excellent,” “very good,” or “good,” significantly higher than the 52.6% of non-heavy drinkers. Therefore, among those with depression, heavy drinkers perceived themselves to be healthier than those who were not heavy drinkers. Nevertheless, in this study, heavy drinkers with depression had a significantly lower life expectancy, a finding consistent with previous reports that excessive alcohol consumption contributed significant loss in premature death and years of life lost [65,68]. This suggests that the depressed heavy drinkers may be misperceiving their health relative to those who are not heavy drinkers.

Although obesity was associated with an 8.6% decrease in HRQOL, the life expectancy of obese persons in this study was nearly the same as the life expectancy of normal-weight or overweight persons for those with depression. Obesity was associated with only a 0.4-year (0.8%) decrease in life expectancy, the smallest of the four risk factors [70].

Ideally, QALE should be estimated from a longitudinal study in which HRQOL and mortality would be estimated simultaneously. But, in practice, such longitudinal data are usually not available or too sparse to provide reliable estimates. This study applied a method that estimated the mean HRQOL (from table BRFSS) and the life-years (from Compressed Mortality File, NHANES, and BRFSS) separately from different data sets. One of the primary advantages of this method is its ability to use currently available data to estimate QALE for those with many chronic conditions and/or risky behaviors.

### Study Limitations

First, the PHQ-8 is not a clinical diagnostic tool for diagnosing major depressive disorder but has been used primarily for estimating the prevalence of depressive symptoms in the general population; nevertheless, misclassification of the occurrence of depression would likely attenuate the associations we report. Second, we estimated HRQOL and depression using cross-sectional point estimates. Current symptoms of depression are correlated with chronic depression, but depression is a state-dependent health outcome and not all individuals with current depressive symptoms will go on to experience chronic

depression. Third, this study relies on estimates of preference-based HRQOL scores indirectly using the Centers for Disease Control and Prevention's unhealthy days questions rather than direct measurements using a scale such as the EQ-5D. Therefore, our estimates of QALE loss are likely to be attenuated because of "regression to the mean" [47]. One study that estimated the bias of the estimated QALE using the estimated EQ-5D scores found a bias less than 1% of that using the actual EQ-5D questions [60]. Fourth, the lack of information about, and the inability to adjust for, potential unmeasured confounders in these data may have affected our estimates of QALE loss especially for heavy alcohol drinkers. Fifth, the NHANES questions used to ascertain physical activity differed in wording but not in time period and setting from the BRFSS questions used to classify persons as physically inactive in this study. Nevertheless, because the prevalence of physical inactivity in those who were depressed markedly exceeded that in those who were not depressed, we doubt that these differences in wording led to misclassification large enough to affect our estimates of decrease in QALE in those who were both depressed and physically inactive. Sixth, we likely underestimated the decrease in QALE associated with cigarette smoking because only current smokers were associated with these decreases because we classified former smokers as nonsmokers in these analyses. Finally, QALE was estimated for each individual risk behavior independent of the other risk behaviors. Because these behaviors are correlated, our estimates may not accurately reflect the potential additive or interactive effects of co-occurring behaviors or any nonlinear effects between depression symptoms and these behaviors.

## Conclusions

Physical inactivity and smoking had stronger associations with QALE among adults with depression than among adults without depression. These relationships were substantially weaker for obesity and heavy alcohol drinking. These findings have at least two important public health implications. First, these findings support using QALE to describe the burden of disease in groups with risky behaviors. Second, these findings suggest the importance of carrying out additional studies of the complex associations between depression and risky behaviors such as alcohol abuse and smoking [71–75].

Source of financial support: H.J. received support from a contract of the Centers for Disease Control and Prevention (No. 200-2011-M-41977). I. I.G. received support from the University of Louisville Grawemeyer Award for Psychology.

## REFERENCES

- [1] Flint J, Kendler KS. The genetics of major depression. *Neuron* 2014;81:484–503.
- [2] Sullivan PF, Neale MC, Kendler KS. Genetic epidemiology of major depression: review and meta-analysis. *Am J Psychiatry* 2000;157:1552–62.
- [3] Widom CS, DuMont K, Czaja SJ. A prospective investigation of major depressive disorder and comorbidity in abused and neglected children grown up. *Arch Gen Psychiatry* 2007;64:49–56.
- [4] Cameron R, Watson P, Friedman M. *Interventions Following Mass Violence and Disasters: Strategies for Mental Health Practice*. New York, NY: Guilford Press, 2006.
- [5] Haushofer J, Fehr E. On the psychology of poverty. *Science* 2014;344:862–7.
- [6] Lindwall M, Gerber M, Jonsdottir IH, et al. The relationships of change in physical activity with change in depression, anxiety, and burnout: a longitudinal study of Swedish healthcare workers. *Health Psychol* 2014;33:1309–18.
- [7] Cooney G, Dwan K, Mead G. Exercise for depression. *JAMA* 2014;311:2432–3.
- [8] Stanton R, Reaburn P. Exercise and the treatment of depression: a review of the exercise program variables. *J Sci Med Sport* 2014;17:177–82.
- [9] Josefsson T, Lindwall M, Archer T. Physical exercise intervention in depressive disorders: meta-analysis and systematic review. *Scand J Med Sci Sports* 2014;24:259–72.
- [10] Dunn AL, Trivedi MH, Kampert JB, et al. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med* 2005;28:1–8.
- [11] Wassink-Vossen S, Collard RM, Oude Voshaar RC, et al. Physical (in)activity and depression in older people. *J Affect Disord* 2014;161:65–72.
- [12] de Rezende LF, Rodrigues Lopes M, Rey-López JP, et al. Sedentary behavior and health outcomes: an overview of systematic reviews. *PLoS One* 2014;9:e105620.
- [13] Hamer M, Stamatakis E. Prospective study of sedentary behavior, risk of depression, and cognitive impairment. *Med Sci Sports Exerc* 2014;46:718.
- [14] Regier DA, Farmer ME, Rae DS, et al. Comorbidity of mental disorders with alcohol and other drug abuse: results from the Epidemiologic Catchment Area (ECA) study. *JAMA* 1990;264:2511–8.
- [15] Ross HE, Glaser FB, Germanson T. The prevalence of psychiatric disorders in patients with alcohol and other drug problems. *Arch Gen Psychiatry* 1988;45:1023–31.
- [16] Wu LT, Kouzis AC, Leaf PJ. Influence of comorbid alcohol and psychiatric disorders on utilization of mental health services in the National Comorbidity Survey. *Am J Psychiatry* 1999;156:1230–6.
- [17] Ruggles KV, Fang Y, Tate J, et al. What are the patterns between depression, smoking, unhealthy alcohol use, and other substance use among individuals receiving medical care? A longitudinal study of 5479 participants. *AIDS Behav* 2017;21:2014–22.
- [18] Dierker LC, Avenevoli S, Stolar M, Merikangas KR. Smoking and depression: an examination of mechanisms of comorbidity. *Am J Psychiatry* 2002;159:947–53.
- [19] Glassman AH, Helzer JE, Covey LS, et al. Smoking, smoking cessation, and major depression. *JAMA* 1990;264:1546–9.
- [20] Luger TM, Suls J, Vander Weg MW. How robust is the association between smoking and depression in adults? A meta-analysis using linear mixed-effects models. *Addict Behav* 2014;39:1418–29.
- [21] Taylor G, McNeill A, Girling A, et al. Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ* 2014;348:g1151.
- [22] Shahab L, Gilchrist G, Hagger-Johnson G, et al. Reciprocal associations between smoking cessation and depression in older smokers: findings from the English Longitudinal Study of Ageing. *Br J Psychiatry* 2015;207:243–9.
- [23] Bisschop MJ, Kriegsman DM, Beekman AT, Deeg DJ. Chronic diseases and depression: the modifying role of psychosocial resources. *Soc Sci Med* 2004;59:721–33.
- [24] Chapman DP, Perry GS, Strine TW. The vital link between chronic disease and depressive disorders. *Prev Chronic Dis* 2005;2:A14.
- [25] Katon W, Lin EH, Kroenke K. The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *Gen Hosp Psychiatry* 2007;29:147–55.
- [26] Katon WJ. Clinical and health services relationships between major depression, depressive symptoms, and general medical illness. *Biol Psychiatry* 2003;54:216–26.
- [27] Jonas BS, Mussolino ME. Symptoms of depression as a prospective risk factor for stroke. *Psychosom Med* 2000;62:463–71.
- [28] Kessler RC, Berglund P, Demler O, et al. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62:593–602.
- [29] Klein DN, Schatzberg AF, McGullough JP, et al. Age of onset in chronic major depression: relation to demographic and clinical variables, family history, and treatment response. *J Affect Disord* 1999;55:149–57.
- [30] Atlantis E, Fahey P, Cochrane B, Smith S. Bidirectional associations between clinically relevant depression or anxiety and COPD: a systematic review and meta-analysis. *Chest* 2013;144:766–77.
- [31] Arroll B, Chin WY, Martis W, et al. Antidepressants for treatment of depression in primary care: a systematic review and meta-analysis. *J Prim Health Care* 2016;8:325–34.
- [32] Barth J, Munder T, Gerger H, et al. Comparative efficacy of seven psychotherapeutic interventions for patients with depression: a network meta-analysis. *PLoS Med* 2013;10:e1001454.
- [33] Cuijpers P, Donker T, Weissman MM, et al. Interpersonal psychotherapy for mental health problems: a comprehensive meta-analysis. *Am J Psychiatry* 2016;173:680–7.
- [34] Trudeau L, Spoth R, Mason WA, et al. Effects of adolescent universal substance misuse preventive interventions on young adult depression symptoms: mediational modeling. *J Abnorm Child Psychol* 2016;44:257–68.
- [35] Goodwin FK, Jamison KR. *Manic-Depressive Illness: Bipolar Disorders and Recurrent Depression* (2nd ed.). New York, NY: Oxford University Press, 2007.

- [36] Krishnan KR, Delong M, Kraemer H, et al. Comorbidity of depression with other medical diseases in the elderly. *Biol Psychiatry* 2002;52:559–88.
- [37] Kupfer DJ, Frank E. Comorbidity in depression. *Acta Psychiatr Scand Suppl* 2003;108(Suppl. 418):57–60.
- [38] Dembling BP, Chen DT, Vachon L. Life expectancy and causes of death in a population treated for serious mental illness. *Psychiatr Serv* 1999;50:1036–42.
- [39] Simon GE, Rutter CM, Peterson D, et al. Does response on the PHQ-9 Depression Questionnaire predict subsequent suicide attempt or suicide death? *Psychiatr Serv* 2013;64:1195–202.
- [40] Murray CJ, Abraham J, Ali MK; US Burden of Disease Collaborators. The state of US health, 1990–2010: burden of diseases, injuries, and risk factors. *JAMA* 2013;310:591–608.
- [41] Ferrari AJ, Charlson FJ, Norman RE, et al. Burden of depressive disorders by country, sex, age, and year: findings from the Global Burden of Disease study 2010. *PLoS Med* 2013;10:e1001547.
- [42] Chang CK, Hayes RD, Perera G, et al. Life expectancy at birth for people with serious mental illness and other major disorders from a secondary mental health care case register in London. *PLoS One* 2011;6:e19590.
- [43] Fortes C, Mastroeni S, Alessandra S, et al. The combination of depressive symptoms and smoking shorten life expectancy among the aged. *Int Psychogeriatr* 2012;24:624–30.
- [44] Jia H, Zack MM, Thompson WW, et al. Impact of depression on quality-adjusted life expectancy (QALE) directly as well as indirectly through suicide. *Soc Psychiatry Psychiatr Epidemiol* 2015;50:939–49.
- [45] Gold MR, Siegel JE, Russell RB, Weinstein MC. *Cost-Effectiveness in Health and Medicine*. New York, NY: Oxford University Press, 1996.
- [46] Rosenberg MA, Fryback DG, Lawrence WF. Computing population-based estimates of health-adjusted life expectancy. *Med Decis Making* 1999;19:90–7.
- [47] Jia H, Zack MM, Thompson WW, Dube SR. Quality-adjusted life expectancy (QALE) loss due to smoking in the United States. *Qual Life Res* 2013;22:27–35.
- [48] Brown DS, Jia H, Zack MM, et al. Using health-related quality of life and quality-adjusted life expectancy for effective public health surveillance and prevention. *Expert Rev Pharmacoecon Outcomes Res* 2013;13:425–7.
- [49] Frazier EL, Franks AL, Sanderson LM. Using behavioral risk factor surveillance data. In: *Using Chronic Disease Data: A Handbook for Public Health Practitioners*. Atlanta, GA: Centers for Disease Control and Prevention, 1992.
- [50] Xu F, Town M, Balluz LS, et al. Surveillance for certain health behaviors among states and selected local areas—United States, 2010. *MMWR Surveill Summ* 2013;62:1–247.
- [51] Centers for Disease Control and Prevention. The Behavioral Risk Factor Surveillance System (BRFSS), 2006, 2008, 2010. Available from: [http://www.cdc.gov/brfss/annual\\_data/annual\\_data.htm](http://www.cdc.gov/brfss/annual_data/annual_data.htm). [Accessed December 2, 2016].
- [52] Dhingra SS, Kroenke K, Zack MM, et al. PHQ-8 days: a measurement option for DSM-5 major depressive disorder (MDD) severity. *Popul Health Metr* 2011;9:11.
- [53] Reeves WC, Strine TW, Pratt LA, et al. Mental illness surveillance among adults in the United States. *MMWR Surveill Summ* 2011;60 (Suppl. 3):1–29.
- [54] Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606–13.
- [55] Kroenke K, Strine TW, Spitzer RL, et al. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord* 2009;114:163–73.
- [56] Corson K, Gerrity MS, Dobscha SK. Screening for depression and suicidality in a VA primary care setting: 2 items are better than 1 item. *Am J Manag Care* 2004;10:839–45.
- [57] Centers for Disease Control and Prevention. *Measuring Healthy Days: Population Assessment of Health-Related Quality of Life*. Atlanta, GA: Centers for Disease Control and Prevention, 2000.
- [58] Jia H, Lubetkin EI. Estimating EuroQol EQ-5D scores from population healthy days data. *Med Decis Making* 2008;28:491–9.
- [59] Jia H, Zack MM, Moriarty DG, Fryback DG. Predicting the EuroQol group's EQ-5D index from CDC's "healthy days" in a US sample. *Med Decis Making* 2011;31:174–85.
- [60] Jia H, Zack MM, Thompson WW. State quality-adjusted life expectancy for U.S. adults from 1993 to 2008. *Qual Life Res* 2011;20:853–63.
- [61] Chiang CL. *The Life Table and Its Applications*. Malabar, FL: Robert E. Krieger Publishers, 1984.
- [62] Jia H, Zack MM, Thompson WW. The effects of diabetes, hypertension, asthma, heart disease, and stroke on quality-adjusted life expectancy. *Value Health* 2013;16:140–7.
- [63] US Department of Health and Human Services. *The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, 2014.
- [64] Centers for Disease Control and Prevention. Health effects of cigarette smoking. Available from: [http://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/health\\_effects/effects\\_cig\\_smoking/](http://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/). [Accessed March 24, 2017].
- [65] Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA* 2004;291:1238–45.
- [66] Vogl M, Wenig CM, Leidl R, Pokhrel S. Smoking and health-related quality of life in English general population: implications for economic evaluations. *BMC Public Health* 2012;12:203.
- [67] Schutzer KA, Graves BS. Barriers and motivations to exercise in older adults. *Prev Med* 2004;39:1056–61.
- [68] Stahre M, Roebor J, Kanny D, et al. Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis* 2014;11:E109.
- [69] Kanny D, Brewer RD, Mesnick JB, et al. Vital signs: alcohol poisoning deaths—United States, 2010–2012. *MMWR Morb Mortal Wkly Rep* 2015;263:1238–42.
- [70] Jia H, Zack MM, Thompson WW. Population-based estimates of decreases in quality-adjusted life expectancy associated with unhealthy body mass index. *Public Health Rep* 2016;131:177–84.
- [71] Lehman AF, Myers CP, Corty E. Assessment and classification of patients with psychiatric and substance abuse syndromes. *Hosp Community Psychiatry* 1989;40:1019–25.
- [72] Morgan D. Mindfulness-based cognitive therapy for depression: a new approach to preventing relapse. *Psychother Res* 2003;13:123–5.
- [73] Cummins RA, Gullone E, Lau ALD. A model of subjective well-being homeostasis: the role of personality. In: Gullone E, Cummins RA, eds. *The Universality of Subjective Wellbeing Indicators*. Dordrecht, The Netherlands: Springer, 2002.
- [74] Cristea IA, Huibers MJ, David D, et al. The effects of cognitive behavior therapy for adult depression on dysfunctional thinking: a meta-analysis. *Clin Psychol Rev* 2015;42:62–71.
- [75] Weitz ES, Hollon SD, Twisk J, et al. Baseline depression severity as moderator of depression outcomes between cognitive behavioral therapy vs pharmacotherapy: an individual patient data meta-analysis. *JAMA Psychiatry* 2015;72:1102–9.