

# Chapter 4: Basic Constraint Reasoning (SEND+MORE=MONEY)

Helmut Simonis

CRT-AI CP Week 2025

# Chapter 4: Basic Constraint Reasoning (SEND+MORE=MONEY)

# Licence

This work is licensed under the Creative Commons Attribution-Noncommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/> or send a letter to Creative Commons, 171 Second Street, Suite 300, San Francisco, California, 94105, USA.



# Acknowledgments

This publication has emanated from research conducted with the financial support of Science Foundation Ireland under Grant number 12/RC/2289-P2 at Insight the SFI Research Centre for Data Analytics at UCC, which is co-funded under the European Regional Development Fund.

A version of this material was developed as part of the ECLIPSe ELearning course:

<https://eclipseclp.org/ELearning/index.html>.

Support from Cisco Systems and the Silicon Valley Community Foundation is gratefully acknowledged.

# Example 1: SEND+MORE=MONEY

- Example of Finite Domain Constraint Problem
- Models and Programs
- Constraint Propagation and Search
- Some Basic Constraints: linear arithmetic, alldifferent, disequality
- A Built-in search
- Visualizers for variables, constraints and search

# Outline

Problem

Program

Constraint Setup

Search

Points to Remember

# Problem Definition

## A Crypt-Arithmetic Puzzle

We begin with the definition of the SEND+MORE=MONEY puzzle. It is often shown in the form of a hand-written addition:

$$\begin{array}{rccccccccc} & & & S & E & N & D & & & \\ & & + & M & O & R & E & & & \\ \hline M & O & N & E & Y & & & & & \end{array}$$

The puzzle was first proposed by Henry Dudeney in the Strand Magazine from 1924.

# Rules

- Each character stands for a digit from 0 to 9.
- Numbers are built from digits in the usual, positional notation.
- Repeated occurrence of the same character denote the same digit.
- Different characters denote different digits.
- Numbers do not start with a zero.
- The equation must hold.

$$\begin{array}{rccccc} & S & E & N & D \\ + & M & O & R & E \\ \hline M & O & N & E & Y \end{array}$$



# Outline

Problem

**Program**

Constraint Setup

Search

Points to Remember

# Model

- Each character is a variable, which ranges over the values 0 to 9.
- An *alldifferent* constraint between all variables, which states that two different variables must have different values. This is a very common constraint, which we will encounter in many other problems later on.
- Two *disequality constraints* (variable  $X$  must be different from value  $V$ ) stating that the variables at the beginning of a number can not take the value 0.
- An arithmetic *equality constraint* linking all variables with the proper coefficients and stating that the equation must hold.

# SEND+MORE=MONEY Models

- ECLiPSe [▶ Show](#)
- MiniZinc [▶ Show](#)
- NumberJack [▶ Show](#)
- CPMpy [▶ Show](#)
- Choco-solver [▶ Show](#)

# ECLiPSe Model

```
:- lib(ic).  
  
sendmore(Digits) :-  
    Digits = [S,E,N,D,M,O,R,Y],  
    Digits :: [0..9],  
    alldifferent(Digits),  
    S #\= 0,  
    M #\= 0,  
    1000*S + 100*E + 10*N + D  
        + 1000*M + 100*O + 10*R + E  
    #= 10000*M + 1000*O + 100*N + 10*E + Y,  
    labeling(Digits).
```

► Continue

# MiniZinc Model

```
include "alldifferent.mzn";
var 0..9: S;
var 0..9: E;
var 0..9: N;
var 0..9: D;
var 0..9: M;
var 0..9: O;
var 0..9: R;
var 0..9: Y;
constraint S != 0;
constraint M != 0;
constraint
    1000 * S + 100 * E + 10 * N + D
    + 1000 * M + 100 * O + 10 * R + E
    = 10000 * M + 1000 * O + 100 * N + 10 * E + Y;
constraint alldifferent([S,E,N,D,M,O,R,Y]);

solve satisfy;
```

► Continue

# NumberJack Model (from <https://github.com/eomahony/Numberjack/>)

```
from Numberjack import *

def get_model():
    model = Model()
    s, m = VarArray(2, 1, 9)
    e, n, d, o, r, y = VarArray(6, 0, 9)
    model.add(
        s*1000 + e*100 + n*10 + d +
        m*1000 + o*100 + r*10 + e ==
        m*10000 + o*1000 + n*100 + e*10 + y)
    model.add(AllDiff((s, e, n, d, m, o, r, y)))
    return s, e, n, d, m, o, r, y, model

def solve(param):
    s, e, n, d, m, o, r, y, model = get_model()
    solver = model.load(param['solver'])
    solver.setVerbosity(param['verbose'])
    solver.solve()
```

► Continue

# CPMpy Model (from <https://github.com/CPMpy/>)

```
from cpmPy import *
import numpy as np

s,e,n,d,m,o,r,y = intvar(0,9, shape=8)
model = Model(
    AllDifferent([s,e,n,d,m,o,r,y]),
    (
        sum([s,e,n,d] * np.array([1000, 100, 10, 1])) \
        + sum([m,o,r,e] * np.array([1000, 100, 10, 1])) \
        == sum([m,o,n,e,y] * np.array([10000, 1000, 100, 10, 1]))),
    s > 0,
    m > 0,
)

model.solve()
```

▶ Continue

# Choco-solver Model (from <https://choco-solver.org/>)

```
Model model = new Model("SEND+MORE=MONEY");
IntVar S = model.intVar("S", 1, 9, false);
IntVar E = model.intVar("E", 0, 9, false);
IntVar N = model.intVar("N", 0, 9, false);
IntVar D = model.intVar("D", 0, 9, false);
IntVar M = model.intVar("M", 1, 9, false);
IntVar O = model.intVar("O", 0, 9, false);
IntVar R = model.intVar("R", 0, 9, false);
IntVar Y = model.intVar("Y", 0, 9, false);

model.allDifferent(new IntVar[]{S, E, N, D, M, O, R, Y}).post();

IntVar[] ALL = new IntVar[]{
    S, E, N, D,
    M, O, R, E,
    M, O, N, E, Y};
int[] COEFFS = new int[]{
    1000, 100, 10, 1,
    1000, 100, 10, 1,
    -10000, -1000, -100, -10, -1};
model.scalar(ALL, COEFFS, "=", 0).post();

Solver solver = model.getSolver();
solver.showStatistics();
solver.showSolutions();
solver.findSolution();
```

► Continue




# A Note on Syntax

- Some formulations may seem simpler than others
- This largely is an artifact of a very simple problem
- In most models, you do not write down constraints one by one
- You create constraints based on data
- Ease of integration becomes more important than syntax
- Debugging tools for those who need a debugger :-)

# Choice of Model

- This is *one* model, not *the* model of the problem
- Many possible alternatives
- Choice often depends on your constraint system
  - Constraints available
  - Reasoning attached to constraints
- Not always clear which is the *best* model
- Often: Not clear what is the *problem*

# Running the Program (MiniZinc IDE)



The screenshot shows the MiniZinc IDE interface. The top menu bar includes File, Edit, MiniZinc, View, and Help. Below the menu is a toolbar with icons for New model, Open, Save, Copy, Paste, Undo, Redo, Shift left, Shift right, Run, and Solver configuration. The Solver configuration dropdown is set to 'Gecode 6.1.1 (built-in)'. The main editor displays the MiniZinc model 'sendmore.mzn' with the following code:

```
1 include "alldifferent.mzn";
2 var 0..9: S;
3 var 0..9: E;
4 var 0..9: N;
5 var 0..9: D;
6 var 0..9: M;
7 var 0..9: O;
8 var 0..9: R;
9 var 0..9: Y;
10 constraint S != O;
11 constraint M != O;
12 constraint
13   1000 * S + 100 * E + 10 * N + D
14   + 1000 * M + 100 * O + 10 * R + E
15   = 10000 * M + 1000 * O + 100 * N + 10 * E + Y;
16 constraint alldifferent([S,E,N,D,M,O,R,Y]);
17 solve satisfy;
```

The Output window at the bottom shows the compilation and execution results:

```
Compiling sendmore.mzn
Running sendmore.mzn
S = 9;
E = 5;
N = 6;
D = 7;
M = 1;
O = 0;
R = 8;
Y = 2;
-----
Finished in 3s 103msec
```

The status bar at the bottom indicates 'Line: 16, Col: 1' and '3s 103msec'.

# Question

- But how did the program come up with this solution?
- We show solution with ECLiPse, other solvers vary slightly

# Outline

Problem

Program

## Constraint Setup

- Domain Definition

- Alldifferent Constraint

- Disequality Constraints

- Equality Constraint

Search

Points to Remember

# Domain Definition

```
var 0..9: S;  
var 0..9: E;  
var 0..9: N;  
var 0..9: D;  
var 0..9: M;  
var 0..9: O;  
var 0..9: R;  
var 0..9: Y;
```

# Domain Visualization

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Domain Visualization

Rows =  
Variables

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										



# Domain Visualization

Columns = Values

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Domain Visualization

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M			Cells= State							
O										
R										
Y										

# Alldifferent Constraint

```
include "alldifferent.mzn";
```

```
constraint alldifferent([S,E,N,D,M,O,R,Y]);
```

- Built-in alldifferent predicate included
- No initial propagation possible
- *Suspends*, waits until variables are changed
- When variable is fixed, remove value from domain of other variables
- *Forward checking*

# All different Visualization

Uses the same representation as the domain visualizer

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Disequality Constraints

```
constraint S != 0;  
constraint M != 0;
```

Remove value from domain

$$S \in \{1..9\}, M \in \{1..9\}$$

Constraints solved, can be removed

# Domains after Disequality

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Equality Constraint

- Normalization of linear terms
  - Single occurrence of variable
  - Positive coefficients
- Propagation

# Normalization

$$\begin{array}{rcccccl} & 1000*S+ & 100*E+ & 10*N+ & D & \\ & +1000*M+ & 100*O+ & 10*R+ & E & \\ \hline 10000*M+ & 1000*O+ & 100*N+ & 10*E+ & Y & \end{array}$$



# Normalization

$$\begin{array}{rrrrr} & 1000*S+ & 100*E+ & 10*N+ & D \\ & +\mathbf{1000*M+} & 100*O+ & 10*R+ & E \\ \hline \mathbf{10000*M+} & 1000*O+ & 100*N+ & 10*E+ & Y \end{array}$$

# Normalization

$$\begin{array}{rcccl} & 1000*S+ & 100*E+ & 10*N+ & D \\ & + & 100*O+ & 10*R+ & E \\ \hline 9000*M+ & 1000*O+ & 100*N+ & 10*E+ & Y \end{array}$$

# Normalization

$$\begin{array}{rcccl} & 1000*S+ & 100*E+ & 10*N+ & D \\ & + & \mathbf{100*O+} & 10*R+ & E \\ \hline 9000*M+ & \mathbf{1000*O+} & 100*N+ & 10*E+ & Y \end{array}$$

# Normalization

$$\begin{array}{rcccl} & 1000 * S+ & 100 * E+ & 10 * N+ & D \\ & & + & 10 * R+ & E \\ \hline 9000 * M+ & \mathbf{900 * O+} & 100 * N+ & 10 * E+ & Y \end{array}$$

# Normalization

$$\begin{array}{rcccc} & 1000^*S+ & 100^*E+ & \mathbf{10^*N+} & D \\ & & + & 10^*R+ & E \\ \hline 9000^*M+ & 900^*O+ & \mathbf{100^*N+} & 10^*E+ & Y \end{array}$$

# Normalization

$$\begin{array}{rcccccc} & & 1000*S+ & 100*E+ & & D \\ & & & + & 10*R+ & E \\ \hline 9000*M+ & 900*O+ & \mathbf{90*N+} & 10*E+ & & Y \end{array}$$

# Normalization

$$\begin{array}{rcccccc} & & 1000*S+ & \mathbf{100}*E+ & & D \\ & & & + & 10*R+ & \mathbf{E} \\ \hline 9000*M+ & 900*O+ & 90*N+ & \mathbf{10}*E+ & & Y \end{array}$$

# Normalization

$$\begin{array}{rcccc} & 1000*S+ & 91*E+ & & D \\ & & + & 10*R & \\ \hline 9000*M+ & 900*O+ & 90*N+ & & Y \end{array}$$



# Simplified Equation

$$1000 * S + 91 * E + 10 * R + D = 9000 * M + 900 * O + 90 * N + Y$$

# Propagation

$$1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9} = \\ 9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}$$

# Propagation

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{1000..9918} = \underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..89919}$$

# Propagation

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{9000..9918} = \underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..9918}$$

# Propagation

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{9000..9918} =$$
$$\underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..9918}$$

Deduction:

$$M = 1, S = 9, O \in \{0..1\}$$

# Propagation

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{9000..9918} = \underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..9918}$$

Deduction:

$$M = 1, S = 9, O \in \{0..1\}$$

Why? [▶ Skip](#)

# Consider lower bound for $S$

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{9000..9918} = \underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..9918}$$

- Lower bound of equation is 9000
- Rest of lhs (left hand side) ( $91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}$ ) is atmost 918
- $S$  must be greater or equal to  $\frac{9000-918}{1000} = 8.082$ 
  - otherwise lower bound of equation not reached by lhs
- $S$  is integer, therefore  $S \geq \lceil \frac{9000-918}{1000} \rceil = 9$
- $S$  has upper bound of 9, so  $S = 9$

# Consider upper bound of $M$

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{9000..9918} = \underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..9918}$$

- Upper bound of equation is 9918
- Rest of rhs (right hand side)  $900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}$  is at least 0
- $M$  must be smaller or equal to  $\frac{9918-0}{9000} = 1.102$
- $M$  must be integer, therefore  $M \leq \lfloor \frac{9918-0}{9000} \rfloor = 1$
- $M$  has lower bound of 1, so  $M = 1$



# Consider upper bound of $O$

$$\underbrace{1000 * S^{1..9} + 91 * E^{0..9} + 10 * R^{0..9} + D^{0..9}}_{9000..9918} = \underbrace{9000 * M^{1..9} + 900 * O^{0..9} + 90 * N^{0..9} + Y^{0..9}}_{9000..9918}$$

- Upper bound of equation is 9918
- Rest of rhs (right hand side)  $9000 * 1 + 90 * N^{0..9} + Y^{0..9}$  is at least 9000
- $O$  must be smaller or equal to  $\frac{9918-9000}{900} = 1.02$
- $O$  must be integer, therefore  $O \leq \lfloor \frac{9918-9000}{900} \rfloor = 1$
- $O$  has lower bound of 0, so  $O \in \{0..1\}$

# Propagation of equality: Result

	0	1	2	3	4	5	6	7	8	9
S		-	-	-	-	-	-	-	-	☀
E										
N										
D										
M		☀	-	-	-	-	-	-	-	-
O			✖	✖	✖	✖	✖	✖	✖	✖
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S		-	-	-	-	-	-	-	-	☀
E										
N										
D										
M		☀	-	-	-	-	-	-	-	-
O			✕	✕	✕	✕	✕	✕	✕	✕
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										☀
E										
N										
D										
M		☀								
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M		☀								
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O	☀									
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

$$O = 0, [E, R, D, N, Y] \in \{2..8\}$$



# Waking the equality constraint

- Triggered by assignment of variables
- *or* update of lower or upper bound

# Removal of constants

$$\begin{aligned} 1000 * 9 + 91 * E^{2..8} + 10 * R^{2..8} + D^{2..8} = \\ 9000 * 1 + 900 * 0 + 90 * N^{2..8} + Y^{2..8} \end{aligned}$$

# Removal of constants

$$\begin{aligned} 1000 * 9 + 91 * E^{2..8} + 10 * R^{2..8} + D^{2..8} = \\ 9000 * 1 + 900 * 0 + 90 * N^{2..8} + Y^{2..8} \end{aligned}$$

# Removal of constants

$$91 * E^{2..8} + 10 * R^{2..8} + D^{2..8} = 90 * N^{2..8} + Y^{2..8}$$

# Propagation of equality (Iteration 1)

$$\underbrace{91 * E^{2..8} + 10 * R^{2..8} + D^{2..8}}_{204..816} = \underbrace{90 * N^{2..8} + Y^{2..8}}_{182..728}$$

# Propagation of equality (Iteration 1)

$$\underbrace{91 * E^{2..8} + 10 * R^{2..8} + D^{2..8}}_{204..728} = 90 * N^{2..8} + Y^{2..8}$$

# Propagation of equality (Iteration 1)

$$\underbrace{91 * E^{2..8} + 10 * R^{2..8} + D^{2..8} = 90 * N^{2..8} + Y^{2..8}}_{204..728}$$

$$N \geq 3 = \lceil \frac{204 - 8}{90} \rceil, E \leq 7 = \lfloor \frac{728 - 22}{91} \rfloor$$

## Propagation of equality (Iteration 2)

$$91 * E^{2..7} + 10 * R^{2..8} + D^{2..8} = 90 * N^{3..8} + Y^{2..8}$$



## Propagation of equality (Iteration 2)

$$\underbrace{91 * E^{2..7} + 10 * R^{2..8} + D^{2..8}}_{204..725} = \underbrace{90 * N^{3..8} + Y^{2..8}}_{272..728}$$

## Propagation of equality (Iteration 2)

$$\underbrace{91 * E^{2..7} + 10 * R^{2..8} + D^{2..8}}_{272..725} = 90 * N^{3..8} + Y^{2..8}$$

## Propagation of equality (Iteration 2)

$$\underbrace{91 * E^{2..7} + 10 * R^{2..8} + D^{2..8}}_{272..725} = 90 * N^{3..8} + Y^{2..8}$$

$$E \geq 3 = \lceil \frac{272 - 88}{91} \rceil$$

## Propagation of equality (Iteration 3)

$$91 * E^{3..7} + 10 * R^{2..8} + D^{2..8} = 90 * N^{3..8} + Y^{2..8}$$

## Propagation of equality (Iteration 3)

$$\underbrace{91 * E^{3..7} + 10 * R^{2..8} + D^{2..8}}_{295..725} = \underbrace{90 * N^{3..8} + Y^{2..8}}_{272..728}$$

## Propagation of equality (Iteration 3)

$$\underbrace{91 * E^{3..7} + 10 * R^{2..8} + D^{2..8}}_{295..725} = 90 * N^{3..8} + Y^{2..8}$$

## Propagation of equality (Iteration 3)

$$\underbrace{91 * E^{3..7} + 10 * R^{2..8} + D^{2..8}}_{295..725} = 90 * N^{3..8} + Y^{2..8}$$

$$N \geq 4 = \lceil \frac{295 - 8}{90} \rceil$$

## Propagation of equality (Iteration 4)

$$91 * E^{3..7} + 10 * R^{2..8} + D^{2..8} = 90 * N^{4..8} + Y^{2..8}$$



## Propagation of equality (Iteration 4)

$$\underbrace{91 * E^{3..7} + 10 * R^{2..8} + D^{2..8}}_{295..725} = \underbrace{90 * N^{4..8} + Y^{2..8}}_{362..728}$$

## Propagation of equality (Iteration 4)

$$\underbrace{91 * E^{3..7} + 10 * R^{2..8} + D^{2..8}}_{362..725} = 90 * N^{4..8} + Y^{2..8}$$

## Propagation of equality (Iteration 4)

$$\underbrace{91 * E^{3..7} + 10 * R^{2..8} + D^{2..8}}_{362..725} = 90 * N^{4..8} + Y^{2..8}$$

$$E \geq 4 = \lceil \frac{362 - 88}{91} \rceil$$

## Propagation of equality (Iteration 5)

$$91 * E^{4..7} + 10 * R^{2..8} + D^{2..8} = 90 * N^{4..8} + Y^{2..8}$$

## Propagation of equality (Iteration 5)

$$\underbrace{91 * E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{386..725} = \underbrace{90 * N^{4..8} + Y^{2..8}}_{362..728}$$

# Propagation of equality (Iteration 5)

$$\underbrace{91 * E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{386..725} = 90 * N^{4..8} + Y^{2..8}$$

## Propagation of equality (Iteration 5)

$$\underbrace{91 * E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{386..725} = 90 * N^{4..8} + Y^{2..8}$$

$$N \geq 5 = \lceil \frac{386 - 8}{90} \rceil$$

## Propagation of equality (Iteration 6)

$$91 * E^{4..7} + 10 * R^{2..8} + D^{2..8} = 90 * N^{5..8} + Y^{2..8}$$



## Propagation of equality (Iteration 6)

$$\underbrace{91 * E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{386..725} = \underbrace{90 * N^{5..8} + Y^{2..8}}_{452..728}$$

## Propagation of equality (Iteration 6)

$$\underbrace{91 * E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{452..725} = 90 * N^{5..8} + Y^{2..8}$$

## Propagation of equality (Iteration 6)

$$\underbrace{91 * E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{452..725} = 90 * N^{5..8} + Y^{2..8}$$

$$N \geq 5 = \lceil \frac{452 - 8}{90} \rceil, E \geq 4 = \lceil \frac{452 - 88}{91} \rceil$$

No further propagation at this point

# Domains after setup

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Outline

Problem

Program

Constraint Setup

Search

Step 1

Step 2

Further Steps

Solution

Points to Remember

solve satisfy;

- Try to find a feasible solution, choice left to solver
- Naive search strategy shown here
  - Try variable in order given
  - Try values starting from smallest value in domain
  - When failing, backtrack to last open choice
  - *Chronological Backtracking*
  - *Depth First search*


# Search Tree Step 1

S  
9  
E

Variable  $S$  already fixed

## Step 2, Alternative $E = 4$

Variable  $E \in \{4..7\}$ , first value tested is 4





# Assignment $E = 4$

	0	1	2	3	4	5	6	7	8	9
S										
E					☀	-	-	-		
N										
D										
M										
O										
R										
Y										

## Propagation of $E = 4$ , equality constraint

$$91 * 4 + 10 * R^{2..8} + D^{2..8} = 90 * N^{5..8} + Y^{2..8}$$

## Propagation of $E = 4$ , equality constraint

$$\underbrace{91 * 4 + 10 * R^{2..8} + D^{2..8}}_{386..452} = \underbrace{90 * N^{5..8} + Y^{2..8}}_{452..728}$$

## Propagation of $E = 4$ , equality constraint

$$\underbrace{91 * 4 + 10 * R^{2..8} + D^{2..8}}_{452} = 90 * N^{5..8} + Y^{2..8}$$

## Propagation of $E = 4$ , equality constraint

$$\underbrace{91 * 4 + 10 * R^{2..8} + D^{2..8}}_{452} = 90 * N^{5..8} + Y^{2..8}$$

$$N = 5, Y = 2, R = 8, D = 8$$

# Result of equality propagation

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of alldifferent


	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

Alldifferent fails!



## Step 2, Alternative $E = 5$

Return to last open choice,  $E$ , and test next value



# Assignment $E = 5$

	0	1	2	3	4	5	6	7	8	9
S										
E					-	☀	-	-		
N										
D										
M										
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E					-	☀	-	-		
N										
D										
M										
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of all different

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

$N \neq 5, N \geq 6$

# Propagation of equality

$$91 * 5 + 10 * R^{2..8} + D^{2..8} = 90 * N^{6..8} + Y^{2..8}$$

# Propagation of equality

$$\underbrace{91 * 5 + 10 * R^{2..8} + D^{2..8}}_{477..543} = \underbrace{90 * N^{6..8} + Y^{2..8}}_{542..728}$$

# Propagation of equality

$$\underbrace{91 * 5 + 10 * R^{2..8} + D^{2..8} = 90 * N^{6..8} + Y^{2..8}}_{542..543}$$



# Propagation of equality

$$\underbrace{91 * 5 + 10 * R^{2..8} + D^{2..8}}_{542..543} = 90 * N^{6..8} + Y^{2..8}$$

$$N = 6, Y \in \{2, 3\}, R = 8, D \in \{7..8\}$$

# Result of equality propagation

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of alldifferent

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of alldifferent

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of alldifferent

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

# Propagation of alldifferent

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										

$$D = 7$$

# Propagation of equality

$$91 * 5 + 10 * 8 + 7 = 90 * 6 + Y^{2..3}$$

# Propagation of equality

$$\underbrace{91 * 5 + 10 * 8 + 7}_{542} = \underbrace{90 * 6 + Y^{2..3}}_{542..543}$$



# Propagation of equality

$$\underbrace{91 * 5 + 10 * 8 + 7}_{542} = 90 * 6 + Y^{2..3}$$


# Propagation of equality

$$\underbrace{91 * 5 + 10 * 8 + 7 = 90 * 6 + Y^{2..3}}_{542}$$
$$Y = 2$$


# Last propagation step

	0	1	2	3	4	5	6	7	8	9
S										
E										
N										
D										
M										
O										
R										
Y										


# Further Steps: Nothing more to do




# Further Steps: Nothing more to do




# Further Steps: Nothing more to do



# Further Steps: Nothing more to do




# Further Steps: Nothing more to do







# Further Steps: Nothing more to do




# Further Steps: Nothing more to do



# Complete Search Tree



# Search Tree with Gecode/GIST



# Some Differences

- Uses binary branching
  - var equal value, var not equal value
- Solutions in green, failure leafs in red, internal nodes in blue
- By default, shows all failed sub trees collapsed
- By default, uses different search strategy

# Solution

$$\begin{array}{r} 9567 \\ + 1085 \\ \hline 10652 \end{array}$$

# Outline

Problem

Program

Constraint Setup

Search

Points to Remember

# Points to Remember

- Constraint models are expressed by variables and constraints.
- Problems can have many different models, which can behave quite differently. Choosing the best model is an art.
- Constraints can take many different forms.
- Propagation deals with the interaction of variables and constraints.
- It removes some values that are inconsistent with a constraint from the domain of a variable.
- Constraints only communicate via shared variables.



# Points to Remember

- Propagation usually is not sufficient, search may be required to find a solution.
- Propagation is data driven, and can be quite complex even for small examples.
- The default search uses chronological depth-first backtracking, systematically exploring the complete search space.
- The search choices and propagation are interleaved, after every choice some more propagation may further reduce the problem.

# For Puzzle Purists Only

- We did not follow the puzzler ethos!
- We should solve the puzzle without making choices
- Even a case analysis should be avoided
- The puzzle has a single solution, we should be able to deduce the solution
- In the process shown, we are limited by the underlying assumptions
  - Treat each constraint on its own, they interact only by domains of variables
  - We only use the constraints that we stated in the model
- Can we do better than that?

► Skip

# Better Reasoning

- Three possible approaches (possibly many more)
  - Full domain reasoning for arithmetic, not just bound reasoning
  - Interaction of *sum* and *alldifferent* constraints
  - Deduced implied constraint

# Looking at more than just bounds

- We only considered the smallest and largest values that can be achieved in the sum constraint
- We can do more
  - Can any of the values between be expressed as the sum of the terms
  - Consider holes in the domains, and in the range of possible values for LHS and RHS
- Usually not done in actual solvers for arithmetic constraints
- Easy to do with Dynamic Programming

# Consider the interaction of multiple constraints

- Usually ignored, as only interaction is via domains of shared variables
- Here: Sum and *alldifferent* interact
  - When considering the bounds, we cannot assume that each variable takes its smallest/largest value independently
  - Find feasible assignment that minimizes/maximizes the total weight
  - To do this properly, we need some non-trivial reasoning
- Do we do this combined reasoning automatically, or only when prompted by the modeler?

# Deduced Implied Constraints

- Look at the partially solved puzzle

$$\begin{array}{r} 9\text{END} \\ +10\text{RE} \\ \hline \end{array}$$

- 10NEY
- In the hundreds position, we have  
 $E + 0 + C_{10} = N + 10 * C_{100}$ , with  $C_{10}$  the 0/1 carry from the tens position
- NB: No carry  $C_{100}$  into the thousands,  $C_{100} = 0$
- $N$  must be equal to  $E + 1$  with  $C_{10} = 1$
- If  $C_{10} = 0$ , then  $N = E$ , not possible
- We can substitute  $N = E + 1$  into our main equation, but keep  $N = E + 1$  as well

# Expert Mode Reasoning

Starting with

$$91 * E^{4..7} + 10 * R^{2..8} + D^{2..8} = 90 * N^{5..8} + Y^{2..8}$$

# Expert Mode Reasoning

we get

$$91 * E^{4..7} + 10 * R^{2..8} + D^{2..8} = 90 * E^{4..7} + 90 + Y^{2..8}$$



# Expert Mode Reasoning

Eliminating duplicate occurrences of E

$$\underbrace{E^{4..7} + 10 * R^{2..8} + D^{2..8}}_{26..95} = \underbrace{90 + Y^{2..8}}_{92..98}$$

shared range 92..95

# Expert Mode Reasoning

To reach 92, R must be equal to 8, therefore N, E, D, Y must be less than 8

As  $N = E + 1$ , E must be less than 7

$$E^{4..6} + 10 * 8 + D^{2..7} = 90 + Y^{2..7}$$

# Expert Mode Reasoning

Simplification yields

$$\underbrace{E^{4..6} + D^{2..7}}_{6..13} = \underbrace{10 + Y^{2..7}}_{12..17}$$

shared range 12..13

# Expert Mode Reasoning

To reach 12 on LHS, E must be greater than 4, D must be greater than 5

To reach 13 in RHS, Y must be smaller than 4

$$E^{5..6} + D^{6..7} = 10 + Y^{2..3}$$

# Expert Mode Reasoning

As both N and D must be in 6..7, no other variable can use those values

NB: This requires better reasoning than *forward checking* on the *alldifferent* constraint

If we remove 6 from the domain of E, E must be  $E = 5$ , and thus  $N = 6$ , due to  $N = E + 1$

*Alldifferent* forces  $D = 7$ , leaving

$$5 + 7 = 10 + Y^{2..4}$$

# Expert Mode Reasoning

This only leaves  $Y = 2$

$$5 + 7 = 10 + 2$$

That is a solution, the constraint is satisfied

# Expert Mode Summary

- Often there is more propagation that can be done
- Can be difficult/expensive to do
- Balancing
  - How much work it done at each step of search?
  - How many steps of search you need?
- For hard problems, doing all possible propagation may be exponential
- Not aware that any CP system does the full reasoning shown here

# This is how people solve the puzzle by hand



Send More Money | Puzzle only a genius can solve | MAX MATH GAMES



Max Math Games  
1.28K subscribers

Subscribe

14



Share



Download



Clip



2K views · 3 years ago · #mathsandphysicsfun

- When writing the first version of this puzzle for CHIP (in 1986), we wanted to mimic the way we solve the puzzle by hand