

# Poincaré conjecture

- ▶ Every simply connected, closed 3-manifold is homeomorphic to the 3-sphere.
- ▶ Proven by Grigori Perelman in 2003.
- ▶ Uses Richard S. Hamilton's Ricci flow.

# Birch and Swinnerton-Dyer conjecture

- Predicts a relationship between the rank of the group of rational points and the behavior of the L-function at  $s = 1$ .

# Birch and Swinnerton-Dyer conjecture

- ▶ Predicts a relationship between the rank of the group of rational points and the behavior of the L-function at  $s = 1$ .
- ▶ Iwasawa theory with Euler systems (Kolyvagin–Rubin) is currently the most promising approach.

# Birch and Swinnerton-Dyer conjecture

- ▶ Predicts a relationship between the rank of the group of rational points and the behavior of the L-function at  $s = 1$ .
- ▶ Iwasawa theory with Euler systems (Kolyvagin–Rubin) is currently the most promising approach.
- ▶ Profound impact on number theory and cryptography; would clarify ranks of elliptic curves and reshape algorithms and security assumptions.

# Hodge conjecture

- ▶ States that certain de Rham cohomology classes are algebraic.

# Hodge conjecture

- ▶ States that certain de Rham cohomology classes are algebraic.
- ▶ Motivic cohomology and derived algebraic geometry (Voevodsky–Deligne program) offer potential pathways.

# Hodge conjecture

- ▶ States that certain de Rham cohomology classes are algebraic.
- ▶ Motivic cohomology and derived algebraic geometry (Voevodsky–Deligne program) offer potential pathways.
- ▶ Would unify topology and algebraic geometry; enable algorithmic methods for classifying algebraic varieties and deepen understanding of the structure of complex manifolds.

# Navier–Stokes existence and smoothness

- ▶ Governs the motion of fluid substances.



# Navier–Stokes existence and smoothness

- ▶ Governs the motion of fluid substances.
- ▶ The existence and smoothness of solutions in three dimensions is unknown.

# Navier–Stokes existence and smoothness

- ▶ Governs the motion of fluid substances.
- ▶ The existence and smoothness of solutions in three dimensions is unknown.
- ▶ Most promising method: convex integration for constructing wild weak solutions (De Lellis–Székelyhidi), and scale-critical regularity via harmonic analysis (Koch–Tataru).

# P versus NP

- ▶ Asks whether every efficiently verifiable problem is also efficiently solvable.

# P versus NP

- ▶ Asks whether every efficiently verifiable problem is also efficiently solvable.
- ▶ Foundational in complexity theory; impacts optimization, cryptography, and AI.

# P versus NP

- ▶ Asks whether every efficiently verifiable problem is also efficiently solvable.
- ▶ Foundational in complexity theory; impacts optimization, cryptography, and AI.
- ▶ Key lines of attack: circuit lower bounds, barriers from natural proofs (Razborov–Rudich), and algebraic/geometric complexity (VP vs VNP).

# P versus NP

- ▶ Asks whether every efficiently verifiable problem is also efficiently solvable.
- ▶ Foundational in complexity theory; impacts optimization, cryptography, and AI.
- ▶ Key lines of attack: circuit lower bounds, barriers from natural proofs (Razborov–Rudich), and algebraic/geometric complexity (VP vs VNP).
- ▶ Consequences: if  $P = NP$ , many cryptosystems collapse; if  $P \neq NP$ , formal hardness underpins secure cryptography and explains intractability.