


**EXERCISE 10.1**

**PAGE NO: 64**

1. AD is a diameter of a circle and AB is a chord. If AD = 34 cm, AB = 30 cm, the distance of AB from the centre of the circle is :

- (A) 17 cm
- (B) 15 cm
- (C) 4 cm
- (D) 8 cm

**Solution:**



**(D) 8 cm**

Explanation:

Given: Diameter of the circle =  $d = AD = 34$  cm

$\therefore$  Radius of the circle =  $r = d/2 = AO = 17$  cm

Length of chord AB = 30 cm

Since the line drawn through the center of a circle to bisect a chord is perpendicular to the chord, therefore AOP is a right angled triangle with L as the bisector of AB.

$\therefore AP = 1/2(AB) = 15$  cm

In right angled triangle AOB, by Pythagoras theorem, we have:

$$(AO)^2 = (OP)^2 + (AP)^2$$

$$\Rightarrow (17)^2 = (OP)^2 + (15)^2$$

$$\Rightarrow (OP)^2 = (17)^2 - (15)^2$$

$$\Rightarrow (OP)^2 = 289 - 225$$

$$\Rightarrow (OP)^2 = 64$$

Take square root on both sides:

$$\Rightarrow (OP) = 8$$

$\therefore$  The distance of AB from the center of the circle is 8 cm.

Hence, option D is the correct answer.


2. In Fig. 10.3, if  $OA = 5$  cm,  $AB = 8$  cm and  $OD$  is perpendicular to  $AB$ , then  $CD$  is equal to:

- (A) 2 cm
- (B) 3 cm
- (C) 4 cm
- (D) 5 cm

**Solution:**

(A) 2 cm

Explanation:



Given:

Radius of the circle =  $r = AO = 5$  cm

Length of chord  $AB = 8$  cm

Since the line drawn through the center of a circle to bisect a chord is perpendicular to the chord, therefore  $AOC$  is a right angled triangle with  $C$  as the bisector of  $AB$ .

$$\therefore AC = \frac{1}{2} (AB) = \frac{8}{2} = 4 \text{ cm}$$

In right angled triangle  $AOC$ , by Pythagoras theorem, we have:

$$(AO)^2 = (OC)^2 + (AC)^2$$

$$\Rightarrow (5)^2 = (OC)^2 + (4)^2$$

$$\Rightarrow (OC)^2 = (5)^2 - (4)^2$$

$$\Rightarrow (OC)^2 = 25 - 16$$

$$\Rightarrow (OC)^2 = 9$$

Take square root on both sides:

$$\Rightarrow (OC) = 3$$

$\therefore$  The distance of  $AC$  from the center of the circle is 3 cm.

Now,  $OD$  is the radius of the circle,  $\therefore OD = 5$  cm

$$CD = OD - OC$$

$$CD = 5 - 3$$

$$CD = 2$$

Therefore,  $CD = 2$  cm

Hence, option A is the correct answer.


3. If  $AB = 12$  cm,  $BC = 16$  cm and  $AB$  is perpendicular to  $BC$ , then the radius of the circle passing through the points  $A$ ,  $B$  and  $C$  is :

- (A) 6 cm
- (B) 8 cm
- (C) 10 cm
- (D) 12 cm

**Solution:**

(C) 10 cm

Explanation:



According to the question,

$AB = 12$  cm,  $BC = 16$  cm,  $AB \perp BC$ .

Therefore,

$AC$  is the diameter of the circle passing through the points  $A$ ,  $B$  and  $C$ .

Now, according to the figure,

We get,

$ABC$  is a right angled triangle.

By Pythagoras theorem:

$$(AC)^2 = (CB)^2 + (AB)^2$$

$$\Rightarrow (AC)^2 = (16)^2 + (12)^2$$

$$\Rightarrow (AC)^2 = 256 + 144$$

$$\Rightarrow (AC)^2 = 400$$

Take square root on LHS and RHS,

We get,

$$(AC) = 20$$

Diameter of the circle = 20 cm

Thus, radius of the circle = Diameter/2

$$= 20/2$$


$$= 10 \text{ cm}$$

Hence, Radius of the circle = 10 cm

Hence, option C is the correct answer.

**4. In Fig.10.4, if  $\angle ABC = 20^\circ$ , then  $\angle AOC$  is equal to:**

- (A)  $20^\circ$
- (B)  $40^\circ$
- (C)  $60^\circ$
- (D)  $10^\circ$



**Fig. 10.4**

**Solution:**

**(B)  $40^\circ$**

Explanation:

According to the question,

$$\angle ABC = 20^\circ$$

We know that,

“The angle subtended by an arc at the center of a circle is twice the angle subtended by it at remaining part of the circle”

According to the theorem, we have,

$$\angle AOC = 2 \times \angle ABC$$

$$= 2 \times 20^\circ$$


$$= 40^\circ$$

Therefore,  $\angle AOC = 40^\circ$

Hence, option B is the correct answer.

**5. In Fig.10.5, if AOB is a diameter of the circle and  $AC = BC$ , then  $\angle CAB$  is equal to:**

- (A)  $30^\circ$
- (B)  $60^\circ$
- (C)  $90^\circ$
- (D)  $45^\circ$



**Fig. 10.5**

**Solution:**

**(D)  $45^\circ$**

Explanation:

According to the question,

We have,

Diameter of the circle = AOB

$AC = BC$

Since, angles opposite to equal sides are equal

$\angle ABC = \angle BAC$

Let,  $\angle ABC = \angle BAC = x$

Also, diameter subtends a right angle to the circle,

$\angle ACB = 90^\circ$

We also know that,

By angle sum property of a triangle, sum of all angles of a triangle =  $180^\circ$ .

$\angle CAB + \angle ABC + \angle ACB = 180^\circ$

$\Rightarrow x + x + 90^\circ = 180^\circ$

$\Rightarrow 2x = 90^\circ$

$\Rightarrow x = 45^\circ$

$\angle CAB = \angle ABC = 45^\circ$

Hence, option D is the correct answer.

**EXERCISE 10.2**

**PAGE NO: 64**

**Write True or False and justify your answer in each of the following:**

**1. Two chords AB and CD of a circle are each at distances 4 cm from the centre. Then  $AB = CD$ .**

**Solution:**

True

Justification:

Given that AB and AC are chords that are at a distance of 4 cm from center of a circle.

Since, chords that are equidistant from the center of a circle are equal in length,

We have,  $AB = CD$ .


**2. Two chords AB and AC of a circle with centre O are on the opposite sides of OA. Then  $\angle OAB = \angle OAC$ .**

**Solution:**

False

Justification:

Let AB and AC be the chord of the circle with center O on the opposite side of OA.



Consider the triangles AOC and AOB:

$AO = AO$  (Common side in both triangles)

$OB = OC$  (Both OB and OC are radius of circle)

But we can't show that either the third side of both triangles are equal or any angle is equal.

Therefore  $\triangle AOB$  is not congruent to  $\triangle AOC$ .

$\therefore \angle OAB \neq \angle OAC$ .

**3. Two congruent circles with centres O and O' intersect at two points A and B. Then  $\angle AOB = \angle AO'B$ .**

**Solution:**

True

Justification:

Equal chords of congruent circles subtend equal angles at the respective centre.

Hence, the given statement is true.

**4. Through three collinear points a circle can be drawn.**

**Solution:**

False

Justification:

A circle through two points cannot pass through a point which is collinear to these two points.  
Hence, the given statement is false.

**5. A circle of radius 3 cm can be drawn through two points A, B such that  $AB = 6$  cm.**

**Solution:**

True

Justification:

According to the question,

Radius of circle = 3 cm

Diameter of circle =  $2 \times r$

$$= 2 \times 3 \text{ cm}$$

$$= 6 \text{ cm}$$

Now,

From the question we have,

$$AB = 6 \text{ cm}$$

So, the given statement is true because AB will be the diameter

**EXERCISE 10.3**

**PAGE NO: 64**

**1. If arcs AXB and CYD of a circle are congruent, find the ratio of AB and CD.**

**Solution:**

According to the question,

We have,

$\text{AXB} \cong \text{CYD}$ .

We know that,

If two arcs of a circle are congruent, then their corresponding arcs are also equal.

So, we have chord  $AB = \text{chord } CD$ .

Hence, we get,


$AB/CD = 1$

$AB/CD = 1/1$

$AB : CD = 1:1$

**2. If the perpendicular bisector of a chord AB of a circle PXAQBY intersects the circle at P and Q, prove that arc PXA  $\angle$  Arc PYB.**

**Solution:**



According to the question,

We have,

PQ is the perpendicular bisect of AB,

So, we get,

$AM = BM \dots \text{eq.}(1)$

In  $\triangle APM$  and  $\triangle BPM$ ,

From eq.(1),

$AM = BM$

$\angle AMP = \angle BMP = 90^\circ$

$PM = PM$  [Common side]

Therefore,  $\triangle APM \cong \triangle BPM$  [By SAS congruence rule]

So,  $AP = BP$  [CPCT]

Hence, arc PXA  $\cong$  Arc PYB




Therefore, if two chords of a circle are equal, then their corresponding arcs are congruent.

**3. A, B and C are three points on a circle. Prove that the perpendicular bisectors of AB, BC and CA are concurrent.**

**Solution:**

According to the question,

Three non-collinear points A, B and C are on a circle.



To prove: Perpendicular bisectors of AB, BC and CA are concurrent.

Construction: Join AB, BC and CA.

Draw:

ST, perpendicular bisector of AB,

PM, perpendicular bisector of BC

And, QR perpendicular bisector of CA

As point A, B and C are not collinear, ST, PM and QR are not parallel and will intersect.

Proof:

O lies on ST, the  $\perp$  bisector of AB

$$OA = OB \dots (1)$$

Similarly, O lies on PM, the  $\perp$  bisector of BC

$$OB = OC \dots (2)$$

And, O lies on QR, the  $\perp$  bisector of CA

$$OC = OA \dots (3)$$

From (1), (2) and (3),

$$OA = OB = OC$$

Let  $OA = OB = OC = r$


Draw circle, with centre O and radius r, passing through A, B and C.

Hence, O is the only point equidistance from A, B and C.

Therefore, the perpendicular bisectors of AB, BC and CA are concurrent.

**4. AB and AC are two equal chords of a circle. Prove that the bisector of the angle BAC passes through the centre of the circle.**

**Solution:**



According to the question,

We have,

AB and AC are two chords which are equal with centre O.

AM is the bisector of  $\angle BAC$ .

To prove: AM passes through O.

Construction: Join BC.

Let AM intersect BC at P.

Proof: In  $\triangle BAP$  and  $\triangle CAP$

$AB = AC$  [Given]

$\angle BAP = \angle CAP$  [Given]

And  $AP = AP$  [Common side]

$\triangle BAP \cong \triangle CAP$  [By SAS]

$\angle BPA = \angle CPA$  [CPCT]

We know that,

$CP = BP$

But, since  $\angle BPA$  and  $\angle CPA$  are linear pair angles,

We have,

$\angle BPA + \angle CPA = 180^\circ$

$\angle BPA = \angle CPA = 90^\circ$

Then,


AP is perpendicular bisector of the chord BC, which will pass through the centre O on being produced.

Therefore, AM passes through O.

**5. If a line segment joining mid-points of two chords of a circle passes through the centre of the circle, prove that the two chords are parallel.**

**Solution:**

Consider AB and CD to be the chords of the circle with center O.



Let L be the midpoint of AB.

Let M be the midpoint of CD.

Let PQ be the line passing through these midpoints and the center of the circle.

Then, PQ is the diameter of the circle.

We know that,

Line joining center to the midpoint of a chord is always perpendicular to the chord.

Since M is the midpoint of CD,

We have,  $OM \perp CD$

$\Rightarrow \angle OMD = 90^\circ$

Similarly, L is the midpoint of AB,

$OL \perp AB$

$\Rightarrow \angle OLA = 90^\circ$

But, we know,

$\angle OLA$  and  $\angle OMD$  are alternate angles.

So,  $AB \parallel CD$ .

Hence, proved.

**6. ABCD is such a quadrilateral that A is the centre of the circle passing through B, C and D.**

**Prove that  $\angle CBD + \angle CDB = \frac{1}{2} \angle BAD$**

**Solution:**


According to the question,

We have,

A quadrilateral ABCD such that A is the centre of the circle passing through B, C and D.

Construction:

Join CA and BD.



We know that,

In a circle, angle subtended by an arc at the center is twice the angle subtended by it at any other point in the remaining part of the circle

So,

The arc DC subtends  $\angle DAC$  at the center and  $\angle CAB$  at point B in the remaining part of the circle,

We get,

$$\angle DAC = 2\angle CBD \dots(1)$$

Similarly,

The arc BC subtends  $\angle CAB$  at the center and  $\angle CDB$  at point D in the remaining part of the circle,

We get,

$$\angle CAB = 2\angle CDB \dots(2)$$

From equations (1) and (2),

We have:

$$\angle DAC + \angle CAB = 2\angle CDB + 2\angle CBD$$

$$\Rightarrow \angle BAD = 2(\angle CDB + \angle CBD)$$

$$\Rightarrow (\angle CDB + \angle CBD) = \frac{1}{2} (\angle BAD)$$


**7. O is the circumcentre of the triangle ABC and D is the mid-point of the base BC. Prove that  $\angle BOD = \angle A$ .**

**Solution:**

According to the question,

We have,

O is the circumcenter of the triangle ABC and D is the midpoint of BC.



To prove:  $\angle BOD = \angle A$

Construction: Join OB and OC.

In  $\triangle OBD$  and  $\triangle OCD$ :

$OD = OD$  (common)

$DB = DC$  (D is the midpoint of BC)

$OB = OC$  (radius of the circle)

By SSS congruence rule,

We get,

$\triangle OBD \cong \triangle OCD$ .

$\angle BOD = \angle COD$  (By CPCT)

Let  $\angle BOD = \angle COD = x$

We know that,

Angle subtended by an arc at the center of the circle is twice the angle subtended by it at any other point in the remaining part of the circle.

So, we have,

$2\angle BAC = \angle BOC$

$\Rightarrow 2\angle BAC = \angle BOD + \angle DOC$

$\Rightarrow 2\angle BAC = x + x$

$\Rightarrow 2\angle BAC = 2x$

$\Rightarrow \angle BAC = x$

$\Rightarrow \angle BAC = \angle BOD$

Hence, proved.


**8. On a common hypotenuse AB, two right triangles ACB and ADB are situated on opposite sides. Prove that  $\angle BAC = \angle BDC$ .**

**Solution:**

According to the question,

We have,

ACB and ADB are two right triangles.



To Prove:  $\angle BAC = \angle BDC$

We know that,

ACB and ADB are right angled triangles,

Then,

$$\angle C + \angle D = 90^\circ + 90^\circ$$

$$\angle C + \angle D = 180^\circ$$

Therefore ADBC is a cyclic quadrilateral as sum of opposite angles of a cyclic quadrilateral =  $180^\circ$

We also have,

$\angle BAC$  and  $\angle BDC$  lie in the same segment BC and angles in the same segment of a circle are equal.

$$\therefore \angle BAC = \angle BDC.$$

Hence Proved.

**9. Two chords AB and AC of a circle subtends angles equal to  $90^\circ$  and  $150^\circ$ , respectively at the centre. Find  $\angle BAC$ , if AB and AC lie on the opposite sides of the centre.**


**Solution:**

According to the question,

We have,

In  $\triangle AOB$ ,

$OA = OB$  (radius of the circle)



Since angle opposite to equal sides are equal, we get,

$$\angle OBA = \angle OAB$$

We know that,

According to angle sum property, sum of all angles of a triangle =  $180^\circ$

Using the angle sum property in  $\triangle AOB$ , we get,

$$\angle OAB + \angle AOB + \angle OBA = 180^\circ$$

$$\Rightarrow \angle OAB + 90^\circ + \angle OAB = 180^\circ$$

$$\Rightarrow 2\angle OAB = 180^\circ - 90^\circ$$

$$\Rightarrow 2\angle OAB = 90^\circ$$

$$\Rightarrow \angle OAB = 45^\circ$$

Now, in  $\triangle AOC$ ,

$OA = OC$  (radius of the circle)

Since, angle opposite to equal sides are equal

$$\therefore \angle OCA = \angle OAC$$

Using the angle sum property in  $\triangle AOB$ , sum of all angles of the triangle is  $180^\circ$ , we have:

$$\angle OAC + \angle AOC + \angle OCA = 180^\circ$$

$$\Rightarrow \angle OAC + 150^\circ + \angle OAC = 180^\circ$$

$$\Rightarrow 2\angle OAC = 180^\circ - 150^\circ$$

$$\Rightarrow 2\angle OAC = 30^\circ$$

$$\Rightarrow \angle OAC = 15^\circ$$

Now,  $\angle BAC = \angle OAB + \angle OAC$

$$= 45^\circ + 15^\circ$$


$$= 60^\circ$$

$$\therefore \angle BAC = 60^\circ$$

**10. If BM and CN are the perpendiculars drawn on the sides AC and AB of the triangle ABC, prove that the points B, C, M and N are concyclic.**

**Solution:**





According to the question,

BM and CN are the perpendiculars drawn on the sides AC and AB of the triangle ABC.

So, we have,

$$\angle BMC = \angle BNC = 90^\circ$$

We know that,

If a line segment joining two points subtends equal angles on the same side of the line containing the segment, then the four points are concyclic.

Considering the question,

Since BC joins the two points, B and C, subtending equal angles,  $\angle BMC$  and  $\angle BNC$ , at M and N on the same side BC containing the segment, then B, C, M and N are concyclic.

Hence, we get that,

B, C, M and N are concyclic.



**EXERCISE 10.4**

**PAGE NO: 64**

**1. If two equal chords of a circle intersect, prove that the parts of one chord are separately equal to the parts of the other chord.**

**Solution:**


According to the question,

AB and CD are two equal chords of a circle with centre O, intersect each other at M.

To prove:

(i)  $MB = MC$  and

(ii)  $AM = MD$



**Proof:**

AB is a chord and  $OE \perp$  to AB from the centre O,

Since, perpendicular from the centre to a chord bisect the chord

We get,

$$AE = \frac{1}{2} AB$$

Similarly,

$$FD = \frac{1}{2} CD$$

It is given that,

$$AB = CD \Rightarrow \frac{1}{2} AB = \frac{1}{2} CD$$

$$\text{So, } AE = FD \dots (1)$$

Since equal chords are equidistance from the centre,

And  $AB = CD$

$$\text{So, } OE = OF$$

Now, as proved, in right triangles MOE and MOF,

hyp.  $OE =$  hyp.  $OF$  [Common side]

$$OM = OM$$

$$\triangle MOE \cong \triangle MOF$$


$$ME = MF \dots (2)$$

Subtracting equations (2) from (1), we get

$$AE - ME = FD - MF$$

$\Rightarrow AM = MD$  [Proved part (ii)]  
 Again,  $AB = CD$  [Given]  
 And  $AM = MD$  [Proved]  
 $AB - AM = CD - MD$  [Equals subtracted from equal]  
 Hence,  $MB = MC$  [Proved part (i)]

**2. If non-parallel sides of a trapezium are equal, prove that it is cyclic.**  
**Solution:**



According to the question,  
 We have,  
 ABCD is a trapezium in which  $AD \parallel BC$   
 Non-parallel sides AB and DC of the trapezium ABCD are equal i.e.,  
 $AB = DC$ .  
 To prove: Trapezium ABCD is cyclic.  
 Construction: Draw AM and DN such that they are perpendicular on BC.  
 Proof: In right triangles AMB and DNC,  
 $\angle AMB = \angle DNC = 90^\circ$   
 $AB = DC$  [Given]  
 Since perpendicular distance between two parallel lines are same,  
 $AM = DN$   
 $\triangle AMB \cong \triangle DNC$  [By RHS congruence rule]  
 $\angle B = \angle C$  [CPCT]  
 And  $\angle 1 = \angle 2$   
 $\angle BAD = \angle 1 + 90$   
 $= \angle 2 + 90$   
 $= \angle CDA$   
 Now, in quadrilateral ABCD  
 $\angle B + \angle C + \angle CDA + \angle BAD = 360$   
 $\angle B + \angle B + \angle CDA + \angle CDA = 360$   
 $2(\angle B + \angle CDA) = 360$   
 $\angle B + \angle CDA = 180$

We know that,


If any pair of opposite angles of a quadrilateral is  $180^\circ$ , then the quadrilateral is cyclic.

Hence, the trapezium ABCD is cyclic.

**3. If P, Q and R are the mid-points of the sides BC, CA and AB of a triangle and AD is the perpendicular from A on BC, prove that P, Q, R and D are concyclic.**

**Solution:**

To prove: R, D, P and Q are concyclic.



Construction: Join RD, QD, PR and PQ.

RP joins the mid-point of AB, i.e., R, and the mid-point of BC, i.e., P.

Using midpoint theorem,

$RP \parallel AC$

Similarly,

$PQ \parallel AB$ .

So, we get,

AR PQ is a parallelogram.

So,  $\angle RAQ = \angle RPQ$  [Opposite angles of a ||gm]...(1)

ABD is a right angled triangle and DR is a median,

$RA = DR$  and  $\angle 1 = \angle 2$  ...(2)

Similarly  $\angle 3 = \angle 4$  ...(3)

Adding equations (2) and (3),

We get,

$$\angle 1 + \angle 3 = \angle 2 + \angle 4$$

$$\Rightarrow \angle RDQ = \angle RAQ$$

$\angle RPQ$  [Proved above]

Since  $\angle D$  and  $\angle P$  are subtended by RQ on the same side of it, we get the points R, D, P and Q concyclic.

Hence, R, D, P and Q are concyclic.

**4. ABCD is a parallelogram. A circle through A, B is so drawn that it intersects AD at P and BC at Q. Prove that P, Q, C and D are concyclic.**

**Solution:**


According to the question,

ABCD is a parallelogram.

A circle through A, B is so drawn that it intersects AD at P and BC at Q.

To prove: P, Q, C and D are concyclic.

Construction: Join PQ.



Extend side AP of the cyclic quadrilateral APQB to D.

External angle,  $\angle 1$  = interior opposite angle,  $\angle B$

Since,  $BA \parallel CD$  and BC cuts them

$$\angle B + \angle C = 180^\circ$$

Since, Sum of interior angles on the same side of the transversal =  $180^\circ$

$$\text{Or } \angle 1 + \angle C = 180^\circ$$

So, PDCQ is cyclic quadrilateral.


Hence, the points P, Q, C and D are concyclic.

**5. Prove that angle bisector of any angle of a triangle and perpendicular bisector of the opposite side if intersect, they will intersect on the circumcircle of the triangle.**

**Solution:**

According to the question,

Triangle ABC and  $l$  is perpendicular bisector of BC.



To prove:

Angles bisector of  $\angle A$  and perpendicular bisector of BC intersect on the circumcircle of  $\triangle ABC$ .

Proof:

Let the angle bisector of  $\angle A$  intersect circumcircle of  $\triangle ABC$  at D.

Construction: Join BP and CP.

Since, angles in the same segment are equal

We have,  $\angle BAP = \angle BCP$

We know that,

AP is bisector of  $\angle A$ .

Then,

$$\angle BAP = \angle BCP = \frac{1}{2} \angle A \dots (1)$$

Similarly,

We have,

$$\angle PAC = \angle PBC = \frac{1}{2} \angle A \dots (2)$$

From equations (1) and (2),

We have

$$\angle BCP = \angle PBC$$

We know that,

If the angles subtended by two Chords of a circle at the centre are equal, the chords are equal.

So,

$$BP = CP$$

Here, P is on perpendicular bisector of BC.

Hence, angle bisector of  $\angle A$  and perpendicular bisector of BC intersect on the circumcircle of  $\triangle ABC$ .