ANATOMY

1. **concerning ureter which is true?**
    a) has constriction when enter pelvic prim
    b) mainly supply by left renal artery
    c) inferior epigastric vein run on lateral side
    d) run in direct contact with the tip of transverse process of the lumber vertebra
2. **female 65 years diagnosed with pelvic tumor metastasis to medial inguinal lymph nodes what is the origin?**
    a) urinary bladder
    b) ovaries
    c) uterus
    d) uterine tube
3. **24 years got trauma to pelvis and penis come with swollen in scrotum the fluid extravasated to:**
    a) deep perineal pouch
    b) tunica vaginalis
    c) douglas pouch
    d) superficial perineal pouch
4. **small (short) saphenous vein which is correct?**
    a) ascend up on medial malleolus
    b) always has valve in terminal end (12 valves)
    c) has relation with lateral sural nerve
    d) perforate deep fascia in upper part of popliteal fossa
5. **In exploration popliteal artery?**
    a) run anterior to sural nerve
    b) run superior to popliteal vein
    c) anterior tibial artery is Infront of popliteal muscle in high division
    d) go round and flat to words knee to be ant tibial artery
6. **Lymph node of trigone of bladder drain to?**
    a) internal iliac lymph nodes

b) external iliac lymph nodes
c) Internal and external iliac lymph nodes
d) Superficial inguinal lymph nodes
e) D-paraaortic lymph nodes

7. Which of the following concerning the penis is incorrect?

a) glans of penis form bulbospongiosus
b) ischiocovernosus cover the pulp of penis
c) penile urethra lies in bulbospongiosus
d) labia majora is bounded the vestibule

8. Which of the following concerning the perineum is incorrect?

a) somatic sensory of clitoris and glans of penis come from internal pudendal nerve
b) erection of penis and clitoris by parasympathetic
c) contraction of pulbospongiosus and covernousus muscles by parasympathetic
d) bulpspongiousus cover RT and LEFT crura of clitoris

9. Internal spermatic fascia made from?

a) external oblique muscle
b) internal oblique muscle
c) cremasteric facia
d) transversalis fascia

10. Concerning surgery the sciatic nerve?

a) come anterior to piriformis
b) has relation with superior gluteal artery
c) save to deal with lateral site in gluteus

11. Regarding lymph draining to lower limbs all true except?

a) medial group of inguinal lymph nodes is much in number than lateral group
b) glans of penis drains to deep inguinal lymph nodes
c) deep inguinal lymph nodes are fewer in number
d) popliteal glands drain entirely to the deep group of inguinal lymph nodes

12. Which of the following ligament attach to uterus and have peritoneum?

a) cardinal ligament
b) broad ligament
c) round ligament
d) suspensory ligament

13. Following statement is incorrect about perineum structure:

a. anorectal ring formed by subcutaneous, superficial and 2deep part of external anal sphincter

b. perineal membrane attaches laterally to pubic arch and ischial spine.

14. cancer from pelvic viscera can deposits cancerous cells in the vertebrae, this communication occurs between the internal iliac veins and vertebral veins through:

a) lateral sacral veins
b) lumbar vein
c) umbilical veins
d) prostatic plexus

15. Concerning pelvic which is true?

a) Femoral ring is 1.25-inch lateral and below the pubic tubercle
b) The anterior superior iliac spine is at spinous level of second sacral vertebra
c) The greater sciatic notch lies at the level of third sacral vertebra
d) The line that meets the anterior superior iliac spine with most prominent point in ischial tuberosity passes directly through the acetabulum

16. Primary support of uterus by:

a) Lateral transverse cervical ligament (cardinal ligament)
b) Round ligament
c) Pubocervial ligament
d) broad ligament

17. Ischiorectal fossa is clinically important space. Which of the Following is site of their connection:

a) anteriorly
b) posteriorly

c) Inferiorly
d) Anterolaterally

18. posterior fornix is a gate way to:

a) Retroperitoneal space
b) Vesciouterine pouch
c) Retcouterine pouch (douglus pouch)

19. 60-year-old female known case of ovarian CA presented with Medial thigh Parathesis the nerve involved by CA is

a) obturator N
b) medial cutaneous N of the thigh
c) intermediate cutaneous N of the thigh
d) femoral trunk
e) lumbosacral trunk

20. Hip joint characterizes by huge capsule. Which of the following is Attached to it posteriorly:

a) greater trochanter
b) lesser trochanter
c) Inter trochanteric line
d) Mid cervical line (Femoral neck)
e) intertubercular line

21. 2 years old boy has hernia sac that lying above and medial to the Pubic tubercle. Which of the following is diagnosis:

a) Femoral hernia
b) Umbilical hernia
c) Post surgical hernia
d) Direct inguinal hernia
e) Indirect inguinal hernia

22. In a patient with small volume of fluid collection in the abdomen Likely site for fluid collection in the erect position is:

a) Lesser sac
b) Greater sac
c) Douglas pouch
d) Rt hepatorenal pouch

e) Paraduedenal recesses

23. Femoral pulsation is detected best at which point:

a) mid inguinal point
b) 2/3 away from the pubic tubercle.
c) Inferior and lateral to pubic tubercle
d) Lateral to femoral nerve
e) Medial and above pubic tubercle

24. Which of the following Muscles not attached to perineal body:

a) Levator ani
b) Puborectalis
c) Pubovaginalis
d) Superficial Transverse perineal muscle
e) External anal sphincter

25. Epiploic foramen bounded anteriorly by:

a) Inferior vena cava
b) Caudate lobe
c) First part duodenum
d) portal vein
e) stomach

26. Concerning urinary bladder

a) Against fascia of levator ani
b) Detrusor muscle supplied by sympathetic motor fiber
c) Connected to remnant of cloaca
d) In female is completely pelvic organ
e) Its sphincter vesicae has parasympathetic motor supply

27. Regarding the anterior abdominal wall:

a) Superficial tissues of the anterolateral wall above the umbilicus to axillary nodes
b) Intertubercular plane lies at the level of L4

c) Mid-inguinal point lying halfway between the pubic tubercle and Anterior Superior Iliac Spine
d) Posterior wall of the inguinal canal is formed by the transversalis abdomius muscle
e) Surgical incision at Linea alba heals faster than that along paramedian abdominal wall

28. Most precancerous organ and can predictable by screening?

a) Ovary
b) Vulva
c) Cervix.
d) Endometrium

29. Anal canal above pectoral line

a) innervated by inferior pudendal nerve
b) contain columns

30. Muscle of urogenital diaphragm include

a) Sphincter urethrae
b) Sphincter vesicae
c) Levator ani
d) Ischiocavernosus
e) Obturator internus

31. lymph node of lower part of uterine body is

a) Para aortic lymph node
b) Internal iliac lymph node
c) External iliac lymph node
d) Deep group

32. What is the glistening membrane that covers the fetal surface of the placenta?

a) Chorion
b) Decidua
c) Amnion
d) Cotyledon

33. 19 years old college student, majoring in biology, just delivered her first baby, she is very interested in studying early pregnancy and asks about the origin of the placental tissue. which of the following develop into the placenta?

a) Trophectoderm
b) Inner cell mass of the blastocyst
c) Decidua
d) Myoblast cells.

34. Inguinal canal has four wall which of the following from the posterior wall:

a) Internal oblique
b) External oblique
c) Transvers abdominus
d) Transversalis fascia

35. Iliac crest at level of

a) L3
b) L4
c) L5
d) S1

36. The dermatomes of umbilical region?

a) T7
b) T10
c) T12
d) L1

37. The superficial inguinal ring from

a) Transverses abdominus
b) 3Aponeurosis external oblique
c) Aponeurosis internal oblique
d) conjoint tendon

38. Main muscle responsible for abduction of hip joint

a) Iliopoas muscle
b) Gluteus medius
c) Gluteus minimus

39. Pudendal nerve before enters the pudendal canal lie on which of the following structure

a) Piriformis
b) Obturator internus muscle and fascia
c) Sacrospinatous ligament
d) Sacrotuberous ligament

40. Which of the following operations involve in cutting rectus sheath anteriorly and posteriorly

a) paramedian
b) Subcostal
c) Transverse suprapubic
d) Vertical incision

41. Medial part of the knee joint supply by

a) Femoral nerve
b) Sciatic nerve
c) tibial nerve
d) Obturator nerve

42. Which of the following nerve supply the anterior 2/3 of labia majora and clitoris

a) Ilioinguinal nerve
b) Genitofemoral nerve
c) perineal branch of pudendal nerve
d) lateral sacral nerve

43. which of the following structure lies medial to femoral artery within femoral canal

a) femoral nerve
b) femoral vein
c) femoral sheath
d) lymph nodes

44. Vagina covering by peritoneum in

a) anterior wall
b) posterior fornix
c) anterolateral wall
d) lateral fornix

45. Which of the following is non glandular organ

a) Uterus
b) Vagina
c) Fallopian tube
d) Cervix

46. Zona pellucida disintegrates just:

a) before fertilization
b) after fertilization
c) midway during cleavage
d) after completion of cleavage

47. Regarding the Ureter:

a) partially retroperitoneal
b) received blood supply by ovarian artery
c) on the right cross ovarian vessels
d) on the left cross inferior vena cava

48. mature placenta composed of:

a) syncytiotrophoblast, cytotrophoblast
b) maternal blood vessels, syncytiotrophoblast, fetal blood vessels
c) syncytiotrophoblast, fetal blood vessels
d) syncytiotrophoblast, cytotrophoblast, mesenchyma

49. regarding Rectus sheath:

a) inferior and superior epigastric arteries anastomosis
b) below umbilicus 3 layers of muscles
c) below umbilicus 2 layers of muscles d.

50. external genitalia development:

a) genital tubercle give rise to clitoris
b) clitoris between two labia majora
c) genital folds become labia majora
d) glans derived from genital swelling

51. Theca cell produces primarily:

a) progesterone

b) androsterone
c) estrogen
d) DHEAS

52. All of the following about the anterior abdominal wall is correct except:

a) Median umbilical ligament represented by urachus
b) ligamentum of teres remnant of umbilical vein
c) pyramidalis m. presented in 20% of population
d) Medial umbilical ligament represented by umbilical arteries

53. Anterior division of internal iliac artery:

a) lateral sacral artery
b) iliolumbar artery
c) obturator artery
d) inferior rectal artery

54. Regarding the Fallopian tube:

a) it contains inner longitudinal muscles that moves the fertilized ovum into the endometrium
b) isthmus run along uterine wall
c) it's diameter about 0.5 to 1.0 cm d

55. About spermatogenesis:

a) nucleus form the cap
b) mitochondria make the neck
c) the centriole elongated to form axial filament
d) tail have two and nine microtubules

56. Regarding female development:

a. vaginal plate forms the lower two third of vagina which connects to para mesonephric duct

57. Ovarian artery:

a) Rt ovarian artery crosses anterior to IVC
b) originating from the aorta above the renal a.
c) enter ovary through ovarian ligament d.

58. At what stage of the following life is compatible Extra uterine:

a) glandular
b) canalicular
c) saccular
d) after birth

59. Regarding early stages of development:

a) zygote contain blastomere
b) blastocyte contain morula
c) implantation occur after 10 days
d) inner cell mass becomes placenta

60. Extrusion of Second polar body from egg occurs:

a) after entry of sperm and before completion of fertilization
b) after completion of fertilization
c) before the entry of sperm
d) without any relation with sperm entry

61. Innervation of majority of Vulva?

a) Ilioinguinal nerve
b) Pudendal nerve
c) Inferior rectal nerve
d) Perineal branch of S4

62. Approximately how many oocytes are present in the ovaries at birth?

a) 1,000
b) 10,000
c) 100,000
d) 1 million

63. What process would you expect a secondary spermatocyte to undergo during spermatogenesis?

a) 1st meiotic division
b) 2nd meiotic division
c) 1st mitotic division
d) differentiation

64. Which of the following would be most likely to be damaged by a stab wound into the ischiorectal (ischioanal) fossa 2 cm lateral to the anal canal?

a) Crus of the Penis
b) Perineal Body
c) Pudendal Nerve
d) Inferior Rectal Artery

65. Regarding the anterior abdominal wall:

a) the arcuate lines lie midway between the symphysis pubis and umbilicus
b) between the umbilicus and the lateral margin transversus aponeurosis lies posterior the rectus
c) the inferior epigastric artery originates from the internal iliac
d) the ilioinguinal nerve passes through the deep inguinal ring

66. Pelvic joints and ligaments:

a) the sacroiliac joint is a fibrous joint between the auricular surfaces of the ilium and sacrum
b) the wedge shape of the sacrum contributes to its stability
c) the sacrotuberous and sacrospinous ligaments enclose the greater sciatic foramen
d) the iliolumbar ligament is V-shaped with the apex attached to L5

67. Within the anal canal are anal cushions (dilated venous spaces and anastomoses)

a) 3, 7 and 11 o'clock
b) 2, 6, and 10 o'clock
c) 1, 7 and 12 o'clock
d) 1, 7 and 9 o'clock

68. Referred pain from which organ may be felt in the cutaneous distribution of the obturator nerve?

a) bladder
b) prostate
c) ovary
d) uterus

69. The pelvic floor

a) Muscles include obturator internus
b) Slopes down and forwards

c) Is supplied by S2, S3 and S4
d) Muscle iliococcygeus arises from the ilium

70. Perineum:

a) lymphatic drainage from the rectum passes to internal iliac nodes from the upper part and to superficial
b) superior rectal branches of the pudendal nerves supply the external and sphincter
c) the internal anal sphincter is composed of skeletal muscle
d) the anal canal consists of inner circular muscle fibers and outer longitudinal fibers

71. Regarding the inguinal canal:

a) the deep inguinal ring lies 1cm above and at the junction of the lateral third and medial two thirds of the inguinal ligament
b) the deep inguinal ring is formed by an opening in the transversalis fascia
c) an indirect inguinal hernia bulges into the inguinal triangle, posterior to the canal and medial to the inferior epigastric artery
d) contains the spermatic cord that has two fascial layers/coverings and eight contents

72. Which pair of structures does NOT differentiate from comparable embryonic structures in the male and female?

a) Bulb of corpus spongiosum and vestibular bulb
b) Shaft of penis and labia majora
c) Glans of penis and glans of clitoris
d) Crus of corpus cavernosum penis and crus of corpus cavernosum clitoris

73. The most inferior extent of the peritoneal cavity in the female is the:

a) Pararectal fossa
b) Paravesical fossa
c) Rectouterine pouch
d) Rectovesical pouch

74. Which of the following structures contribute to both of the pelvic inlet and outlet?

a) Pubic symphysis
b) Iliopectineal line
c) Ischiopubic ramus

d) Sacral promontory

75. All of the following are parts of the lateral pelvic wall except:

a) Piriformis muscle
b) Obturator internus
c) Sacro tuberous ligament
d) Sacrospinous ligament

76. pain from vagina is lie on:

a) T10 - L1
b) T12 - L2
c) T12 - L3
d) S2 - S4

77. Cancer from ovary Mets to:

a) omentum
b) liver
c) lung
d) Stomach

78. Content of broad ligament include:

a) Ureter
b) Round ligament of the uterus
c) suspensory ligament of the ovary
d) Ovarian ligament

79. Which nerve provide sensation the skin over the suprapupic area:

a) superior hypogastric plexus
b) illiohypogastric plexus
c) pudendal nerve
d) femoral nerve

80. A female comes after normal vaginal delivery complaining of dribbling of urine with sneezing, coughing and straining Which structure is most likely to be damaged?

a) levator ani muscle
b) Transverse muscle
c) Urethral sphincter
d) perineal body

81. Inguinal canal has four wall which of the following from the posterior wall:

a) Internal oblique

b) External oblique
c) Transverse abdominis
d) conjoint tendon
e) lacunar ligament

82. Regarding The anterior abdominal wall:

a) the deep surface above the umbilicus has lymphatic drainage to mediastinal nodes
b) the deep surface below the umbilicus has lymphatic drainage to the internal iliac and paraaortic nodes
c) rectus abdominis is supplied by lower intercostal and subcostal n. (T8-T10)
d) internal oblique and transversus abdominis are supplied solely by the iliohypogastric and ilioinguinal nerves
e) the superior and inferior epigastric vessels (arteries) anastomose within the body of the rectus muscle

83. Genitofemoral nerve is:

a) Branch from femoral
b) Pass in deep inguinal canal
c) Pass anterior to iliacus
d) Supply sensation of medial portion of thigh

84. Obstetrician with anaesthetist perform episiotomy. block pudendal nerve at sacrospinous ligament, has he blocked cutaneous or dermatome field?

a) Cutaneous field because it arises from single spinal level
b) Dermatomal because it arises from single spinal level
c) Cutaneous field because it arises from many spinal level
d) Dermatome because it arises from many spinal level

85. Superior gluteal artery and inferior gluteal artery named for their relation to:

a) Gluteus maximus
b) Gluteus Medius
c) Sciatic nerve
d) Piriformis muscle

86. Trauma by bicycle injured perineal body and structure attached to it, which structure will be spared:

a) levator ani
b) external anal sphincter
c) coccygeus
d) Bulbospongiosus muscle
e) Perineal membrane

87. Infection and bleeding below umbilicus in deep peritoneum spread to perineum by:

a) Scarpa s fascia
b) Camper fascia
c) Parietal peritoneum
d) Deep fascia

88. The ureter passes deep to the:

a) femoral nerve
b) gonadal artery
c) psoas muscle
d) genitofemoral nerve
e) common iliac artery

89. In differentiation between male and female what is true:

a) Lymph drainage from the gonads can reach the para-aortic lymph node in both
b) in male the cavities open outside where in female it's not
c) ureter has the same path across the pelvic in both

90. Lymph drainage of the lower anal canal

a) Superficial inguinal LN
b) Internal iliac LN
c) External iliac LN
d) Deep inguinal LN

91. Regarding Lower limb veins:

a) dermatitis characteristic in Buerger disease
b) thrombosis more in superficial veins
c) Superficial veins is source of pulmonary thrombus
d) 20% of people have thrombosis