PHARMACOLOGY

1. **reversible inhibition of cyclooxygenase in platelet is caused by?**
    a) alprostadil
    b) aspirin
    c) paracetamol
    d) ibuprofen
2. **Cardiotoxicity limits the clinical usefulness of which one of the following antitumor antibiotics:**
    a) Doxorubicin
    b) Bleomycin
    c) Dactinomycin
    d) Plicamycin
3. **A nucleophilic attack on deoxyribonucleic acid which cause disruption of base pairing occurs as a result:**
    a) Cyclophosphamide
    b) Thioguanine
    c) Prednisone
    d) Methotrexate
4. **19 yrs female diagnosed with TB. before prescribing treatment, you should be aware of that one antitubercular induce cytochrome P450 enzyme in the liver:**
    a) rifampicin
    b) isoniazid
    c) pyrazinamide
    d) ethambutol
5. **Which of the following antiviral agents is a purine nucleoside analogue that has shown promise with Lassa fever, influenza A and B, and respiratory syncytial virus (RSV)?**
    a) Amantadine
    b) Rimantadine
    c) Ribavirin
    d) Acyclovir
6. **drug use to treat angina consistently cause reduction of HR and prevent Vaso spasmodic angina?**
    a) nitro glycerate

b) propranolol
c) verapamil
d) isosorbide dinitrate

7. Coombs test is positive and associated with hemolytic anemia in?

a) methyldopa
b) captopril
c) clonidine

8. atropine overdose led to:

a) bronchoconstriction
b) increase HR
c) increased gastric secretion
d) pupillary constriction

9. 45 years old treated for ovarian cancer developed difficulty hearing:

a) paclitaxel
b) bleomycin
c) cisplatin

10. benzodiazepine action:

a) enhance GABA B neurotransmitters
b) enhance GABA A in spinal cord
c) enhanced Cl channels that act on GABA neurotransmitters

11. amiloride potent diuretic action by:

a) aldosterone antagonist
b) acts on ENaC
c) Na -K blocker

12. drug to use in lowering blood pressure, analgesic and spasmolytic and given intrathecally:

a) timolol
b) clonidine
c) morphine
d) nitroprusside

13. primary mechanism of furosemide:

a) inhibition of sodium potassium ATPase
b) inhibition of Na k cl co- transporter
c) inhibition of Na Cl co-transporter

d) inhibition of Cl transporter

14. anaphylactic shock treatment is by

a) epinephrine
b) chlorpheniramine maleate
c) atropine
d) hydrocortisone

15. A patient with chronic renal failure receiving an intravenous Diamorphine infusion has had a respiratory arrest. Which of the Following is the most likely cause?

a) Decreased plasma clearance of diamorphine
b) Decreased plasma volume following dialysis leading to increased plasma concentration of diamorphine
c) Impaired renal excretion of diamorphine
d) Increased plasma concentration of 6-monoacetyl morphine
e) Increased plasma concentration of morphine-6- glucuronide

16. An experiment is being designed to assess speed of gastric Emptying. It is planned to administer a standard dose of a marker rug orally to healthy volunteers and then measure plasma drug Concentrations at standard times after the oral dose. Which of the Following drugs is likely to be most useful in this experiment?

a) Aspirin
b) Morphine
c) Paracetamol
d) Propranolol
e) Vancomycin

17. An intravenous infusion of a drug is administered to a 60 kg Subject and, at steady state, urine is collected over four hours from a Urinary catheter. The following data are obtained:

- **Plasma concentration at steady state = 25 mcg ml**
- **Urinary concentration = 2.5 mg ml**
- **Total urinary volume = 240 ml**

What is the renal clearance of the substance?

a) 24 ml min
b) 60 ml min

c) 100 ml min
d) 120 ml min
e) 240 ml min

18. Drug of choice for dermal, oral, vaginal candidiasis is

a) Griseofulvin
b) Amphotericin B
c) Gentian violet
d) Nystatin

19. B- Lactam ring is present in

a) Erythromycin
b) Penicillin
c) Tetracycline
d) Chloramphenicol

20. Erythromycin is obtained from

a) s. griseus
b) s. rimosus
c) s. scabies
d) s. erythraeus.

21. Mode of action of quinolone antibiotics (ciprofloxacin) on growing bacteria was thought to be

a) Inhibition of lactamase
b) Prevention of cross linking of glycine
c) Inhibition of DNA gyrase
d) Inhibition of reverse transcriptase

22. Drug of choice for methicillin resistant staph. Aureus is

a) Ampicillin
b) Erythromycin
c) Neomycin
d) Vancomycin

23. The anti-tumor antibiotic act by inhibiting

a) Cell wall synthesis
b) RNA synthesis
c) Cell membrane synthesis
d) The DNA structure and function

24. Which of the following does not inhibit cell wall synthesis?

a) Penicillin
b) Carbenicillin
c) Amikacin
d) Vancomycin

25. Which of the following is penicillinase resistant, acid labile penicillin?

a) Amoxycillin
b) Cloxacillin
c) Carbenicillin
d) Methicillin

26. Female complain of dysmenorrhea you give her indomethacin, İndomethacin inhibit the formation of:

a) Bradykinin
b) histamine
c) PG F2
d) PG E2

27. Which of the following medications is recommended most commonly for the treatment of maternal hyperthyroidism in pregnancy?

a) Carbimazole
b) Propylthiouracil
c) Levothyroxine
d) Radioactive iodine

28. Which of the following drugs is a known fetal teratogen?

a) Marijuana
b) Alcohol
c) Heroin
d) Caffeine

29. Which of the following agents is a known teratogen in humans?

a) Vitamin A
b) Vancomycin
c) Oral contraceptives
d) Ampicillin

30. Which of the following medications is not associated with central nervous system (CNS) abnormalities?

a) Valproic acid
b) Methotrexate
c) Tegretol

d) Vitamin A
e) Thalidomide

31. Pt after injected spinal anaesthesia became hypotensive, first treatment is:

a) Sympathomimetic infusion
b) Blood transfusion
c) Plasma
d) Fluids